

CHROMOSOMAL ABNORMALITIES ASSOCIATED WITH NEURAL TUBE DEFECTS (II): PARTIAL ANEUPLOIDY

Chih-Ping Chen*

Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Department of Medical Research, Mackay Memorial Hospital, Taipei, Department of Biotechnology, Asia University, College of Chinese Medicine, China Medical University, Taichung, and Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan.

SUMMARY

Fetuses with neural tube defects (NTDs) carry a risk of chromosomal abnormalities. The risk varies with maternal age, gestational age at diagnosis, association with other structural abnormalities, and family history of chromosome aberrations. This article provides a comprehensive review of structural chromosomal abnormalities associated with NTDs, such as del(13q), r(13), dup(2p), del(2q), del(1p), del(1q), dup(1q), del(3p), dup(3p), del(3q), dup(3q), del(4p), dup(4p), del(4q), dup(4q), del(5p), del(6p), dup(6q), del(6q), dup(7p), del(7q), dup(8q), del(9p), del(10q), del(11q), dup(11q), dup(12p), dup(14q), del(14q), del(15q), dup(16q), del(18q), r(18), dup(20p), +i(20p), del(22q), del(Xp), and dup(Xq). NTDs may be associated with aneuploidy. Perinatal identification of NTDs should alert one to the possibility of chromosomal abnormalities and prompt a thorough cytogenetic investigation and genetic counseling. [*Taiwan J Obstet Gynecol* 2007;46(4):336-351]

Key Words: chromosomal abnormalities, neural tube defects, partial aneuploidy

Introduction

Neural tube defects (NTDs) have an incidence of 1–2 per 1,000 births and are considered to be a heterogeneous condition resulting from failure of normal neural tube closure between the third and fourth week of embryonic development. The three common types of NTDs are anencephaly, spina bifida, and encephalocele. The uncommon types of NTDs include amniotic band syndrome, limb-body wall complex, cloacal exstrophy or OEIS complex, and other types of spinal abnormalities. The incidence of NTDs varies with race, geographic variation, socioeconomic classes, nutritional status, and multiple predisposing factors such as single gene disorders (Meckel syndrome, median cleft face syndrome, Roberts syndrome, Jarcho-Levin syndrome and HARD [hydrocephalus, agyria, retinal dysplasia] syndrome),

chromosomal abnormalities (trisomy 18, trisomy 13, triploidy and other structural abnormalities), teratogens (valproic acid, aminopterin/amethopterin and thalidomide), maternal diabetes, family history of NTDs, thermolabile mutation in the *MTHFR* gene, and others [1]. There is considerable evidence that genetics contributes to the etiology of NTDs. Fetuses with NTDs carry a risk of chromosomal abnormalities. The risk varies with maternal age, gestational age at diagnosis, association with other structural abnormalities, and family history of chromosome aberrations.

Partial Aneuploidy

Structural chromosomal abnormalities associated with NTDs include del(13q), r(13), dup(2p), del(2q), del(1p), del(1q), dup(1q), del(3p), dup(3p), del(3q), dup(3q), del(4p), dup(4p), del(4q), dup(4q), del(5p), del(6p), dup(6q), del(6q), dup(7p), del(7q), dup(8q), del(9p), del(10q), del(11q), dup(11q), dup(12p), dup(14q), del(14q), del(15q), dup(16q), del(18q), r(18), dup(20p), +i(20p), del(22q), del(Xp), and dup(Xq).

*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei 104, Taiwan.
E-mail: cpc_mmh@yahoo.com
Accepted: October 11, 2007

Del(13q) and r(13)

The reported NTDs associated with del(13q) and r(13) include spina bifida, encephalocele, anencephaly, atelencephaly, exencephaly, and aprosencephaly [2–20] (Table 1). Schmid et al [3] first reported the prenatal diagnosis of ring chromosome 13 by amniocentesis at 33 weeks' gestation in a fetus with anencephaly, limb deformities, and congenital heart defects. Rudelli et al [8] reported the pathologic characteristics of two 18-week gestation fetuses with anencephaly, exencephaly, multiple limb defects, and urogenital defects carrying chromosome 13q deletions (distal to bands q13 and q22, respectively). Benn et al [7] reported the prenatal diagnosis of ring chromosome 13 (46,XY,r(13)(p11q13)) by amniocentesis at 20 weeks' gestation in a fetus with anencephaly, imperforate anus and urethral meatus, severe talipes, syndactyly, cardiac defects, and other anomalies. Palmer et al [9] reported the prenatal diagnosis of ring chromosome 13 (46,XX,r(13)) by

amniocentesis at 18 weeks' gestation in a fetus with anencephaly. Brown et al [13] reported prenatal diagnosis of chromosome 13q deletion (46,XX,del(13)(q31.2q34)) by amniocentesis at 25 weeks' gestation in a fetus with a large occipital encephalocele, severe microcephaly, anteriorly displaced anus, and bilaterally absent thumbs. Chen et al [16] reported the prenatal diagnosis of a chromosome 13q deletion (46,XX,del(13)(q21)) by amniocentesis at 20 weeks' gestation in a fetus with severe intrauterine growth restriction, cardiomegaly, an occipital encephalocele, and a calvarial defect. Chen et al [17] reported the prenatal diagnosis of mosaic ring chromosome 13 (46,XX,r(13)(p11q32)/45,XX,-r(13); 77%/23%) by amniocentesis at 23 weeks' gestation in a fetus with intrauterine growth restriction and anencephaly. Molecular genetic analysis of the case reported by Chen et al [17] revealed that the breakpoint was distal to the *ZIC2* gene on 13q32, suggesting the association of other genes on 13q32–q34 with

Table 1. Reported neural tube defect (NTD) cases with partial aneuploidy of chromosome 13

Authors	Chromosome aberration	Deletion or duplication extent	NTDs and brain anomalies
Del(13q) and r(13)			
Fear and Briggs [4]	der(13)t(3;13)(q21;q34)mat	13q34 → qter 3q21 → qter	Lumbar meningocele
Kennedy et al [18]	der(13)t(11;13)(q23.1;q34)mat	13q34 → qter 11q23.1 → qter	Lumbosacral meningocele
Jones et al [6]	r(13)(p11q34)/-r(13) (84.5%/6.5%)	13q34 → qter	Lumbar spina bifida
Luo et al [20]	del(13)(q33.2)	13q33.2 → qter	Lumbosacral meningocele
Chen et al [17]	r(13)(p11q32)/-r(13) (77%/23%)	13q32 → qter	Anencephaly
Telfer et al [5]	del(13)(q32)	13q32 → qter	Parietal encephalocele
Brown et al [14]	der(13)t(13;22)(q31.2;q13.3)pat	13q31.2 → qter 22q13.3 → qte	Partial anencephaly, posterior encephalocele
Kennedy et al [18]	der(13)t(13;22)(q31.2;q13.3)pat	13q31.2 → qter 22q13.3 → qter	Anencephaly
Brown et al [13]	del(13)(q31.2q34)	13q31.2 → qter	Occipital encephalocele
Brown et al [13]	der(13)t(13;15)(q22;q26)pat	13q22 → qter 15q26 → qter	Anencephaly
Rudelli et al [8]	del(13)(q22)	13q22 → qter	Anencephaly
Towfighi et al [10]	del(13)(q22q31)	13q22 → q31	Atelencephaly
Lam et al [19]	der(13)t(4;13)(p15.2;q21.2)pat	13q21.2 → qter 4p15.2 → pter	Exencephaly
Brandt et al [12]	r(13)(p11q21.1)/-r(13) (92%/8%)	13q21.1 → qter	Anencephaly
Chen et al [16]	del(13)(q21)	13q21 → qter	Occipital encephalocele
Drugan et al [11]	del(13)(q13)	13q13 → qter	Anencephaly
Benn et al [7]	r(13)(p11q13)	13q13 → qter	Anencephaly
Rudelli et al [8]	del(13)(q13)	13q13 → qter	Anencephaly
Goldsmith et al [15]	r(13)/-r(13) (67%/37%)	Unknown	Aprosencephaly
Palmer et al [9]	r(13)	Unknown	Anencephaly
Schmid et al [3]	r(13)/-r(13) (93%/7%)	Unknown	Anencephaly
Allderdice et al [2]	r(13)	Unknown	Encephalocele

Del = deletion; *r* = ring chromosome; *der* = derivative chromosome; *mat* = maternal origin; *-r(13)* = monosomy 13 with absence of *r(13)*; *pat* = paternal origin.

NTDs. The *ZIC2* gene is well known to be responsible for human holoprosencephaly (HPE) [21]. Recent studies, however, found no supporting evidence for an association between *ZIC2* and the risks of NTDs [22,23]. The human *ZIC5* on 13q32 has been proposed as a strong candidate gene responsible for NTDs [24]. Inoue et al [24] found that mice with disrupted *ZIC5* manifested incomplete neural tube closure and suggested that *ZIC5* is involved in the generation of the neural tube tissue in mouse development. Other possible candidate genes of NTDs in the involved segment of chromosome 13q include *MAB21L1* (13q13). *Cdx2* maps to 13q12.3 and is essential for axial elongation in mouse development [25].

Dup(2p)

The reported central nervous system anomalies associated with *dup(2p)* include spina bifida, anencephaly, encephalocele, atelencephaly, and agenesis of corpus callosum [26–38] (Table 2). Walbaum et al [31] reported a fetus with anencephaly and truncus arteriosus as the unbalanced product of a translocation carrier mother with *t(2;5)(p13;p15)* and a previous child with spina bifida. The fetus had *dup(2)(pter → p13)* and *del(5)(pter → p15)*. Singer et al [32] reported the prenatal diagnosis of a duplication of 2p (*2p24 → pter*) by amniocentesis in the third pregnancy of a woman with fetal anencephaly and congenital cystic disease of the right kidney. The maternal karyotype was

Table 2. Reported neural tube defect (NTD) cases with partial aneuploidy of chromosome 2

Authors	Chromosome aberration	Duplication or deletion extent	NTDs
Dup(2p)			
Singer et al [32]	<i>der(10)t(2;10)(p24;q26)mat</i>	<i>2p24 → pter</i> <i>10q26 → qter</i>	Anencephaly
Wellesley and Boyle [36]	<i>der(3)t(2;3)(p23.1;q29)pat</i>	<i>2p23.1 → pter</i> <i>3q29 → qter</i>	Sacral meningomyelocele
Carriere [27]	<i>der(10)t(2;10)(p23;q26)mat*</i>	<i>2p23 → pter</i> <i>10q26 → qter</i>	Encephalocele
Francke and Jones [28]	<i>der(7)t(2;7)(p23;q36)mat*</i>	<i>2p23 → pter</i> <i>7q36 → qter</i>	Anencephaly
Cassidy et al [29]	<i>der(3)t(2;3)(p23;p27)*</i>	<i>2p23 → pter</i> <i>3p27 → pter</i>	Anencephaly
Winsor et al [34] two cases	<i>der(5)t(2;5)(p23;p15)mat</i>	<i>2p23 → pter</i> <i>5p15 → pter</i>	Anencephaly
Thangavelu et al [38]	<i>dup(2)(p25.3p22)</i> <i>dup(2)(p25p22)</i>	<i>2p22 → p25.3</i> <i>2p22 → p25</i>	Anencephaly Anencephaly
Doray et al [37]	<i>der(15)t(2;15)(p22;q26)mat</i>	<i>2p22 → pter</i> <i>15q26 → qter</i>	Craniorachischisis
	<i>der(15)t(2;15)(p22;q26)mat</i>	<i>2p22 → pter</i> <i>15q26 → qter</i>	Anencephaly
Hahm et al [35]	<i>der(15)t(2;15)(p21;q26)</i>	<i>2p21 → pter</i> <i>15q26 → qter</i>	Anencephaly
Therkelsen et al [26] two cases	<i>ins(2)(q34p13p24)pat</i>	<i>2p13 → p24</i>	Spina bifida
Fineman et al [30]	<i>der(9)t(2;9)(p13;p24)mat</i>	<i>2p13 → pter</i> <i>9p24 → pter</i>	Spina bifida
Walbaum et al [31]	<i>der(5)t(2;5)(p13;p15)mat</i>	<i>2p13 → pter</i> <i>5p15 → pter</i>	Anencephaly
Sarda et al [33]	<i>der(X)t(X;2)(q27;p13)mat</i>	<i>2p13 → pter</i> <i>Xq27 → qter</i>	Lumbar meningomyelocele
Del(2q)			
Melnyk and Muraskas [43]	<i>del(2)(q36q36)</i>	<i>2q36</i>	Lumbar meningomyelocele
Nye et al [45]	<i>del(2)(q35q36.2)</i>	<i>2q35 → q36.2</i>	Lumbosacral meningomyelocele
Chinen et al [44]	<i>del(2)(q24.2q31)</i>	<i>2q24.2 → q31</i>	Occipital encephalocele
McConnell et al [42]	<i>del(2)(q22q31)</i>	<i>2q22 → q31</i>	Occipital encephalocele

*Presumed rearrangement. *Dup* = duplication; *der* = derivative chromosome; *mat* = maternal origin; *pat* = paternal origin; *ins* = insertion; *del* = deletion.

46,XX,t(2;10)(p24;q26) and the fetal karyotype was 46,XY,der(10)t(2;10)(p24;q26)mat. The woman's first pregnancy resulted in a female fetus with anencephaly and left hydronephrosis. Pedigree analysis revealed that the woman had a maternal aunt with spina bifida. Hahm et al [35] reported the prenatal diagnosis of partial trisomy 2p (2p21 → pter) and partial monosomy 15q (15q26 → qter) by amniocentesis at 19 weeks' gestation in a fetus with minor facial anomalies, musculoskeletal defects, postaxial polydactyly, and anencephaly. The fetal karyotype was 46,XY,der(15)t(2;15)(p21;q26). Winsor et al [34] reported recurrent anencephaly with partial trisomy 2p (2p23 → pter). The maternal karyotype was 46,XX,t(2;5)(p23;p15). Her first pregnancy resulted in a male fetus with anencephaly, lumbosacral dysraphism at L3-S1, left microphthalmia, preauricular skin tags, absent pituitary gland and left adrenal gland, rocker-bottom feet, bilateral simian creases, hypoplastic penis and scrotum, undescended testes, and the karyotype of 46,XY,der(5)t(2;5)(p23;p15)mat. Her third pregnancy resulted in a fetus with anencephaly, polydactyly, and the karyotype of 46,XY,der(5)t(2;5)(p23;p15)mat. Her fifth pregnancy resulted in a fetus with the karyotype of 46,XY,der(5)t(2;5)(p23;p15)mat, and multiple anomalies but no anencephaly or gross central nervous system abnormalities. Wellesley and Boyle [36] reported the prenatal diagnosis of partial trisomy 2p (2p23.1 → pter) in a fetus with anencephaly and sacral meningocele by amniocentesis at 15 weeks' gestation. The fetal karyotype was 46,XX,der(3)t(2;3)(p23.1;q29)pat. Thangavelu et al [38] reported dup(2)(p25.3p22) in a fetus with anencephaly diagnosed at 16 weeks' gestation, and dup(2)(p25p22) in another fetus with anencephaly diagnosed at 17 weeks' gestation. Doray et al [37] reported the prenatal diagnosis of partial trisomy 2p (2p22 → pter) in two male sib fetuses by chorionic villus sampling. Both fetuses had NTDs, with craniorachischisis in the first fetus and anencephaly in the second fetus, and the karyotype of 46,XY,der(15)t(2;15)(p22;q26)mat. Doray et al [37] reviewed 64 cases with partial trisomy 2p and found that 15 cases (23.44%) had NTDs, including spina bifida, anencephaly and encephalocele. Lurie et al [39] suggested that the common triplicated region of chromosome 2p associated with NTDs is the band 2p24. Chromosome 2p24 contains the two genes *GDF7* (growth/differentiation factor 7) and *DDEF2* (development and differentiation enhancing factor 2) that are important for neural tube development. *GDF7* is involved in the generation of a discrete class of commissural interneurons in the mouse spinal cord [40], and *DDEF2* is involved in cell communication and structure [41].

Del(2q)

Deletions of 2q have been associated with spina bifida and encephalocele [42–45] (Table 2). McConnell et al [42] reported a partial deletion of chromosome 2, del(2)(q22q31), in a newborn with a prominent occiput, a short mandible, a short sternum, overlapping fingers, rocker-bottom feet, cleft palate, a heart defect, and occipital meningocele. Chinen et al [44] reported a female infant with an interstitial deletion involving 2q24.3 or del(2)(q24.2q31), micrognathia, facial dysmorphism, ocular ptosis, low-set ears, a beaked nose, congenital heart defects, and occipital encephalocele. Melnyk and Muraskas [43] reported an infant with a *de novo* deletion of 2q36, multiple minor anomalies, lumbar meningocele, hydrocephalus, and congenital heart defects. Nye et al [45] reported lumbosacral myelomeningocele and Waardenburg syndrome in patients with an interstitial deletion of 2q35 and the *PAX3* gene. The *PAX3* gene was mapped to chromosome 2q35. In humans, neural tube defects have occasionally been associated with Waardenburg syndrome [46–56]. Hol et al [55] reported a girl with spina bifida, mild signs of Waardenburg syndrome, and a frameshift mutation in the gene for *PAX3*. Shim et al [57] reported a 6-year-old boy with bilateral sensorineural deafness, lateral displacement of inner canthi, a bulbous nasal tip, synophrys, cryptorchidism, Waardenburg syndrome type I, a lumbar spina bifida, and the karyotype of 46,XY,der(2)inv(2)(q13q21)inv(2)(q21q24.2)ins(2)(q24.2q33q35). Fluorescence *in situ* hybridization study showed a breakpoint at 2q35 being proximal to and without involvement of the *PAX* gene. Kujat et al [56] reported the prenatal diagnosis of spina bifida in a fetus, leading to the initial diagnosis of Waardenburg syndrome type I in the members of a four-generation family and a novel splice site mutation within the *PAX3* gene in intron 5 in all affected family members but in none of the unaffected relatives. Stamm et al [58] used a high-density single nucleotide polymorphism screen and identified linkage to loci at 7p21.1–pter and 2q33.1–q35 in a large multiplex NTD family. Other possible candidate genes of NTDs in the involved segment of chromosome 2q include *HOXD* (2q31–q32).

Del(1p), del(1q), and dup(1q)

Partial deletions and duplications of chromosome 1 can be associated with spina bifida and encephalocele [59–66] (Table 3). Thauvin-Robinet et al [65] reported the prenatal diagnosis of anhydramnios, occipital encephalocele, and enlarged bladder at 12 weeks' gestation in a fetus with the karyotype of 46,XY,der(1)t(1;14)(p35;q32)mat, a 1p35–pter deletion and a

Table 3. Reported neural tube defect (NTD) cases with partial aneuploidy of chromosome 1

Authors	Chromosome aberration	Deletion or duplication extent	NTDs
Del(1p)			
Thauvin-Robinet et al [65]	der(1)t(1;14)(p35;q32)mat	1p35 → pter 14q32 → qter	Occipital encephalocele
Del(1q)			
Golabi et al [60]	der(1)t(1;12)(q43;p13)mat	1q43 → qter	Occipital encephalocele,
Case 1 and 2		12p13 → pter	lumbar spina bifida
Ribeiro and Brunoni [62]	del(1)(q43)	1q43 → qter	Occipital meningocele
Wen et al [66]	del(1)(q42)	1q42 → qter	Encephalocele
Al-Awadi et al [61]	del(1)(q32q42)	1q32 → q42	Encephalocele
Takano et al [63]	del(1)(q24q25.3)	1q24 → q25.3	Spina bifida, meningomyelocele
Dup(1q)			
Dagna Bricarelli et al [59]	der(6)t(1;6)(q32;q26)mat*	1q32 → qter 6q26 → qter	Occipital meningomyelocele
Pettenati et al [64]	der(14)t(1;14)(q11;p11.1) [†]	1q11 → qter 14p11.1 → pter	Parietal encephalocele

*Presumed rearrangement; [†]12% (3/25) aberrant cells/analyzed cells in cord blood. Del = deletion; der = derivative chromosome; mat = maternal origin; dup = duplication.

14q32–qter duplication. Golabi et al [60] reported double NTDs of occipital encephalocele and lumbar spina bifida in two sibs with del(1)(q43 → qter), dup(12)(p13 → pter), and the karyotype of 46,XX,der(1)t(1;12)(q43;p13)mat. Ribeiro and Brunoni [62] reported a newborn with hydrocephalus, occipital meningocele, and a terminal deletion 1q43 → qter. Wen et al [66] reported a case with encephalocele and del(1)(q42). Al-Awadi et al [61] reported a malformed infant with a *de novo* interstitial deletion of 1q (1q32 → q42), microcephaly, encephalocele, small eyes with unilateral esotropia, hypertelorism, highly arched palate, micrognathia, low-set ears, a short neck, congenital heart defects, and single umbilical artery. Takano et al [63] reported a 12-month-old boy with the karyotype of 46,XY,del(1)(q24q25.3), spina bifida, meningomyelocele, hydrocephalus, anal atresia, an atrial septal defect, left renal agenesis, bilateral cryptorchidism, talipes equinovarus, low birth weight, growth/developmental retardation, and minor anomalies. Dagna Bricarelli et al [59] reported occipital myelomeningocele in a case with partial trisomy 1q (1q32 → qter) and partial monosomy 6q (6q26 → qter) because of unbalanced translocation inherited from maternal t(1;6)(q32;q26). Pettenati et al [64] reported the prenatal diagnosis of mosaic trisomy 1q in a fetus with mosaic der(14)t(1;14)(q11;p11.1) with nuchal thickening, bitemporal narrowing, a single choroid plexus cyst, mild ventriculomegaly, hyperechoic bowel, hydrops fetalis, and left parietal encephalocele. Possible candidate genes of NTDs in the involved segments of

chromosome 1 include *CRABP2* (1q21.3), *PRKACB* (1p36.1), *PAX7* (1p36.2–p36.12), *MTHFR* (1p36.3) and *MTR* or *MS* (1q43).

Del(3p), dup(3p), del(3q), and dup(3q)

Partial deletions and duplications of chromosome 3 can be associated with spina bifida and encephalocele [4,67–73] (Table 4). Allderdice et al [67] reported spina bifida or sacral dimple in the rec(3)dup(3)(q21 → qter) del(p25 → pter) children of known inv(3)(p25q21) carriers. Fear and Briggs [4] reported absence of the right eye and lumbar spina bifida in an infant with the karyotype of 46,XX,der(13)t(3;13)(q21;q34), and partial trisomy 3q (3q21 → qter) and partial monosomy 13q (13q34 → qter) born to a carrier mother with t(3;13)(q21;q34). Schinzel [72] reported recurrent NTDs in two unbalanced offsprings with dup(3)(q21 → qter) and del(5)(pter → p15.2). The first underweight girl had lumbar myelomeningocele, Arnold-Chiari malformation, agenesis of the corpus callosum, anal stenosis, intestinal malrotation, Meckel's diverticulum, and uterus bicornis. The following prenatally diagnosed fetus had gastroschisis, posterior encephalocele, and uterus bicornis. Kennedy et al [71] reported lumbosacral meningomyelocele, holoprosencephaly, premaxillary agenesis, tetralogy of Fallot, ventricular septal defect, and pulmonary stenosis in a fetus with the karyotype of 46,XY,der(3)del(3)(p26)dup(3)(p26p21.3). Suzumori et al [68] reported the prenatal diagnosis of partial trisomy 3p (3p21 → pter) in a fetus with cleft lip and palate, absent genitalia, lumbosacral

Table 4. Reported neural tube defect (NTD) cases with partial aneuploidy of chromosome 3

Authors	Chromosome aberration	Deletion or duplication extent	NTDs
Del(3p) Bader et al [69]	der(3)t(3;11)(p26;q21)mat	3p26 → pter 11q21 → qter	Lumbosacral meningomyelocele
Dup(3q) and del(3p) Allderdice et al [67] Case 1, 7, 8 and 13	rec(3)dup(3)(q21qter)del(3)(p25)	3q21 → qter 3p25 → pter	Spina bifida or sacral dimple
Dup(3q) Fear and Briggs [4]	der(13)t(3;13)(q21;q34)mat	3q21 → qter 13q34 → qter	Lumbar meningomyelocele
Schintel [72]	dup(3)(q21qter), del(5)(p15.2)	3q21 → qter 5p15.2 → pter	Lumbar meningomyelocele
	dup(3)(q21qter), del(5)(p15.2)	3q21 → qter 5p15.2 → pter	Posterior encephalocele
Dup(3p) and del(3p) Kennedy et al [71]	der(3)del(3)(p26)dup(3)(p26p21.3)	3p21.3 → p26 3p26 → pter	Lumbosacral meningomyelocele, holoprosencephaly
Dup(3p) Suzumori et al [68]	der(11)t(3;11)(p21;q25)mat	3p21 → pter 11q25 → qter	Lumbosacral spina bifida
Del(3q) Kosaki et al [73]	del(3)(q12.2q13.2)	3q12.2 → q13.2	Spina bifida, OEIS complex
Jokiaho et al [70]	del(3)(q27)	3q27 → qter	Parietal meningocele

Del = deletion; der = derivative chromosome; mat = maternal origin; dup = duplication; rec = recombinant chromosome; OEIS = omphalocele-exstrophy-imperforate anus-spinal defects.

spina bifida, and the karyotype of 46,XX,der(11)t(3;11)(p21;q25)mat. Kosaki et al [73] reported an infant with del(3)(q12.2q13.2), spina bifida and omphalocele-exstrophy-imperforate anus-spinal defects (OEIS) complex. Jokiaho et al [70] reported a deletion of 3q27 → qter in an infant with mild dysmorphism, parietal meningocele, and neonatal miliaria rubra-like lesion. Possible candidate genes of NTDs in chromosome 3 include *TERC* (telomerase RNA component) (3q21–q28), *DVL3* (Dishevelled 3) (3q27), *ZIC1* (3q24), and *hOGG1* (3p26.2). Dup(3)(q21 → qter) has been associated with spina bifida and encephalocele [4,67]. The region of 3q21–qter contains the *ZIC1*, *TERC* and *DVL3* genes, and the region of chromosome 3p contains the genes of cell adhesion molecules *CNTN4* and *CNTN6* (3p25–p26) [74], and *CHL1* (3p26.1) [75].

Del(4p), dup(4p), del(4q), and dup(4q)

Partial deletions and duplications of chromosome 4 can be associated with spina bifida and encephalocele [19,76–80] (Table 5). Of interest is the association of Wolf-Hirschhorn syndrome with NTDs. Tachdjian et al

[77] reported three fetuses with distal 4p deletions of 4p14 → pter, 4p15 → pter or 4p16 → pter, Wolf-Hirschhorn syndrome, and sacral dimples. Carbonell et al [76] reported Wolf-Hirschhorn syndrome, cleft lip and palate, polydactyly, multicystic kidneys, and occipital encephalocele in a fetus with a deletion of 4p (4p15 → pter) mimicking Meckel syndrome. Lam et al [19] reported a fetus with exencephaly, dup(4)(p15.2 → pter), and del(13)(q21.2 → qter). The *SLIT2* gene maps to 4p15.2. In *Drosophila* embryogenesis, the *slit* gene has been shown to play a critical role in central nervous system midline formation; and in humans, *Slit-2* mRNA is expressed exclusively in the spinal cord [81]. Quadrelli et al [79] reported a 5-year-old girl with del(4)(q33), facial and digital dysmorphism, a complex congenital heart defect, a large occipital encephalocele, and postnatal growth deficiency. Mikelsaar et al [78] reported a 7-year-old girl with a duplication of 4q (4q25 → qter), mental retardation, facial dysmorphism, a small umbilical hernia, and spina bifida occulta of the 10th and 11th thoracic vertebrae. Nowaczyk et al [80] reported a male infant with an interstitial deletion of chromosome 4,

Table 5. Reported neural tube defect (NTD) cases with partial aneuploidy of chromosomes 4–12, 14–16, 18, 20, 22, and X

Authors	Chromosome aberration	Deletion or duplication extent	NTDs
Del(4p) Tachdjian et al [77] Case 1 Case 2 Case 3 Carbonell et al [76] Dup(4p) Lam et al [19]	del(4)(p14) del(4)(p16) del(4)(p15) del(4)(p15) der(13)t(4;13)(p15.2;q21.2)pat	4p14 → pter 4p16 → pter 4p15 → pter 4p15 → pter 4p15.2 → pter 13q21.2 → qter	Sacral dimple Sacral dimple Sacral dimple Occipital encephalocele Exencephaly
Del(4q) Quadrelli et al [79] Nowaczyk et al [80] Dup(4q) Mikelsaar et al [78] Del(5p) Schinzel [72]	del(4)(q33) del(4)(q13.2q23) der(22)t(4;22)(q25;p11) del(5)(p15.2),dup(3)(q21qter) del(5)(p15.2),dup(3)(q21qter)	4q33 → qter 4q13.2 → q23 4q25 → qter 5p15.2 → pter 3q21 → qter	Occipital encephalocele Occipital encephalocele Spina bifida occulta Lumbar meningocele
Schinzel [72] Mita et al [82] Winsor et al [34] two cases Walbaum et al [31]	del(5)(p15) del(5p) der(5)t(2;5)(p23;p15)mat der(5)t(2;5)(p13;p15)mat	3q21 → qter 5p15 → pter 5p 5p15 → pter 2p23 → pter 5p15 → pter 2p13 → pter	Posterior encephalocele Lumbosacral meningocele Lumbosacral meningocele Anencephaly Anencephaly
Del(6p) Wright et al [85]	der(6)t(6;11)(p?;q?)pat	6p 11q	Craniorachischisis
Dup(6q) Stamberg et al [86] Del(6q) Lukusa et al [87] Dagna Bricarelli et al [59]	der(22)t(6;22)(q21;p13)mat del(6)(q25.3) der(6)t(1;6)(q32;q26)mat*	6q21 → qter 6q25.3 → qter 6q26 → qter 1q32 → qter	Occipital encephalocele Lumbar spina bifida Occipital meningocele
Dup(7p) Carnevale et al [90]	der(14)t(7;14)(p11;p11)mat	7p11 → pter	Frontoethmoidal encephalocele

Case	Cytogenetic	Phenotype
Del(7q)		
Rodríguez et al [93]	del(7)(q36)	Anterior meningocele, sacral agenesis
Drugan et al [11]	del(7)(q32)	Encephalocele
Bogart et al [92]	del(7)(q32)	Anencephaly, craniorachischisis
Case 4		
Stevenson et al [91]	del(7)(q22.1q22.3)	Anencephaly
Dup(8q)		
Williams et al [98]	rec(8)dup q, inv(8)(p23q22)	Occipital encephalocele
Case 3		
Schintel [97]		
Case 1	der(18)t(8;18)(q23;p11.3)mat	Frontal meningocele of the skull, lower thoracic and lumbar spina bifida
Case 2	der(18)t(8;18)(q23;p11.3)mat	Lower thoracic spina bifida
Del(9p)		
Fineman et al [30]	der(9)t(2;9)(p13;p24)mat	Spina bifida
Del(10q)		
Singer et al [32]	der(10)t(2;10)(p24;q26)mat	Anencephaly
Carriere [27]	der(10)t(2;10)(p23;q26)mat*	Encephalocele
Del(11q)		
Suzumori et al [68]	der(11)t(3;11)(p21;q25)mat	Lumbosacral spina bifida
Dup(11q)		
Wright et al [85]	der(6)t(6;11)(p?;q?)pat	Craniorachischisis
Kennedy et al [18]	der(13)t(11;13)(q23.1;q34)mat	Lumbosacral meningocele
Park et al [100]	+der(13)t(11;13)(q21;q14)mat	Sacral spina bifida
Bader et al [69]	der(3)t(3;11)(p26;q21)mat	Lumbosacral meningocele
Dup(12p)		
Golabi et al [60]	der(1)t(1;12)(q43;p13)mat	Occipital encephalocele, lumbar spina bifida
Case 1 and 2		
Dup(14q)		
Thauvin-Robinet et al [65]	der(1)t(1;14)(p35;q32)mat	Occipital encephalocele

(continued on next page)

Table 5. (continued)

Authors	Chromosome aberration	Deletion or duplication extent	NTDs
De(14q) Claussen et al [105]	del(14)(q31)	14q31 → qter	Encephalocele
De(15q) Doray et al [37]	der(15)t(2;15)(p22;q26)mat	15q26 → qter 2p22 → pter	Craniorachischisis
	der(15)t(2;15)(p22;q26)mat	15q26 → qter 2p22 → pter	Anencephaly
Hahm et al [35]	der(15)t(2;15)(p21;q26)	15q26 → qter 2p21 → pter	Anencephaly
Dup(16q) Gustavsson et al [106]	der(11)ins(11;16)(q13;q12.1q22.1)	16q12.1 → q22.1	Lumbosacral meningocele
De(18q) Downton et al [107]	del(18)(q22.2)	18q22.2 → qter	Thoracolumbar meningocele
r(18) Bird et al [109]	r(18)(p11q12)	18p11 → pter 18q12 → qter	Anencephaly
Cohen et al [108]	r(18)	18p	Encephalocele, cyclopia
Dup(20p) Zumel et al [112]	der(15)t(15;20)(p11.2;p12)	20p12 → pter	Anencephaly
+i(20p) Wu et al [113]	+i(20)(p10)	tetrasomy 20p	Occipital encephalocele
De(22q) Nickel et al [116]	del(22)(q11.21q11.23)	22q11.21 → q11.23	Lumbar or lumbosacral meningocele
three cases Forrester et al [117]	del(22)(q11.21q11.23)	22q11.2	Lumbosacral meningocele
two cases Seller et al [118]	del(22)(q11.2q11.2)	22q11.2	Sacral spina bifida
Maclean et al [119]	del(22)(q11.2q11.2)	22q11.2	Sacral meningocele
two cases De(Xp) Plaja et al [124]	del(X)(p22.1)	Xp22.1 → pter	Acrania, cervicodorsal rachischisis
Dup(Xq) Hol et al [125]	dup(X)(q26q27) dup(X)(q26q27)	Xq26 → q27 Xq26 → q27	Lower lumbar spina bifida Lumbosacral meningocele

*Presumed rearrangement. Del = deletion; dup = duplication; der = derivative chromosome; pat = paternal origin; mat = maternal origin; rec = recombinant chromosome; ins = insertion; r = ring chromosome; i = isochromosome.

del(4)(q13.2q23), occipital encephalocele, cleft palate, short limbs, and congenital heart defects.

Del(5p)

Distal 5p deletion or cri du chat syndrome can be associated with spina bifida and anencephaly [31,34,72,82] (Table 5). Schinzel [72] reported a 3-week-old girl with large lumbosacral meningocele, cri du chat syndrome, and del(5)(pter → p15). Mita et al [82] reported a 48-day-old girl with deletion of 5p, cri du chat syndrome, characteristic cry, mental retardation, microcephaly, facial dysmorphism, umbilical hernia, prolapsus ani, dislocation of the hip joint, and lumbosacral meningomyelocele. Of interest is the occurrence of NTDs in concomitant dup(2p) and del(5p), and in concomitant dup(3q) and del(5p). Walbaum et al [31] reported recurrent anencephaly in sib fetuses with partial trisomy 2p (2p13 → pter) and partial monosomy 5p (5p15 → pter). Winsor et al [34] reported recurrent anencephaly in sib fetuses with partial trisomy 2p (2p23 → pter) and partial monosomy 5p (5p15 → pter). Schinzel [72] reported recurrent NTDs in two unbalanced offsprings with dup(3)(q21 → qter) and del(5)(pter → p15.2). The first underweight girl had lumbar myelomeningocele, Arnold-Chiari malformation, agenesis of the corpus callosum, anal stenosis, intestinal malrotation, Meckel's diverticulum, and uterus bicornis. The following prenatally diagnosed fetus had gastroschisis, posterior encephalocele, and uterus bicornis. Chromosome 5p contains the two genes *CDH18* (cadherin-18) (5p15.2–p15.1) and *IRX1* (5p15.3) that are important for neural morphogenesis. *CDH18* is expressed through the central nervous system [83], and *IRX1* is involved in neurogenesis [84].

Del(6p), dup(6q), and del(6q)

Partial deletions and duplications of chromosome 6 can be associated with spina bifida, encephalocele, and craniorachischisis [59,85–87] (Table 5). Wright et al [85] reported craniorachischisis in a fetus with partial trisomy 11q and partial monosomy 6p due to an unbalanced translocation inherited from a carrier father with a reciprocal translocation, t(6p+;11q-). The human *TFAP2A* (transcription factor AP-2α) gene maps to 6p24 and the AP-2 knockout mice have been shown to exhibit anencephaly, craniofacial abnormalities, thoraco-abdominoschisis, and skeletal defects [88]. *JUMONJI*, the human homologue of the mouse *jumonji* gene, is required for the closure of the neural tube and maps to 6p23–p24 [89]. Stamberg et al [86] reported a stillborn infant with occipital encephalocele, ambiguous genitalia, imperforate anus, omphalocele, unilateral hydronephrosis, and the karyotype of 46,XY,der(22)

t(6;22)(q21;p13) or (q21;pter)mat and partial trisomy 6q (6q21 → qter). Lukusa et al [87] reported a 10-year-old mentally retarded boy with terminal 6q25.3 deletion, dysmorphism, abnormal behavior, and lumbar spina bifida. Dagna Bricarelli et al [59] reported occipital myelomeningocele in a case with partial trisomy 1q (1q32 → qter) and partial monosomy 6q (6q26 → qter) because of unbalanced translocation inherited from maternal t(1;6)(q32;q26). Other possible candidate genes of NTDs in the involved segments of chromosome 6 include *Cited 2* (6q23.3), *MACS* (6q22.2), and *T box* (human analogue of Brachyury gene in mice) (6q27).

Dup(7p) and del(7q)

Partial duplications of chromosome 7p can be associated with encephalocele, and partial deletions of chromosome 7q can be associated with spina bifida, encephalocele, and anencephaly [11,90–93] (Table 5). Carnevale et al [90] reported frontoethmoidal encephalocele in an infant with dup(7)(p11 → pter) due to der(14)t(7;14)(p11;p11)mat. Stevenson et al [91] reported a male fetus with anencephaly, a beaked nose, bifid thumbs, hypoplastic genitalia, and the karyotype of 46,XY,del(7)(q22.1q22.3). Rodríguez et al [93] reported an 18-month-old boy with mental retardation, microcephaly, a distinctive face, bilateral coloboma, café-au-lait spot in the abdomen, sacral agenesis, anterior myelomeningocele, and a terminal deletion of the long arm of chromosome 7 (7q36 → qter). Drugan et al [11] reported encephalocele in a case with del(7)(q32). Bogart et al [92] reported a 16-week gestation fetus with anencephaly, complete craniorachischisis, and del(7)(q32). The human *HLXB9* gene maps to 7q36 and is involved in Currarino syndrome with autosomal dominant sacral agenesis and the associated features of anorectal malformation, a presacral mass, and urogenital malformation [94]. *SHH* gene also maps to 7q36 and is responsible for holoprosencephaly 3 (HPE3) [95,96]. Another possible candidate gene of NTDs in the segment involved in chromosome 7 is *HOXA1* (7p15.3).

Dup(8q)

Partial duplications of chromosome 8q can be associated with encephalocele and spina bifida [97,98] (Table 5). Schinzel [97] reported frontal meningocele of the skull and spina bifida of the lower thoracic and lumbar spine, bilateral talipes equinovarus, mental retardation in a girl, and lower thoracic spina bifida in her half-sister, both with dup(8)(q23 → qter) due to der(18)t(8;18)(q23;p11.3)mat. Williams et al [98] reported occipital encephalocele in an infant with

recombinant chromosome 8 syndrome, atrial septal defect, patent ductus arteriosus, bilateral abdominal cryptorchidism, polycystic kidneys and hypoplastic ureters, hydrocephalus, prominent pons, valgus deformity of foot, and dup(8)(q22 → qter) due to paternal inv(8)(p23q22).

Del(9p)

Fineman et al [30] reported spina bifida associated with del(9)(p24 → pter) and dup(2)(p13 → pter) due to der(9)t(2;9)(p13;p24)mat.

Del(10q)

Singer et al [32] reported prenatal diagnosis of del(10)(q26 → qter) and dup(2)(p24 → pter) by amniocentesis in the third pregnancy of a woman with fetal anencephaly and congenital cystic disease of the right kidney. The maternal karyotype was 46,XX,t(2;10)(p24;q26) and the fetal karyotype was 46,XY,der(10)t(2;10)(p24;q26)mat. The woman's first pregnancy resulted in a female fetus with anencephaly and left hydro-nephrosis. Pedigree analysis revealed that the woman had a maternal aunt with spina bifida.

Del(11q) and dup(11q)

Suzumori et al [68] reported the prenatal diagnosis of del(11)(q25 → qter) and dup(3)(p21 → pter) in a fetus with cleft lip and palate, absent genitalia, lumbosacral spina bifida, and the karyotype of 46,XX,der(11)t(3;11)(p21;q25)mat. Craniorachischisis and spina bifida have been described in dup(11q) [18,69,85,99,100]. Rott et al [99] first observed familial NTDs and possible duplication of distal 11q in three cases with myelomeningocele, spina bifida, and spina bifida with meningocele, respectively. Kennedy et al [18] reported a duplication of 11q and a deletion of 13q associated with spina bifida. Wright et al [85] reported a duplication of 11q and a deletion of 6p associated with craniorachischisis. Bader et al [69] reported a fetus with a duplication of the segment 11q21 → qter and a deletion of the segment 3p26 → pter associated with lumbosacral meningomyelocele. Park et al [100] reported the prenatal diagnosis of an 18-week gestation fetus with prominent scoliosis involving the lumbar to sacral regions, sacral rachischisis aperta (spina bifida) with myeloschisis, low-set ears, micrognathia, occipital hypoplasia, a large left congenital diaphragmatic hernia, and dup(11)(q21 → qter) and dup(13)(pter → q14) due to unbalanced 3:1 disjunction of a maternal t(11;13)(q21;q14) translocation. Chromosome 11q contains *NAP1L2* (11q23.1), which is a possible NTD candidate gene, and other genes such as *FOLR1* and *FOLR2* (11q13.3–q13.5) for human folate

receptor [101,102], *MKS2* (11q13) [103], and *Barx2* (11q25) [104].

Dup(12p)

Golabi et al [60] reported double NTDs of occipital encephalocele and lumbar spina bifida in two sibs with dup(12)(p13 → pter) and del(1)(q43 → qter) due to 46,XX,der(1)t(1;12)(q43;p13)mat.

Del(14q) and dup(14q)

Thauvin-Robinet et al [65] reported the prenatal diagnosis of anhydramnios, occipital encephalocele, and enlarged bladder in a 12-week gestation fetus with the karyotype of 46,XY,der(1)t(1;14)(p35;q32)mat, a 1p35–pter deletion and a 14q32–qter duplication. Claussen et al [105] reported encephalocele in a 34-week gestation fetus with del(14)(q31 → qter). The DNA repair gene *XRCC3* maps to 14q32.3.

Del(15q)

Doray et al [37] reported the prenatal diagnosis of del(15)(q26 → qter) and dup(2)(p22 → pter) due to 46,XY,der(15)t(2;15)(p22;q26)mat in two male sib fetuses by chorionic villus sampling. Both fetuses had NTDs, with craniorachischisis in the first fetus and anencephaly in the second fetus. Hahm et al [35] reported the prenatal diagnosis of del(15)(q26 → qter) and dup(2)(p21 → pter) in a 19-week gestation fetus with minor facial anomalies, musculoskeletal defects, postaxial polydactyly, and anencephaly. The fetal karyotype was 46,XY,der(15)t(2;15)(p21;q26).

Dup(16q)

Gustavsson et al [106] reported a girl with a duplication of 16q12.1–q22.1, the karyotype of 46,XX,der(11)ins(11;16)(q13;q12.1q22.1), lumbosacral myelomeningocele, Arnold-Chiari II malformation, mental retardation, strabismus, micrognathia, epileptic seizures, and growth retardation.

Del(18q) and r(18)

Downton et al [107] reported an infant with asymmetry of the left hemithorax, widely spaced nipples, scoliosis, a skin-covered thoracolumbar myelomeningocele, left talipes equinovarus, severe right paraxial fibular hemimelia and the karyotype of 46,XX,del(18)(q22.2). Anencephaly and encephalocele have been described in r(18) [108,109]. Cohen et al [108] reported cyclopia and encephalocele in a case with r(18). Bird et al [109] reported an infant with anencephaly, premaxillary agenesis and the karyotype of 46,XX,r(18)(p11q12). *TGIF* maps to 18p11.3 [110,111] and is responsible for HPE4.

Dup(20p) and +i(20p)

Zumel et al [112] reported prenatal diagnosis of dup(20)(p12 → pter) due to der(15)t(15;20)(p11.2;p12) in a 16-week gestation fetus with anencephaly. Wu et al [113] reported the prenatal diagnosis of pure tetrasomy 20p, or 47,XX,+i(20)(p10) by cord blood sampling in a 24-week gestation fetus with occipital encephalocele, mega-cisterna magna, mesomelic shortening, and clubfeet. Possible candidate genes of NTDs in chromosome 20p include *PAX1* (20p11.2). Helwig et al [114] reported a mouse with spina bifida and double heterozygosity for mutations in *PAX1* and *PGDFRA*. Hol et al [115] reported an amino acid substitution in *PAX1* in a patient with spina bifida.

Del(22q)

All reported NTDs associated with a 22q11.2 microdeletion are spina bifida [116–119]. Kousseff [120] first reported three sibs with sacral myelomeningocele, conotruncal cardiac defects, and minor head and neck anomalies. Toriello et al [121] reported an additional similar case and designated the disorder as “Kousseff syndrome”. Nickel et al [116] reported three unrelated children with meningocele, one with DiGeorge syndrome, two with velocardiofacial syndrome, and all with a 22q11.2 deletion. Seller et al [118] reported a 22-week gestation fetus with sacral spina bifida, conotruncal cardiac defects, cleft palate, facial dysmorphism, Kousseff syndrome, and del(22)(q11.2q11.2). Forrester et al [117] studied the family described by Kousseff [120] and found a deletion of chromosome 22q11–q13 in the proband, his diseased brother, and his father. The proband, his brother and his sister had myelomeningocele and clinical findings consistent with DiGeorge syndrome/velocardiofacial syndrome. Maclean et al [119] reported two cases of sacral meningocele, conotruncal cardiac anomalies, Kousseff syndrome, and a 22q11.2 microdeletion and suggested that Kousseff syndrome is a distinct clinical entity that is genetically heterogeneous. The region of chromosome 22q11 contains *DVL1L1*, which is expressed in several fetal and adult tissues including thymus and heart [122], and *MRPL40* (*NLVCF*), which is expressed in the first and second pharyngeal arches and the neural tube [123].

Del(Xp) and dup(Xq)

Partial deletions or duplications of chromosome X can be associated with anencephaly, spina bifida, and craniorachischisis [124,125] (Table 5). Plaja et al [124] reported an anencephalic fetus with acrania, cervico-dorsal rachischisis, a left diaphragmatic hernia, ipsilateral lung hypoplasia, intestinal malrotation, horseshoe

kidneys, adrenal gland hypoplasia, del(X)(p22.1 → pter), and the karyotype of 46,X,del(X)(p22.1). Hol et al [125] reported familial duplication of Xq26–q27 in two brothers, their mother, and their grandmother. The first brother had lower lumbar spina bifida occulta and deep sacral dimples, and the second brother had a large lumbosacral meningocele. Cytogenetic analysis revealed the karyotype of 46,XY,dup(X)(q26q27), and fluorescence *in situ* hybridization analysis revealed that the duplication extended from Xq26.1 to Xq27.3. Fryns et al [126] reported a case of spina bifida with a *de novo* X/autosomal translocation of t(X;22)(q27;q12.1) pointing to a possible causative gene in Xq27. X-linked inheritance of NTDs has been postulated [127,128]. However, two studies of linkage analysis excluded linkage to the X chromosome [129,130]. *NLGN4* maps to Xq22.33 and is associated with X-linked mental retardation and autism [131]. *ZIC3* maps to Xq26.2 and is a possible candidate gene for NTDs [132,133].

Conclusion

This article provides a comprehensive review of structural chromosomal abnormalities associated with NTDs, such as del(13q), r(13), dup(2p), del(2q), del(1p), del(1q), dup(1q), del(3p), dup(3p), del(3q), dup(3q), del(4p), dup(4p), del(4q), dup(4q), del(5p), del(6p), dup(6q), del(6q), dup(7p), del(7q), dup(8q), del(9p), del(10q), del(11q), dup(11q), dup(12p), dup(14q), del(14q), del(15q), dup(16q), del(18q), r(18), dup(20p), +i(20p), del(22q), del(Xp), and dup(Xq). NTDs may be associated with aneuploidy. Perinatal identification of NTDs should alert one to the possibility of chromosomal abnormalities and prompt a thorough cytogenetic investigation and genetic counseling.

References

1. McGahan JP, Pilu G, Nyberg DA. Neural tube defects and the spine. In: Nyberg DA, McGahan JP, Pretorius DH, Pilu G, eds. *Diagnostic Imaging of Fetal Anomalies*. Philadelphia: Lippincott Williams & Wilkins, 2003:291–334.
2. Allderdice PW, Davis JG, Miller OJ, et al. The 13q⁻ deletion syndrome. *Am J Hum Genet* 1969;21:499–512.
3. Schmid W, Mühlethaler JP, Briner J, Knechtli H. Ring chromosome 13 in a polymalformed anencephalic. *Humangenetik* 1975;27:63–6.
4. Fear C, Briggs A. Familial partial trisomy of the long arm of chromosome 3 (3q). *Arch Dis Child* 1979;54:135–8.
5. Telfer MA, Clark CE, Casey PA, Cowell HR, Stroud HH. Long arm deletion of chromosome 13 with exclusion of esterase D from 13q32–13qter. *Clin Genet* 1990;17:428–32.

6. Jones IM, Palmer CG, Weaver DD, Hodes ME. Study of two cases of ring 13 chromosome using high-resolution banding. *Am J Hum Genet* 1981;33:252-61.
7. Benn PA, Warburton D, Byrne JM, et al. A fetus with a chromosome 13 ring and placenta with chromosome 13 rod/ring mosaicism. *Prenat Diagn* 1983;3:297-302.
8. Rudelli R, Byrne J, Warburton D, et al. Anencephaly in the 13q⁻ syndrome. *Teratology* 1983;27:74A.
9. Palmer CG, Miles JH, Howard-Peebles PN, Magenis RE, Patil S, Friedman JM. Fetal karyotype following ascertainment of fetal anomalies by ultrasound. *Prenat Diagn* 1987;7:551-5.
10. Towfighi J, Ladda RL, Sharkey FE. Purkinje cell inclusions and "atelencephaly" in 13q⁻ chromosomal syndrome. *Arch Pathol Lab Med* 1987;111:146-50.
11. Drugan A, Johnson MP, Dvorin E, Moody J, Krivchenia EL, Schwartz D, Evans MI. Aneuploidy with neural tube defects: another reason for complete evaluation in patients with suspected ultrasound anomalies or elevated maternal serum alpha-fetoprotein. *Fetal Ther* 1989;4:88-92.
12. Brandt CA, Hertz JM, Petersen MB, Vogel F, Noer H, Mikkelsen M. Ring chromosome 13: lack of distinct syndromes based on different breakpoints on 13q. *J Med Genet* 1992;29:704-8.
13. Brown S, Gersen S, Anyane-Yeboah K, Warburton D. Preliminary definition of a "critical region" of chromosome 13 in q32: report of 14 cases with 13q deletions and review of the literature. *Am J Med Genet* 1993;45:52-9.
14. Brown S, Russo J, Chitayat D, Warburton D. The 13q⁻ syndrome: the molecular definition of a critical deletion region in band 13q32. *Am J Med Genet* 1995;57:859-66.
15. Goldsmith CL, Tawagi GF, Carpenter BF, Speevak MD, Hunter AGW. Mosaic r(13) in an infant with aprosencephaly. *Am J Med Genet* 1993;47:531-3.
16. Chen CP, Liu FF, Jan SW, Wang KG, Lan CC. Prenatal diagnosis of partial monosomy 13q associated with occipital encephalocele in a fetus. *Prenat Diagn* 1996;16:664-6.
17. Chen CP, Chern SR, Lee CC, Chen WL, Wang W. Prenatal diagnosis of mosaic ring chromosome 13 with anencephaly. *Prenat Diagn* 2001;21:102-5.
18. Kennedy D, Chitayat D, Winsor EJT, Silver MM, Toi A. Prenatally diagnosed neural tube defects: ultrasound, chromosome, and autopsy or postnatal findings in 212 cases. *Am J Med Genet* 1998;77:317-21.
19. Lam YH, Tang MHY, Ng LKL. 13q⁻ in a fetus with ultrasonographic diagnosis of exencephaly in the first trimester. *Prenat Diagn* 1998;18:634-5.
20. Luo J, Balkin N, Stewart JF, Sarwark JF, Charrow J, Nye JS. Neural tube defects and the 13q deletion syndrome: evidence for a critical region in 13q33-34. *Am J Med Genet* 2000;91:227-30.
21. Brown SA, Warburton D, Brown LY, et al. Holoprosencephaly due to mutations in *ZIC2*, a homologue of *Drosophila* odd-paired. *Nat Genet* 1998;20:180-3.
22. Zhu H, Junker WM, Finnell RH, et al. Lack of association between *ZIC2* and *ZIC3* genes and the risk of neural tube defects (NTDs) in Hispanic populations. *Am J Med Genet A* 2003;116:414-5.
23. Klootwijk R, Groenen P, Schijvenaars M, et al. Genetic variants in *ZIC1*, *ZIC2*, and *ZIC3* are not major risk factors for neural tube defects in humans. *Am J Med Genet A* 2004;124:40-7.
24. Inoue T, Hatayama M, Tohmonda T, Itoharu S, Aruga J, Mikoshiba K. Mouse *Zic5* deficiency results in neural tube defects and hypoplasia of cephalic neural crest derivatives. *Dev Biol* 2004;270:146-62.
25. Chawengsaksophak K, de Graaff W, Rossant J, Deschamps J, Beck F. *Cdx2* is essential for axial elongation in mouse development. *Proc Natl Acad Sci USA* 2004;101:7641-5.
26. Therkelsen AJ, Hulten M, Jonasson J, Lindsten J, Christensen NC, Iversen T. Presumptive direct insertion within chromosome 2 in man. *Ann Hum Genet* 1973;36:367-73.
27. Carriere C. *Etude Clinique et Cytogénétique d'une Enfant Trisomique pour le Segment p23.3-pter du Chromosome 2*. Lyon: Thèse, 1975.
28. Francke U, Jones KL. The 2p partial trisomy syndrome. Duplication of region 2p23 → 2pter in two members of a t(2;7) translocation kindred. *Am J Dis Child* 1976;130:1244-9.
29. Cassidy SB, Heller RM, Chazen EM, Engel E. The chromosome 2 distal short arm trisomy syndrome. *J Pediatr* 1977;91:934-8.
30. Fineman RM, Buyse M, Morgan M. Variable phenotype associated with duplication of different regions of 2p. *Am J Med Genet* 1983;15:451-6.
31. Walbaum R, Peyrat MF, van de Velde MF, Guevin du Masgenet B, Bethouart M, Farriaux JP. Faut-il carotyper les parents d'enfants atteints d'anomalies de fermeture du tube neural. *J Genet Hum* 1984;32:307.
32. Singer N, Gersen S, Warburton D. The value of chromosome analysis in cases of neural tube defects: a case of anencephaly associated with fetal dup(2)(p24-pter). *Prenat Diagn* 1987;7:567-71.
33. Sarda P, Lefort G, Devaux P, Humeau C, Rieu D. Multiple congenital anomalies due to partial 2p13-2pter duplication resulting from an unbalanced X;2 translocation. *Ann Genet* 1992;35:117-20.
34. Winsor SH, McGrath MJ, Khalifa M, Duncan AM. A report of recurrent anencephaly with trisomy 2p23-2pter: additional evidence for the involvement of 2p24 in neural tube development and evaluation of the role for cytogenetic analysis. *Prenat Diagn* 1997;17:665-9.
35. Hahn GK, Barth RF, Schauer GM, Reiss R, Opitz JM. Trisomy 2p syndrome: a fetus with anencephaly and postaxial polydactyly. *Am J Med Genet* 1999;87:45-8.
36. Wellesley D, Boyle T. Regarding trisomy 2p syndrome. *Am J Med Genet* 2000;92:295.
37. Doray B, Favre R, Gasser B, Girard-Lemaire F, Schluth C, Flori E. Recurrent neural tube defects associated with partial trisomy 2p22-pter: report of two siblings and review of the literature. *Genet Couns* 2003;14:165-72.
38. Thangavelu M, Frolich G, Rogers D. Partial duplication 2p as the sole abnormality in two cases with anencephaly. *Am J Med Genet A* 2004;124:170-2.
39. Lurie IW, Ilyina HG, Gurevich DB, et al. Trisomy 2p: Analysis of unusual phenotypic findings. *Am J Med Genet* 1995;55:229-36.
40. Lee KJ, Mendelsohn M, Jessell TM. Neuronal patterning by BMPs: a requirement for GDF7 in the generation of a discrete class of commissural interneurons in the mouse spinal cord. *Genes Dev* 1998;12:3394-407.

41. Ishikawa K, Nagase T, Nakajima D, et al. Prediction of the coding sequences of unidentified human genes. VIII. 78 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res* 1997;4:307-13.
42. McConnell TS, Kornfeld M, McClellan G, Aase J. Partial deletion of chromosome 2 mimicking a phenotype of trisomy 18: case report with autopsy. *Hum Pathol* 1980;11:202-5.
43. Melnyk AR, Muraskas J. Interstitial deletion of chromosome 2 region in a malformed infant. *Am J Med Genet* 1993;45:49-51.
44. Chinen Y, Tohma T, Izumikawa Y, Iha T, Goya Y, Naritomi K. Small interstitial deletion of the long arm of chromosome 2 (2q24.3): further delineation of 2q medial monosomy syndrome. *Jpn J Hum Genet* 1996;41:323-8.
45. Nye JS, Balkin N, Lucas H, Knepper PA, McLone DG, Charrow J. Myelomeningocele and Waardenburg syndrome (type 3) in patients with interstitial deletions of 2q35 and the *PAX3* gene: possible digenic inheritance of a neural tube defect. *Am J Med Genet* 1998;75:401-8.
46. Pantke O, Cohen M. The Waardenburg syndrome. *Birth Defects Orig Artic Ser* 1971;7:147-52.
47. de Saxe M, Kromberg JG, Jenkins T. Waardenburg syndrome in South Africa. Part I. An evaluation of the clinical findings in 11 families. *S Afr Med J* 1984;66:256-61.
48. da-Silva EO. Waardenburg I syndrome: a clinical and genetic study of two large Brazilian kindreds, and literature review. *Am J Med Genet* 1991;40:65-74.
49. Begleiter ML, Harris DJ. Waardenburg syndrome and meningocele. *Am J Med Genet* 1992;44:541.
50. Carezani-Gavin M, Clarren SK, Steege T. Waardenburg syndrome associated with meningomyelocele. *Am J Med Genet* 1992;42:135-6.
51. Chatkupt S, Chatkupt S, Johnson WG. Waardenburg syndrome and myelomeningocele in a family. *J Med Genet* 1993;30:83-4.
52. Hoth CF, Milunsky A, Lipsky N, Sheffer R, Clarren SK, Baldwin CT. Mutations in the paired domain of the human *PAX3* gene cause Klein-Waardenburg syndrome (WS-III) as well as Waardenburg syndrome type I (WS-I). *Am J Hum Genet* 1993;52:455-62.
53. Moline ML, Sandlin C. Waardenburg syndrome and meningomyelocele. *Am J Med Genet* 1993;47:126.
54. Aymé S, Philip N. Possible homozygous Waardenburg syndrome in a fetus with exencephaly. *Am J Med Genet* 1995;59:263-5.
55. Hol FA, Hamel BC, Geurds MP, Mullaart RA, Barr FG, Macina RA, Mariman EC. A frameshift mutation in the gene for *PAX3* in a girl with spina bifida and mild signs of Waardenburg syndrome. *J Med Genet* 1995;32:52-6.
56. Kujat A, Veith VP, Faber R, Froster UG. Prenatal diagnosis and genetic counseling in a case of spina bifida in a family with Waardenburg syndrome type I. *Fetal Diagn Ther* 2007;22:155-8.
57. Shim SH, Wyandt HE, McDonald-McGinn DM, Zackai EZ, Milunsky A. Molecular cytogenetic characterization of multiple intrachromosomal rearrangements of chromosome 2q in a patient with Waardenburg's syndrome and other congenital defects. *Clin Genet* 2004;66:46-52.
58. Stamm DS, Rampersaud E, Slifer SH, et al. High-density single nucleotide polymorphism screen in a large multiplex neural tube defect family refines linkage to loci at 7p21.1-pter and 2q33.1-q35. *Birth Defects Res A Clin Mol Teratol* 2006;76:499-505.
59. Dagna Bricarelli F, Lamba Carbone LD, Caffarena G, Cresta E, Frateschi M, Rasore-Quartino A. Iq32qter trisomy due to maternal translocation t(1;6)(q32;q26). *Pathologica* 1979;71:342. [In Italian]
60. Golabi M, Ito M, Hadley D. Double neural tube defects in two siblings: new features of chromosome 1 deletion syndrome—1q43. *Am J Hum Genet* 1982;34:125A.
61. Al-Awadi SA, Farag TI, Usha R, El-Khalifa MY, Sundareshan TS, Al-Othman SA. Interstitial deletion of the long arm of chromosome 1 [del(1)(q32q42)]. *Am J Med Genet* 1986;23:931-3.
62. Ribeiro MC, Brunoni D. Terminal deletion 1q43 in a newborn with hydrocephalus. *Ann Genet* 1987;30:126-8.
63. Takano T, Yamanouchi Y, Mori Y, et al. Interstitial deletion of chromosome 1q [del(1)(q24q25.3)] identified by fluorescence *in situ* hybridization and gene dosage analysis of apolipoprotein A-II, coagulation factor V, and antithrombin III. *Am J Med Genet* 1997;68:207-10.
64. Pettenati MJ, Berry M, Shashi V, Bowen JH, Harper M. Prenatal diagnosis of complete sole trisomy 1q. *Prenat Diagn* 2001;21:435-40.
65. Thauvin-Robinet C, Callier P, Laurent N, et al. Syndromic encephalocele in a fetal case with a 1p35-pter deletion and a 14q32-qter duplication inherited from a maternal balanced translocation. *Prenat Diagn* 2007;27:555-9.
66. Wen S, Ethen M, Langlois PH, Mitchell LE. Prevalence of encephalocele in Texas, 1999-2002. *Am J Med Genet A* 2007;143:2150-5.
67. Allderdice PW, Browne N, Murphy DP. Chromosome 3 duplication q21 → qter deletion p25 → pter syndrome in children of carriers of a pericentric inversion inv(3)(p25q21). *Am J Hum Genet* 1975;27:699-718.
68. Suzumori K, Koishi T, Manzai M, Yagami Y. A fetus with partial trisomy 3 (p21-pter) detected by prenatal diagnosis. *Jpn J Hum Genet* 1983;28:45-53.
69. Bader PI, Haney SM, Munsick RA, Schubert SR, Hodes ME. Brief clinical report: neural tube defects in dup(11q). *Am J Med Genet* 1984;19:5-8.
70. Jokiaho I, Salo A, Niemi KM, Blomstedt GC, Pihkala J. Deletion 3q27 → 3qter in an infant with mild dysmorphism, parietal meningocele, and neonatal miliaria rubra-like lesions. *Hum Genet* 1989;83:302-4.
71. Kennedy D, Silver MM, Winsor EJT, et al. Inverted duplication of the distal short arm of chromosome 3 associated with lobar holoprosencephaly and lumbosacral meningomyelocele. *Am J Med Genet* 2000;91:167-70.
72. Schinzel A. *Catalogue of Unbalanced Chromosome Aberrations in Man*, 2nd edition. Berlin: Walter de Gruyter, 2001:145, 223.
73. Kosaki R, Fukuhara Y, Kosuga M, et al. OEIS complex with del(3)(q12.2q13.2). *Am J Med Genet A* 2005;135:224-6.
74. Zeng L, Zhang C, Xu J, et al. A novel splice variant of the cell adhesion molecule contactin 4 (*CNTN4*) is mainly expressed in human brain. *J Hum Genet* 2002;47:497-9.
75. Wei MH, Karavanova I, Ivanov SV, et al. *In silico*-initiated cloning and molecular characterization of a novel human member of the *L1* gene family of neural cell adhesion molecules. *Hum Genet* 1998;103:355-64.

76. Carbonell JG, Angel J, Carlan S, Papenhausen PR, Essig YP, Kousseff BG. Deletion 4 (p15 → pter) mimicking *in utero* Meckel-Gruber syndrome. *Proc Greenwood Genet Center* 1991;10:138.
77. Tachdjian G, Fondacci C, Tapia S, Hutten Y, Blot P, Nessmann C. The Wolf-Hirschhorn syndrome in fetuses. *Clin Genet* 1992;42:281-7.
78. Mikelsaar RV, Lurie IW, Illus TE. "Pure" partial trisomy 4q25-qter owing to a *de novo* 4;22 translocation. *J Med Genet* 1996;33:344-5.
79. Quadrelli R, Strehle EM, Vaglio A, et al. A girl with del(4)(q33) and occipital encephalocele: clinical description and molecular genetic characterization of a rare patient. *Genet Test* 2007;11:4-10.
80. Nowaczyk MJ, Teshima IE, Siegel-Bartelt J, Clarke JT. Deletion 4q21/4q22 syndrome: two patients with *de novo* 4q21.3q23 and 4q13.2q23 deletions. *Am J Med Genet* 1997;69:400-5.
81. Itoh A, Miyabayashi T, Ohno M, Sakano S. Cloning and expressions of three mammalian homologues of *Drosophila slit* suggest possible roles for *Slit* in the formation and maintenance of the nervous system. *Brain Res Mol Brain Res* 1998;62:175-86.
82. Mita R, Moriyama T, Sekiya T, Takebe Y. A case of "cri-du-chat" syndrome with meningomyelocele. *No Shinkei Geka* 1980;8:761-5. [In Japanese]
83. Shibata T, Shimoyama Y, Gotoh M, Hirohashi S. Identification of human cadherin-14, a novel neurally specific type II cadherin, by protein interaction cloning. *J Biol Chem* 1997;272:5236-40.
84. Bosse A, Zülch A, Becker MB, Torres M, Gómez-Skarmeta JL, Modolell J, Gruss P. Identification of the vertebrate Iroquois homeobox gene family with overlapping expression during early development of the nervous system. *Mech Dev* 1997;69:169-81.
85. Wright YM, Clark WE, Breg WR. Craniorachischisis in a partially trisomic 11 fetus in a family with reproductive failure and a reciprocal translocation, t(6p plus;11q minus). *J Med Genet* 1974;11:69-75.
86. Stamberg J, Shapiro J, Valle D, Kuhajda FP, Thomas G, Wisnow L. Partial trisomy 6q, due to balanced maternal translocation (6;22) (q21; p13) or (q21; pter). *Clin Genet* 1981;19:122-5.
87. Lukusa T, Willekens D, Lukusa N, De Cock F, Fryns JP. Terminal 6q25.3 deletion and abnormal behaviour. *Genet Couns* 2001;12:213-21.
88. Zhang J, Hagopian-Donaldson S, Serbedzija G, et al. Neural tube, skeletal and body wall defects in mice lacking transcription factor AP-2. *Nature* 1996;381:238-41.
89. Bergé-LeFranc JL, Jay P, Massacrier A, et al. Characterization of the human *jumonji* gene. *Hum Mol Genet* 1996;5:1637-41.
90. Carnevale A, Frias S, del Castillo V. Partial trisomy of the short arm of chromosome 7 due to a familial translocation rcp(7;14)(p11;p11). *Clin Genet* 1978;14:202-6.
91. Stevenson RE, Phelan MC, Stanislovitis P, Klinger K, Schwartz CE. Interstitial deletion of 7q22 in a fetus with anencephaly. *Proc Greenwood Genet Center* 1989;8:56-9.
92. Bogart MH, Cuniff C, Bradshaw C, Jones KL, Jones OW. Terminal deletions of the long arm of chromosome 7: five new cases. *Am J Med Genet* 1990;36:53-5.
93. Rodríguez L, Cuadrado Pérez I, Herrera Montes J, Lorente Jareño ML, López Grondona F, Martínez-Frías ML. Terminal deletion of the chromosome 7(q36-qter) in an infant with sacral agenesis and anterior myelomeningocele. *Am J Med Genet* 2002;110:73-7.
94. Lynch SA, Wang Y, Strachan T, Burn J, Lindsay S. Autosomal dominant sacral agenesis: Currarino syndrome. *J Med Genet* 2000;37:561-6.
95. Belloni E, Muenke M, Roessler E, et al. Identification of Sonic hedgehog as a candidate gene responsible for holoprosencephaly. *Nat Genet* 1996;14:353-6.
96. Roessler E, Belloni E, Gaudenz K, et al. Mutations in the human Sonic Hedgehog gene cause holoprosencephaly. *Nat Genet* 1996;14:357-60.
97. Schinzel A. Partial trisomy 8q in half-sisters with distinct dysmorphic patterns not similar to the trisomy 8 mosaicism syndrome. *Hum Genet* 1977;37:17-26.
98. Williams TM, McConnell TS, Martinez F Jr, Smith ACM, Sujansky E. Clinicopathologic and dysmorphic findings in recombinant chromosome 8 syndrome. *Hum Pathol* 1984;15:1080-4.
99. Rott HD, Schwanitz G, Grosse KP, Alexandrow G. C 11/D 13-translocation in four generations. *Humangenetik* 1972;14:300-5.
100. Park JP, McDermet MK, Doody AM, Marin-Padilla JM, Moeschler JB, Wurster-Hill DH. Familial t(11;13)(q21;q14) and the duplication 11q, 13q phenotype. *Am J Med Genet* 1993;45:46-8.
101. Ragoussis J, Senger G, Trowsdale J, Campbell IG. Genomic organization of the human folate receptor genes on chromosome 11q13. *Genomics* 1992;14:423-30.
102. De Marco P, Moroni A, Merello E, et al. Folate pathway gene alterations in patients with neural tube defects. *Am J Med Genet* 2000;95:216-23.
103. Roume J, Genin E, Cormier-Daire V, et al. A gene for Meckel syndrome maps to chromosome 11q13. *Am J Hum Genet* 1998;63:1095-101.
104. Jones FS, Kiousi C, Copertino DW, Kallunki P, Holst BD, Edelman GM. *Barx2*, a new homeobox gene of the *Bar* class, is expressed in neural and craniofacial structures during development. *Proc Natl Acad Sci USA* 1997;94:2632-7.
105. Claussen U, Ulmer R, Beinder E, Voigt HJ. Six years' experience with rapid karyotyping in prenatal diagnosis: correlations between phenotype detected by ultrasound and fetal karyotype. *Prenat Diagn* 1994;14:113-21.
106. Gustavsson P, Schoumans J, Staaf J, Borg Å, Nordenskjöld M, Annerén G. Duplication 16q12.1-q22.1 characterized by array CGH in a girl with spina bifida. *Eur J Med Genet* 2007;50:237-41.
107. Dowton SB, Hing AV, Sheen-Kaniecki V, Watson MS. Chromosome 18q22.2 → qter deletion and a congenital anomaly syndrome with multiple vertebral segmentation defects. *J Med Genet* 1997;34:414-7.
108. Cohen MM, Storm DF, Capraro VJ. A ring chromosome (No. 18) in a cyclops. *Clin Genet* 1972;3:249-52.
109. Bird LM, Pretorius DH, Mendoza AE, Jones MC. Anencephaly with holoprosencephalic facies due to ring chromosome 18. *Clin Dysmorphol* 1997;6:351-8.
110. Overhauser J, Mitchell HF, Zackai EH, Tick DB, Rojas K, Muenke M. Physical mapping of the holoprosencephaly

- critical region in 18p11.3. *Am J Hum Genet* 1995;57:1080-5.
111. Gripp KW, Edwards MC, Mowat D, et al. Mutations in the transcription factor TGIF in holoprosencephaly. *Am J Hum Genet* 1998;63:A32. [Abstract]
 112. Zumel RM, Darnaude MT, Delicado A, Diaz de Bustamante A, de Torres ML, Lopez Pajares I. Trisomy 20p from maternal translocation and anencephaly. Case report and genetic review. *Ann Genet* 1989;32:247-9.
 113. Wu YC, Fang JS, Lee KF, Estipona J, Yang ML, Yuan CC. Prenatal diagnosis of occipital encephalocele, mega-cisterna magna, mesomelic shortening, and clubfeet associated with pure tetrasomy 20p. *Prenat Diagn* 2003;23:124-7.
 114. Helwig U, Imai K, Schmahl W, Thomas BE, Varnum DS, Nadeau JH, Balling R. Interaction between undulated and Patch leads to an extreme form of spina bifida in double-mutant mice. *Nat Genet* 1995;11:60-3.
 115. Hol FA, Geurds MP, Chatkupt S, et al. PAX genes and human neural tube defects: an amino acid substitution in PAX1 in a patient with spina bifida. *J Med Genet* 1996;33:655-60.
 116. Nickel RE, Pillers DA, Merckens M, Magenis RE, Driscoll DA, Emanuel BS, Zonana J. Velo-cardio-facial syndrome and DiGeorge sequence with meningomyelocele and deletions of the 22q11 region. *Am J Med Genet* 1994;52:445-9.
 117. Forrester S, Kovach MJ, Smith RE, Rimer L, Wesson M, Kimonis VE. Kousseff syndrome caused by deletion of chromosome 22q11-13. *Am J Med Genet* 2002;112:338-42.
 118. Seller MJ, Mohammed S, Russell J, Ogilvie C. Microdeletion 22q11.2, Kousseff syndrome and spina bifida. *Clin Dysmorphol* 2002;11:113-5.
 119. Maclean K, Field MJ, Colley AS, Mowat DR, Sparrow DB, Dunwoodie SL, Kirk EP. Kousseff syndrome: a causally heterogeneous disorder. *Am J Med Genet A* 2004;124:307-12.
 120. Kousseff BG. Sacral meningocele with conotruncal heart defects: a possible autosomal recessive trait. *Pediatrics* 1984;74:395-8.
 121. Toriello HV, Sharda JK, Beaumont EJ. Autosomal recessive syndrome of sacral and conotruncal developmental field defects (Kousseff syndrome). *Am J Med Genet* 1985;22:357-60.
 122. Perrimon N, Mahowald AP. Multiple functions of segment polarity genes in *Drosophila*. *Dev Biol* 1987;119:587-600.
 123. Funke B, Puech A, Saint-Jore B, Pandita R, Skoultschi A, Morrow B. Isolation and characterization of a human gene containing a nuclear localization signal from the critical region for velo-cardio-facial syndrome on 22q11. *Genomics* 1998;53:146-54.
 124. Plaja A, Vendrell T, Sarret E, Toran N, Mediano C. Terminal deletion of Xp in a dysmorphic anencephalic fetus. *Prenat Diagn* 1994;14:410-2.
 125. Hol FA, Schepens MT, van Beersum SE, et al. Identification and characterization of an Xq26-q27 duplication in a family with spina bifida and panhypopituitarism suggests the involvement of two distinct genes. *Genomics* 2000;69:174-81.
 126. Fryns JP, Devriendt K, Moerman P. Lumbosacral spina bifida and myeloschizis in a female foetus with *de novo* X/autosomal translocation (t(X;22)(q27;q121)). *Genet Couns* 1996;7:159-60.
 127. Baraitser M, Burn J. Brief clinical report: neural tube defects as an X-linked condition. *Am J Med Genet* 1984;17:383-5.
 128. Jensson O, Arnason A, Gunnarsdottir H, Petursdottir I, Fossdal R, Hreidarsson S. A family showing apparent X linked inheritance of both anencephaly and spina bifida. *J Med Genet* 1988;25:227-9.
 129. Hol FA, Geurds MP, Jensson O, et al. Exclusion mapping of the gene for X-linked neural tube defects in an Icelandic family. *Hum Genet* 1994;93:452-6.
 130. Newton R, Stanier P, Loughna S, et al. Linkage analysis of 62 X-chromosomal loci excludes the X chromosome in an Icelandic family showing apparent X-linked recessive inheritance of neural tube defects. *Clin Genet* 1994;45:241-9.
 131. Laumonnier F, Bonnet-Brilhault F, Gomot M, et al. X-linked mental retardation and autism are associated with a mutation in the NLGN4 gene, a member of the neuroligin family. *Am J Hum Genet* 2004;74:552-7.
 132. Boyles AL, Hammock P, Speer MC. Candidate gene analysis in human neural tube defects. *Am J Med Genet C Semin Med Genet* 2005;135:9-23.
 133. Kibar Z, Capra V, Gros P. Toward understanding the genetic basis of neural tube defects. *Clin Genet* 2007;71:295-310.