

THE EFFECTS OF CONTINUOUS COMBINED ORAL ESTRADIOL AND NORETHISTERONE ON PULSATILITY INDEX IN INTERNAL CAROTID AND UTERINE ARTERIES IN EARLY POSTMENOPAUSAL TAIWANESE WOMEN—A PRELIMINARY STUDY

Chen-Yu Huang, Chong-Yau Fu¹, Jen-Yu Tseng, Tzay-Shing Yang*, Kuan-Chong Chao

Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, and ¹Division of Biostatistics, Institute of Public Health and School of Medicine, National Yang Ming University, Taipei, Taiwan.

SUMMARY

Objective: This study analyzed whether continuous combined oral estradiol and norethisterone had any effect on the pulsatility index (PI) of the internal carotid and uterine arteries in Taiwanese early postmenopausal women.

Materials and Methods: A group of 40 healthy postmenopausal women with no history of hormone therapy (HT) participated in this study and were randomly subdivided into two groups: HT treatment group ($n=20$) and placebo group ($n=20$). PI was evaluated with color Doppler ultrasound at the beginning of the study and after 4 months of HT (2 mg 17 β -estradiol + 1 mg norethisterone acetate) or placebo.

Results: There was no significant change in the PI of the internal carotid and the uterine arteries after 4 months of HT.

Conclusion: This HT regimen showed no significant negative impact on vascular resistance in Taiwanese early postmenopausal women. Results are compatible with the updated recommendations on HT stating that there is little cardiovascular risk when HT is initiated within a few years of the menopause. [*Taiwan J Obstet Gynecol* 2009;48(1):60–64]

Key Words: estradiol, estrogen/progesterone therapy, hormone therapy, menopause, norethisterone, postmenopause, pulsatility index

Introduction

After the menopause, cardiovascular disease increases and is the leading cause of death among postmenopausal women [1,2]. Over the last decade, many observational studies have shown that hormone therapy

(HT) appears to reduce the risks of vascular disease [3–9]. In 1998, however, the findings of the Heart and Estrogen/Progestin Replacement Study, the first blinded, placebo-controlled, randomized clinical trial of HT for the secondary prevention of cardiac disease in postmenopausal women, found no overall reduction in risk of coronary heart disease events among postmenopausal women [10]. Results of the Women's Health Initiative regarding primary prevention of coronary heart disease showed similar results or even some harm during the first year using HT consisting of estrogen/progesterone therapy (EPT) [11]. However, the population in both clinical trials mainly included patients aged 60–80 years,



ELSEVIER

*Correspondence to: Dr Tzay-Shing Yang, Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, 201, Shih-Pai Road, Section 2, Taipei 112, Taiwan.

E-mail: shc1214clean@gmail.com

Accepted: April 14, 2008

who were more than 10 years from the beginning of the menopause, as opposed to younger women receiving HT in Taiwan.

An increase in vascular resistance, as assessed by Doppler ultrasound, is related to a higher cardiovascular risk among postmenopausal women. The mechanism in which estrogen reduces the pulsatility index (PI), which is an expression of resistance to flow, was first observed in the uterine artery [12], then in the internal carotid artery [13–16]. The effect on peripheral vascular resistance might explain the cardiovascular protection attributed to estrogen. In women with an intact uterus, the concomitant use of progestogen has been suggested to protect the endometrium; women who receive continuous combined HT have a reduced risk of endometrial cancer [17]. Whether there is a difference in the coronary effects when the treatment includes adding progestogen rather than just estrogen alone has not been established. The effects, however, may be more favorable with the use of estrogen alone [18].

The aim of this study was to investigate whether EPT had beneficial or harmful effects on vascular resistance in the uterine and internal carotid arteries in early postmenopausal Taiwanese women.

Materials and Methods

This was a single centre, double-blind, placebo-controlled, randomized study.

Subjects

A group of 40 healthy postmenopausal women without any previous history of hormone use (aged 47–52 years; criteria of inclusion and exclusion shown in Table 1) participated in the study with informed consent. Menopause was diagnosed by natural amenorrhea of at least 6 months, follicle-stimulating hormone (FSH) level > 40 mIU/mL, and estradiol < 20 pg/mL.

Treatments and assessments

First, we evaluated the PI, defined as the ratio of the difference between maximal systolic velocity and final diastolic velocity to mean velocity, all obtained in the same cardiac cycle using color Doppler velocimetry of the internal carotid and uterine arteries, and obtained a baseline. Subsequently, the 40 women were randomized to either the experimental group ($n=20$) or control group ($n=20$). The former received 4 months of HT (2 mg 17 β -estradiol + 1 mg norethisterone acetate), while the latter received matched placebo tablets. We re-evaluated the PI after 4 months of therapy.

Table 1. Criteria of inclusion and exclusion

Inclusion criteria—women who fulfilled all the criteria below:

1. Spontaneous menopause within 3 years and intact uterus
2. Follicle-stimulating hormone > 40 mIU/mL
3. Estradiol < 20 pg/mL
4. Menopausal symptoms

Exclusion criteria—women with any one of the below:

1. Hormone-related neoplasm, such as breast cancer or endometrial cancer
2. Hepatic disease with irreversible liver function impairment or cancer of liver
3. Thromboembolism or cerebral vessel disease
4. Abnormal uterine bleeding of unknown etiology
5. Uncontrolled hypertension or diabetes mellitus
6. History of prior hormone therapy within 3 months
7. Steroid-taking

Color Doppler ultrasound was performed by a well-trained technologist and supervised by an ultrasound specialist.

Statistical analysis

The statistical methods used included descriptive statistics, paired t test and two-sample t test. Treatment effect comparison between the two study groups employed the ratio T_4/T_0 (where T_4 = PI at 4 months and T_0 = PI at baseline), thus individual bias was deleted from the numerator and the denominator of the ratio.

Results

Of the 40 women enrolled, 37 completed the clinical trial, 20 of which were in the HT group and 17 in the placebo group. There was no significant difference between the groups in terms of age, time of menopause, height, and weight (Table 2). The menopause had occurred within 2.5 years (mean, 1.5 years). Baseline PI values for the internal carotid and uterine arteries were similar between the HT group and the placebo group ($p=0.32$ and $p=0.17$ for the internal carotid artery and uterine artery, respectively), in comparison of the mean values (Table 3). Although the results were insignificant in both groups, there was a slight increase in the PI value of the internal carotid artery and a slight decrease in the PI value of the uterine artery from baseline to 4 months in both the HT and the placebo groups (Figure 1 and Table 3). Table 4 and Figure 2 show T_4/T_0 , where the geometric mean of


Table 2. Summary of patient demographics

Characteristics	Hormone therapy (<i>n</i> = 20)	Placebo (<i>n</i> = 17)	<i>p</i>
Age (yr)	51.5 ± 3.70	50.5 ± 2.79	0.359
Time since menopause (mo)	17.73 ± 8.38	19.58 ± 7.71	0.769
Body weight (kg)	54.29 ± 5.91	55.88 ± 6.73	0.449
Body height (cm)	151.5 ± 23.45	155 ± 5.40	0.514

Table 3. Internal carotid artery pulsatility index (PI) and uterine artery PI in different treatment groups and time points

PI	Placebo (<i>n</i> = 17)			Hormone therapy (<i>n</i> = 20)		
	Baseline (T_0^-)	4 months (T_4^-)	Paired difference ($T_0^- - T_4^-$)*	Baseline (T_0^+)	4 months (T_4^+)	Paired difference ($T_0^+ - T_4^+$)†
Internal carotid artery‡						
Mean	1.02	1.13	−0.10	0.96	1.13	−0.16
SD	0.15	0.19	0.15	0.15	0.35	0.36
Uterine artery§						
Mean	2.52	2.08	0.44	3.16	2.58	0.57
SD	1.04	1.30	1.39	1.61	1.17	1.85

*Paired *t* test for $H_1: \mu_D > 0$, where $\mu_D = \text{mean of } (T_0^- - T_4^-)$, $p = 0.99$ (internal carotid artery) and 0.10 (uterine artery); †paired *t* test for $H_1: \mu_D > 0$, where $\mu_D = \text{mean of } (T_0^+ - T_4^+)$, $p = 0.97$ (internal carotid artery) and 0.09 (uterine artery); ‡ $p = 0.32$, comparing the mean values of placebo and hormone treatment groups at baseline; § $p = 0.17$, comparing the mean values of placebo and hormone treatment groups at baseline. SD = standard deviation.

**Figure 1.** The box plots of pulsatility index from the internal carotid artery and uterine artery, with and without hormone therapy, and time points (baseline and 4 months). HRT+ = hormone therapy group; HRT- = placebo group.**Table 4.** Comparing hormone therapy with placebo at T_4/T_0 for the internal carotid artery pulsatility index (PI) and uterine artery PI

T_4/T_0	Hormone therapy on internal carotid artery PI			Hormone therapy on uterine artery PI		
	+	−	<i>p</i> *	+	−	<i>p</i> *
Mean†	1.13	1.10	0.32	0.82	0.77	0.38
95% CI	(1.00, 1.29)	(1.03, 1.18)		(0.63, 1.07)	(0.55, 1.07)	

**t* test $H_0: \mu_- < \mu_+$, in the log transform of the ratio, $\log(T_4/T_0)$, where the analyzed data will be more symmetric and closer to the normal distribution; †geometric mean, the anti-logarithm of the arithmetic mean of $\log(T_4/T_0)$. T_4 = PI at 4 months; T_0 = PI at baseline; + = with hormone therapy; − = without hormone therapy; CI = confidence interval.

the internal carotid artery PI was 1.13 (HT group) and 1.10 (placebo group), and the geometric mean of the uterine artery PI was 0.82 (HT group) and 0.77 (placebo group). Note that the trends from baseline to 4 months

are consistent with the above findings. In addition, from the *t* test, the mean of T_4/T_0 in the HT group was not significantly larger than in the placebo group in either the internal carotid artery PI ($p = 0.32$) or uterine artery




Figure 2. The box plots of pulsatility index (PI) ratio T_4/T_0 , where T_4 =PI at 4 months and T_0 =PI at baseline, in the internal carotid and uterine arteries between hormone treatment (treatment 1) and placebo treatment (treatment 0).

PI ($p=0.38$), i.e. the extent of increase in the internal carotid artery and the decrease in the uterine artery in the HT group was similar to that of the placebo group.

From this study, there is no significant evidence that HT increases PI in the internal carotid or uterine arteries.

Discussion

In this study, after 4 months of HT (2 mg 17β -estradiol + 1 mg norethisterone acetate), no significant changes were noted regarding the PI values of the uterine and internal carotid arteries. This implies that, at least for early postmenopausal Taiwanese women, this regimen of HT does not have a harmful impact on vascular resistance. This is compatible with the consensus that the cardiovascular risks are small when HT is initiated within a few years of menopause [18].

Estrogen benefits the vascular system, including improvement of the lipid profile [19], and production of prostacyclin and nitrogen oxide [20], the latter of which may cause the vasodilatory effects of estrogen (i.e. to reduce the PI). Nearly all studies involving the uterine arteries have demonstrated an important reduction in PI after estrogen administration [15,16,21]. However, the administration of a progestin to protect the endometrium may reduce the positive cardiovascular effects of estradiol. This reduction depends on the dose and the androgenicity of the progestin [22]. Nevertheless, no consensus has been reached regarding duration of use. In our present study, results after 4 months of HT showed no significant reduction in PI on the uterine artery and even a slight increase in the internal carotid artery. As we know, norethisterone is an estrane derivative which has relatively high androgenicity.

In past studies, norethisterone partially antagonized the reduction in the carotid artery PI in postmenopausal women under estrogen replacement therapy [23]. Recent reviews demonstrated that the least antagonistic effect on lipid profile was seen with medrogestone, progesterone and cyproterone acetate. On the other hand, a marked antagonistic effect was observed with medroxyprogesterone acetate, norgestrel and norethindrone [24–30].

In conclusion, not all HT preparations (e.g. EPT) exert the same effects on the vasculature, and the choice of progestogen may influence vascular risk at different levels. In addition, it is possible that a lower dose of progestogen reduces the vascular benefit of estrogen to a lesser extent.

In the present study we observed that there is a tendency towards a reduction in the uterine artery PI value and an increase in that of the internal carotid artery. Similar results were also obtained in a short-term placebo-controlled trial carried out with 50 postmenopausal women [31]. The HT group also received 2 mg of estradiol plus 1 mg of norethisterone acetate and showed no significant decrease in the internal carotid artery PI. This might suggest that the increased flow obtained in the uterine bed cannot be extended to other organs. That is probably because, by comparison to other systems, the reproductive system is much more sensitive to the action of estrogens by having a larger amount of estrogen receptors. These results may explain why EPT had no effect in the prevention of coronary heart disease during the Women's Health Initiative clinical trial [11]. Further clinical trials are needed to clarify if other combinations, dosage, duration and routes of HT are involved in the complex and multiple mechanisms of cardiovascular action in Taiwanese postmenopausal women.

References

1. Mosca L, Collins P, Herrington DM, et al. Hormone replacement therapy and cardiovascular disease: a statement for healthcare professionals from the American Heart Association. *Circulation* 2001;104:499-503.
2. Matthews KA, Kuller LH, Sutton-Tyrrell K, Chang YF. Changes in cardiovascular risk factors during the perimenopause and postmenopause and carotid artery atherosclerosis in healthy women. *Stroke* 2001;32:1104-11.
3. Stampfer MJ, Colditz GA. Estrogen replacement therapy and coronary heart disease: a quantitative assessment of the epidemiologic evidence. *Prev Med* 1991;20:47-63.
4. Sullivan JM, Fowlkes LP. The clinical aspects of estrogen and the cardiovascular system. *Obstet Gynecol* 1996;87(2 Suppl): 36S-43S.
5. Henderson BE, Paganini-Hill A, Ross RK. Decreased mortality in users of estrogen replacement therapy. *Arch Intern Med* 1991;151:75-8.
6. Stampfer MJ, Colditz GA, Willett WC, Manson JE, Rosner B, Speizer FE, Hennekens CH. Postmenopausal estrogen therapy and cardiovascular disease: ten-year follow-up from the Nurses' Health Study. *N Engl J Med* 1991;325:756-62.
7. Bush TL, Barrett-Connor E, Cowan LD, et al. Cardiovascular mortality and noncontraceptive use of estrogen in women: results from the Lipid Research Clinics Program Follow-up Study. *Circulation* 1987;75:1102-9.
8. Barrett-Connor E, Wingard DL, Criqui MH. Postmenopausal estrogen use and heart disease risk factors in the 1980s. *JAMA* 1989;261:2095-100.
9. Matthews KA, Meilahn E, Kuller LH, Kelsey SF, Caggiula AW, Wing RR. Menopause and risk factors for coronary heart disease. *New Engl J Med* 1989;321:641-6.
10. Hulley S, Grady D, Bush T, Furberg C, Herrington D, Riggs B, Vittinghoff E. Randomized trial of estrogen plus progestin for secondary prevention of coronary heart disease in postmenopausal women: the Heart and Estrogen/Progestin Replacement Study (HERS). *JAMA* 1998;280:605-13.
11. Writing Group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *JAMA* 2002;288:321-33.
12. Bourne TH, Hillard TC, Whitehead MI, Crook D, Campbell S. Oestrogens, arterial status and postmenopausal women. *Lancet* 1990;335:1470-1.
13. Gangar KF, Vyas S, Whitehead M, Crook D, Meire H, Campbell S. Pulsatility index in internal carotid artery in relation to transdermal oestradiol and time since menopause. *Lancet* 1991;338:839-42.
14. Sendag F, Terek MC, Karadas N, Bilgin O. Effects of oral and transdermal hormone replacement therapy on internal carotid artery pulsatility indices in postmenopausal women: a prospective, randomized, comparative study. *J Reprod Med* 2001;46:962-8.
15. Battaglia C, Regnani G, Artini PG, Genazzani AR, Primavera MR, Salvatori M, Volpe A. Uterine and cerebral vascularization in postmenopausal women treated with hormone-replacement therapy. *Gynecol Endocrinol* 1999;13:223-9.
16. Cacciato B, Paakkari I, Toivonen J, Tikkanen MJ, Ylikorkala O. Randomized comparison of oral and transdermal hormone replacement on carotid and uterine artery resistance to blood flow. *Obstet Gynecol* 1998;92:563-8.
17. Doherty JA, Cushing-Haugen KL, Saltzman BS, et al. Long-term use of postmenopausal estrogen and progestin hormone therapies and the risk of endometrial cancer. *Am J Obstet Gynecol* 2007;197:139.e1-7.
18. Pines A, Sturdee DW, Birkhauser MH, Schneider HP, Gambacciani M, Panay N. IMS updated recommendations on postmenopausal hormone therapy. *Climacteric* 2007;10: 181-94.
19. Stevenson JC, Crook D, Godsland IF. Influence of age and menopause on serum lipids and lipoproteins in healthy women. *Atherosclerosis* 1993;98:83-90.
20. Ylikorkala O, Cacciato B, Paakkari I, Tikkanen MJ, Viinikka L, Toivonen J. The long-term effects of oral and transdermal postmenopausal hormone replacement therapy on nitric oxide, endothelin-1, prostacyclin, and thromboxane. *Fertil Steril* 1998;69:883-8.
21. Bonilla-Musoles F, Martí MC, Ballester MJ, Raga F, Osborne NG. Normal uterine arterial blood flow in postmenopausal women assessed by transvaginal color doppler sonography: the effect of hormone replacement therapy. *J Ultrasound Med* 1995;14:497-501.
22. Godsland IF. Effects of postmenopausal hormone replacement therapy on lipid, lipoprotein, and apolipoprotein (a) concentrations: analysis of studies published from 1974 to 2000. *Fertil Steril* 2001;75:898-915.
23. Luckas MJ, Gleeve T, Biljan MM, Buckett WM, Aird IA, Drakeley A, Kingsland CR. The effect of progestagens on the carotid artery pulsatility index in postmenopausal women on oestrogen replacement therapy. *Eur J Obstet Gynecol Reprod Biol* 1998;76:221-4.
24. Sitruk-Ware R. Progestogens in hormonal replacement therapy: new molecules, risks, and benefits. *Menopause* 2002; 9:6-15.
25. Rosano GM, Fini M. Postmenopausal women and cardiovascular risk: impact of hormone replacement therapy. *Cardiol Rev* 2002;10:51-60.
26. Rosano GM, Fini M. Comparative cardiovascular effects of different progestins in menopause. *Int J Fertil Womens Med* 2001;46:248-56.
27. Stefanick ML. Estrogen, progestogens and cardiovascular risk. *J Reprod Med* 1999;44:221-6.
28. Clarkson TB. Progestogens and cardiovascular disease: a critical review. *J Reprod Med* 1999;44:180-4.
29. Sitruk-Ware R. Progestins and cardiovascular risk markers. *Steroids* 2000;65:651-8.
30. Rosano GM, Sarais C, Zoncu S, Mercuro G. The relative effects of progesterone and progestins in hormone replacement therapy. *Hum Reprod* 2000;15:60-73.
31. Bain CA, Walters MR, Lees KR, Lumsden MA. The effect of HRT on cerebral haemodynamics and cerebral vasomotor reactivity in post-menopausal women *Hum Reprod* 2004; 19:2411-4.