

USE OF A GONADOTROPIN-RELEASING HORMONE AGONIST TO MANAGE PERIMENOPAUSAL WOMEN WITH SYMPTOMATIC UTERINE MYOMAS

Peng-Hui Wang^{1,2*}, Wen-Ling Lee^{2,3}, Ming-Huei Cheng^{1,2}, Ming-Shyen Yen¹,
Kuan-Chong Chao¹, Hsiang-Tai Chao^{1,2}

¹Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, ²Institute of Clinical Medicine, National Yang-Ming University School of Medicine, ³Division of Endocrinology and Metabolism, Department of Medicine, Cheng Hsin Rehabilitation Center—Taipei, Taipei, Taiwan.

SUMMARY

Objective: To determine the acceptability and effectiveness of a gonadotropin-releasing hormone (GnRH) agonist for the treatment of perimenopausal women with symptomatic uterine myomas.

Materials and Methods: The participants included 43 women with symptomatic myomas who wished to retain their uteri. All the women were older than 45 years old, agreed to use the GnRH agonist for menopause induction, and were without any underlying malignancy. They were treated with six courses of GnRH agonist between 2004 and 2005. The definition of re-intervention included: (1) surgical intervention, such as hysterectomy, myomectomy or laparoscopic uterine vessel occlusion, or (2) modification of GnRH agonist use. Modification of GnRH agonist use included either failure to complete a 6-month GnRH agonist treatment course, or re-use of GnRH agonist with/without interruption of continuity. Failure was defined as women who underwent surgical intervention or failed to complete the 6-month GnRH agonist treatment. Evaluations were performed every 6 months, for up to 2 years.

Results: Re-intervention rates were 14.0% ($n=6$), 23.3% ($n=10$) and 32.6% ($n=14$), and failure rates were 7.0% ($n=3$), 11.6% ($n=5$) and 16.3% ($n=7$), at the end of the 6-, 12- and 24-month follow-up periods, respectively. Three patients failed to complete the 6-month GnRH agonist treatment, and four received surgical interventions.

Conclusion: More than 80% of women in this study benefited from the use of GnRH agonist to produce menopause, suggesting that this can be an alternative choice for managing perimenopausal women with symptomatic uterine myomas. [*Taiwan J Obstet Gynecol* 2009;48(2):133–137]

Key Words: anemia, gonadotropin-releasing hormone agonist, perimenopause, uterine myoma

Introduction

Uterine myomas represent the most common benign tumors of the female reproductive system [1,2]. They are

mostly asymptomatic [3], and expectant management is, therefore, a common choice [4]. Some myomas, however, can cause symptoms and require treatment [5,6]. For women who wish to retain their childbearing potential, organ-preserving strategies, such as myomectomy, uterine artery ligation or embolization, focal ultrasound surgery, or various kinds of medical treatment, have been the preferred treatments [7–12]. Hysterectomy is still considered to be the treatment of choice in women with symptomatic uterine myomas who have completed their child bearing [13]. However,



*Correspondence to: Dr Peng-Hui Wang, Department of Obstetrics and Gynecology, Taipei Veterans General Hospital and National Yang-Ming University, 201, Section 2, Shih-Pai Road, Taipei 112, Taiwan.
E-mail: phwang@vghtpe.gov.tw
Accepted: March 27, 2008

the uterus has been psychologically regarded as a regulator and controller of important physiologic functions, a sexual organ, a source of energy and vitality, and a maintainer of youth and attractiveness [14].

Reproductive sex hormones such as estrogen and progesterone appear to be critical factors related to uterine myomas. Evidence for this includes the fact that: (1) myomas are rarely observed before puberty; (2) they are most prevalent during the reproductive years, and regress after menopause; and (3) factors that increase overall lifetime exposure to estrogen, such as obesity and early menarche, increase the incidence of myomas [15]. This provides us with the possibility of manipulating sex hormones when treating women with symptomatic uterine myomas. However, because the effects of medical treatment are often transient [15] and the symptoms and tumor frequently recur after medication is stopped [16], the use of medical treatment for symptomatic uterine myomas is limited [17,18].

Although medical treatment is seldom considered a definitive treatment, it may be beneficial in a given population; women who are near menopausal status (age ≥ 45 years, defined as perimenopausal) with symptomatic uterine myomas might gain from medical treatment [19,20]. Many drugs are currently available for the management of symptomatic uterine myomas, including gonadotropin-releasing hormone (GnRH) agonists, GnRH antagonists, the progesterone-mediated intrauterine system, selective estrogen receptor modulators, selective progesterone receptor modulators, androgen-mediated therapy, growth factor-mediated therapy, and others [15]. Among these medical treatments, the use of GnRH agonists is possibly one of the best choices. Some perimenopausal women might transit to natural menopause in an unexpectedly short period, and the use of a GnRH agonist, leading to a natural menopause, is a rational choice. In this study, we explored the effectiveness and limitations of a GnRH agonist for the management of perimenopausal women with symptomatic uterine myomas.

Materials and Methods

Between January 2004 and December 2005, women with symptomatic uterine myomas scheduled for uterine-sparing treatment were invited to participate in the present study. All the uterine-sparing treatments used for symptomatic uterine myomas were well-accepted procedures in our hospital. They included myomectomy (performed by traditional exploratory laparotomy, mini-laparotomy, ultramini-laparotomy, laparoscopy

or hysteroscopy), laparoscopic uterine vessel occlusion (LUVVO), combination therapy, and medical treatment, including GnRH agonists, progestins, nonsteroidal anti-inflammatory drugs, or pure consultation and follow-up [5,6,10–12,15]. All patients had uterine fibroids with symptoms comprising either menstrual problems, such as menorrhagia and pain, or compression syndrome, including a bulge-like sensation and frequency. These women were informed that they could choose to be treated with any one of the above-mentioned procedures, based on their preference.

This study was designed to assess the effectiveness and limitation of a GnRH agonist for the treatment of fibroids in older women (age ≥ 45 years). Inclusion criteria were: agreeing to use a GnRH agonist for menopause induction for at least 6 months without surgical intervention, no underlying malignancy, an initial strong willingness to avoid surgical treatment if possible, and commitment to completing a thorough, 2-year follow-up record. Surgical interventions included hysterectomy, myomectomy, LUVVO or uterine artery embolization. A total of 43 patients were included, and all of them received monthly treatment with a depot GnRH agonist (leuporelin; Leuplin-Depot; Takeda Pharmaceuticals, Osaka, Japan).

Leiomyoma-related symptoms, including either menstrual problems such as menorrhagia and pain, or compression syndrome (including a bulge-like sensation and frequency), were assessed using a five-point scale to rate changes in symptoms relative to baseline. An initial score of 3 was given if the symptom was present in these women, and subsequent scores ranged from 1 (marked exacerbation), 2 (slight exacerbation), 3 (consistent with the baseline), 4 (slight improvement), to 5 (significant improvement). The same five-point scale was also used to record satisfaction with a baseline score of 3, and ranging from 1 (very dissatisfied) to 5 (very satisfied).

At each follow-up evaluation (6, 12 and 24 months), any medical or surgical treatment of the myoma was recorded. Re-intervention included: (1) failure, defined by the need for hysterectomy, myomectomy, uterine vessel occlusion for uterine myomas, or discontinuation of GnRH agonist treatment (treatment for < 6 cycles) due to poor patient compliance; and (2) fulfillment of the criteria of success, such as re-use of GnRH agonist after completing the 6-month GnRH agonist treatment, and continuous use of GnRH agonist for 24 months without cessation.

SPSS version 12.0 (SPSS Inc., Chicago, IL, USA) was used to analyze patient data. Paired *t* tests were used to analyze changes in each score from baseline. Statistical significance was defined as $p < 0.05$.

Results

The mean age of the patients was 47.4 ± 1.9 years. The mean baseline hemoglobin level was 9.2 ± 1.9 g/dL. Additional baseline characteristics are provided in Table 1.

In this study, continuous vaginal spotting was noted in >50% of the women ($n=24$) after the first dose of GnRH agonist. However, after the second dose (1 month later), the incidence of continuous vaginal spotting had decreased to 21% ($n=9$), and after the third dose, nearly all women had amenorrhea (98%, $n=42$). Three patients (7%) stopped GnRH agonist treatment after the third dose owing to intolerable or severe climacteric symptoms. Although additional hormone supplements were prescribed at the second dose of GnRH agonist, these patients decided to stop any further treatment. Finally, all women had amenorrhea by the sixth dose when they continued with GnRH agonist treatment during the study period.

The re-intervention rate was 14% ($n=6$) at the end of the 6-month follow-up, including three patients who

failed to complete the 6-month GnRH agonist treatment, and three patients who continued to receive GnRH agonist monthly treatment after the 6-month period. The re-intervention rate increased to 23.3% ($n=10$) at the end of the 12-month follow-up. This included the above-mentioned patients ($n=6$), as well as two patients who were treated with re-administration of the GnRH agonist, one who underwent LUVU, and one who underwent myomectomy. At the end of the 24-month follow-up, the re-intervention rate was 32.6% ($n=14$). This included the previous patients, plus a further two who were re-administered GnRH agonist, and two who underwent hysterectomy. The symptom change and satisfaction scores improved significantly between the successive 6-month follow-ups (at the end of 6, 12 and 24 months) and baseline ($p=0.001$), suggesting that the women who were treated with the GnRH agonist benefited from symptom improvement, with a resultant high level of satisfaction. A summary of the clinical outcomes is presented in Table 2.

Discussion

GnRH agonists are among the most effective and well-accepted drugs for the management of myomas. They are analogs of endogenous GnRH and bind to pituitary GnRH receptors (GnRH-R), resulting in the synthesis and release of luteinizing hormone and follicle-stimulating hormone. Compared with GnRH, GnRH agonists have a longer half-life, leading to continuous exposure of GnRH-R under the activity of GnRH agonists. Downregulation of the GnRH-R consequently results in a decreased level of gonadotropin and a

Table 1. Clinical characteristics of the patients ($n=43$)*

Age (yr)	47.4 ± 1.9
Body mass index	22.46 ± 1.56
Baseline hemoglobin level (g/dL)	9.2 ± 1.9
Symptom	
Pain	8 (18.6)
Menorrhagia	30 (69.8)
Bulge sensation	17 (39.5)
Frequency	15 (34.9)

*Data are presented as mean \pm standard deviation or n (%).

Table 2. Characteristics of subjects ($n=43$) during follow-up*

	Follow-up			
	Baseline	6 months	12 months	24 months
Re-intervention	0 (0)	6 (14)	10 (23.3)	14 (32.6)
Hemoglobin level (g/dL)	9.16 ± 2.18	12.87 ± 0.55	11.22 ± 0.22	11.16 ± 0.27
Symptom [†]				
Pain	0.56 ± 1.18	$0.91 \pm 0.29^{\ddagger}$	$0.81 \pm 0.26^{\ddagger}$	$0.79 \pm 0.26^{\ddagger}$
Menorrhagia	2.09 ± 1.39	$3.49 \pm 0.35^{\ddagger}$	$2.74 \pm 0.21^{\ddagger}$	$2.88 \pm 0.34^{\ddagger}$
Bulge sensation	1.19 ± 1.48	$1.81 \pm 0.35^{\ddagger}$	$1.65 \pm 0.32^{\ddagger}$	$1.60 \pm 0.32^{\ddagger}$
Frequency	1.05 ± 1.45	$1.58 \pm 0.39^{\ddagger}$	$1.47 \pm 0.32^{\ddagger}$	$1.44 \pm 0.32^{\ddagger}$
Amenorrhea	0 (0)	40 (93.0)	14 (32.5)	20 (46.5)
Satisfaction [†]	3.00 ± 0.00	$4.44 \pm 0.73^{\ddagger}$	$4.12 \pm 0.96^{\ddagger}$	$4.28 \pm 0.98^{\ddagger}$
Failure		3 (7.0)	5 (11.6)	7 (16.3)

*Data are presented as n (%) or mean \pm standard deviation; [†]the definitions of symptom (pain, menorrhagia, bulge sensation, frequency) change and satisfaction are available in the text; [‡]compared with baseline, all $p < 0.005$.

suppressed production of ovarian hormones [21]. The administration of GnRH agonists leads to extremely low “pseudomenopausal” or “pseudopremenarchal” estradiol levels, resulting in amenorrhea.

In terms of alleviating menorrhagia, it has been reported that bleeding problems or related anemia were controlled in women after the first month of GnRH agonist treatment [22,23].

Pain and pressure symptoms were reported to be relieved within the first 3 months of treatment. This effect can be achieved within the first 12 weeks of therapy, because it is considered to be a consequence of reducing the size of the uterus and myoma [24]. In addition, it has been reported that the effect of GnRH agonists on reducing myoma size is transient; reversal of estrogen deprivation occurs about 4 weeks after discontinuation of the GnRH agonist [25] and most myoma nodules returned to their initial size within about 6 months after discontinuation of treatment. The rate of enlargement was then rapid compared with natural enlargement [22], and regrowth of the uterus and myomas leads to a recurrence of the initial symptoms [26]. We did not routinely monitor uterine and myoma size in the current study, so were unable to assess any changes. However, all but three of the women (who failed to complete the 6-month GnRH agonist treatment) were satisfied with their symptom relief, including pain, bulging sensation and frequency, especially after the initial 6 months. The symptom improvement was maintained up to the end of the 12- and 24-month follow-ups (Table 2), suggesting that the use of GnRH agonist treatment for perimenopausal women with symptomatic uterine myomas should be considered before choosing a more invasive surgical intervention.

The significant disadvantages of GnRH agonists include hypoestrogenism-related effects resembling postmenopausal climacteric disturbances and bone loss [27]. Partial restoration of the estrogenic state by reduced-dose therapy or steroid “add-back” has been shown to be effective for the relief of annoying side effects, without any loss of the beneficial effects on myoma size and cycle suppression [28,29]. The addition of oral veralipride, a benzamide derivative, can reduce the vasomotor symptoms induced by GnRH agonists [30]. Raloxifene or tibolone may also be used as add-back therapy to prevent bone loss whilst preserving the efficacy of GnRH agonists [31,32]. However, raloxifene did not reduce vasomotor symptoms related to GnRH agonists [31], and could even exacerbate climacteric symptoms [33,34]. In contrast, a significant reduction in hot flushes was observed when tibolone was added to GnRH agonist treatment [32]. Either a lower dose of hormone replacement therapy or use of tibolone

can be successfully applied in symptomatic menopausal women [35–39]. In this study, only three women who failed to complete the 6-month GnRH agonist treatment were reported to have intolerable climacteric symptoms. Although add-back therapy was initiated during the third month, the patients were still anxious about the risk of breast cancer and finally decided to stop any medication, except for nonsteroidal anti-inflammatory drugs and iron replacement. In contrast, more than 90% of the perimenopausal women tolerated the treatment well. In addition, three women chose to continue monthly GnRH agonist treatment up to 24 months and two women were re-treated with GnRH agonist 12 months later.

Amenorrhea was reported in only 50% of the women in this study, at the end of the 24-month follow-up period. This was in contrast to the results of Imai et al [19], who found that the depot GnRH agonist, leuporelin acetate, produced an excellent success rate (>90%) by inducing natural menopause in 19 of 21 perimenopausal women with uterine myomas [19].

In conclusion, more than 80% of women in this study experienced significant symptom relief after treatment with a GnRH agonist. These results indicate that GnRH agonists should be considered as an option for the management of perimenopausal women with uterine myomas, in order to reduce the use of more invasive surgical interventions.

Acknowledgments

This study was supported, in part, by grants (V96ED1-003, V97ED1-008) from Taipei Veterans General Hospital.

References

1. Practice Committee of the American Society for Reproductive Medicine. Myomas and reproductive function. *Fertil Steril* 2006;86(Suppl 5):S194–9.
2. Parker WH. Etiology, symptomatology, and diagnosis of uterine myomas. *Fertil Steril* 2007;87:725–36.
3. Cheng MH, Chao HT, Wang PH. Unusual clinical presentation of uterine myomas. *Taiwan J Obstet Gynecol* 2007;46:323–4.
4. Cheng MH, Chao HT, Wang PH. Medical treatment for uterine myomas. *Taiwan J Obstet Gynecol* 2008;47:18–23.
5. Wen KC, Sung PL, Chao KC, Lee WL, Liu WM, Wang PH. A prospective short-term evaluation of uterine leiomyomas treated by myomectomy through conventional laparotomy or ultraminilaparotomy. *Fertil Steril* 2008;90:2361–6.
6. Wang PH, Liu WM, Fuh JL, Chao HT, Yuan CC, Chao KC. Comparison of ultraminilaparotomy for myomectomy

- through midline vertical incision or modified Pfannenstiell incision—a prospective short-term follow-up. *Fertil Steril* 2009;91:1945–50.
7. Parker WH. Uterine myomas: management. *Fertil Steril* 2007;88:255–71.
8. Taylor E, Gomel V. The uterus and fertility. *Fertil Steril* 2008;89:1–16.
9. Di Spiezio Sardo A, Mazzon I, Gargano V, et al. Hysteroscopic treatment of atypical polypoid adenomyoma diagnosed incidentally in a young infertile woman. *Fertil Steril* 2008;89:456,9–12.
10. Liu WM, Tzeng CR, Chen YJ, Wang PH. Combining the uterine depletion procedure and myomectomy may be useful for treating symptomatic fibroids. *Fertil Steril* 2004;82:205–10.
11. Liu WM, Ng HT, Wu YC, Yen YK, Yuan CC. Laparoscopic bipolar coagulation of uterine vessels: a new method for treating symptomatic fibroids. *Fertil Steril* 2001;75:417–22.
12. Liu WM, Wang PH, Chou CS, Tang WL, Wang IT, Tzeng CR. Efficacy of combined laparoscopic uterine artery occlusion and myomectomy via minilaparotomy in the treatment of recurrent uterine myomas. *Fertil Steril* 2007;87:356–61.
13. Wang PH, Chao HT, Lee WL. Rationale of myomectomy for perimenopausal women. *Maturitas* 2007;58:406–7.
14. Gavai M, Berkes E, Takacs ZF, Papp Z. Can myomectomy be suggested for perimenopausal women before administering hormone replacement therapy? *Maturitas* 2007;58:107–10.
15. Cheng MH, Wang PH. Uterine myoma: a condition amenable to medical therapy? *Expert Opin Emerg Drugs* 2008;13:119–33.
16. Lee WL, Yuan CC, Wang PH. Positron emission tomography and uterine leiomyomas. *Gynecol Oncol* 2007;107:593–4.
17. Lee WL, Liu RS, Yuan CC, Chao HT, Wang PH. Relationship between gonadotropin-releasing hormone agonist and myoma cellular activity: preliminary findings on positron emission tomography. *Fertil Steril* 2001;75:638–9.
18. Chia CC, Huang SC, Chen SS, et al. Ultrasonographic evaluation of the change in uterine fibroids induced by treatment with a GnRH analog. *Taiwan J Obstet Gynecol* 2006;45:124–8.
19. Imai A, Sugiyama M, Furui T, Tamaya T. Treatment of perimenopausal women with uterine myoma: successful use of a depot GnRH agonist leading to a natural menopause. *J Obstet Gynaecol* 2003;23:518–20.
20. Fedele L, Bianchi S, Frontino G. Hormonal treatments for adenomyosis. *Best Pract Res Clin Obstet Gynaecol* 2008;22:333–9.
21. Clayton RN. Gonadotrophin-releasing hormone: its actions and receptors. *J Endocrinol* 1989;120:11–9.
22. Friedman AJ, Lobel SM, Rein MS, Barbieri RL. Efficacy and safety considerations in women with uterine leiomyomas treated with gonadotropin-releasing hormone agonists: the estrogen threshold hypothesis. *Am J Obstet Gynecol* 1990;163:1114–9.
23. Lethaby A, Vollenhoven B, Sowter M. Efficacy of pre-operative gonadotrophin hormone releasing analogues for women with uterine fibroids undergoing hysterectomy or myomectomy: a systematic review. *Br J Obstet Gynaecol* 2002;109:1097–108.
24. Cirkel U, Ochs H, Schneider HP, et al. Experience with leuprorelin acetate depot in the treatment of fibroids: a German multicentre study. *Clin Ther* 1992;14:37–50.
25. Matta WH, Shaw RW, Nye M. Long-term follow-up of patients with uterine fibroids after treatment with the LHRH agonist buserelin. *Br J Obstet Gynaecol* 1989;96:200–6.
26. Rein MS, Friedman AJ, Stuart JM, MacLaughlin DT. Fibroid and myometrial steroid receptors in women treated with gonadotropin-releasing hormone agonist leuprolide acetate. *Fertil Steril* 1990;53:1018–23.
27. Donnez J, Hervais Vivancos B, Kudela M, Audebert A, Jadoul P. A randomized, placebo-controlled, dose-ranging trial comparing fulvestrant with goserelin in premenopausal patients with uterine fibroids awaiting hysterectomy. *Fertil Steril* 2003;79:1380–9.
28. Broekmans FJ, Hompes PG, Heitbrink MA, Netelenbos CC, Roos JC, Falke TM, Schoemaker J. Two-step gonadotropin-releasing hormone agonist treatment of uterine leiomyomas: standard-dose therapy followed by reduced-dose therapy. *Am J Obstet Gynecol* 1996;175:1208–16.
29. Friedman AJ, Daly M, Juneau-Norcross M, Gleason R, Rein MS, Leboff M. Long-term medical therapy for leiomyomata uteri: a prospective, randomized study of leuprolide acetate depot plus either oestrogen-progestin or progestin “add back” for 2 years. *Hum Reprod* 1994;9:1618–25.
30. Vercellini P, Sacerdote P, Trespidi L, Manfredi B, Panerai AE, Crosignani PG. Veralipride for hot flushes induced by a gonadotropin-releasing hormone agonist: a controlled study. *Fertil Steril* 1994;62:938–42.
31. Palomba S, Orio F Jr, Russo T, et al. Long-term effectiveness and safety of GnRH agonist plus raloxifene administration in women with uterine leiomyomas. *Hum Reprod* 2004;19:1308–14.
32. Palomba S, Orio F Jr, Falbo A, Oppedisano R, Tolino A, Zullo F. Tibolone reverses the cognitive effects caused by leuprolide acetate administration, improving mood and quality of life in patients with symptomatic uterine leiomyomas. *Fertil Steril* 2008;90:165–73.
33. Lee WL, Cheng MH, Chao HT, Wang PH. The role of selective estrogen receptor modulators on breast cancer: from tamoxifen to raloxifene. *Taiwan J Obstet Gynecol* 2008;47:24–31.
34. Wang PH, Chao HT. A reconsideration of tamoxifen use for breast cancer. *Taiwan J Obstet Gynecol* 2007;46:93–5.
35. Wang PH, Horng HC, Cheng MH, Chao HT, Chao KC. Standard and low-dose hormone therapy for postmenopausal women—focus on the breast. *Taiwan J Obstet Gynecol* 2007;46:127–34.
36. Wang PH, Cheng MH, Chao HT, Chao KC. Effects of tibolone on the breast of postmenopausal women. *Taiwan J Obstet Gynecol* 2007;46:121–6.
37. Wang PH, Yen MS, Chao KC, Liu CM, Lee WL. The short-term effect of increased body weight in menopausal Chinese women taking tibolone. *Maturitas* 2008;59:281–2.
38. Chao KC, Wang PH, Yen MS, Huang CY, Juang CM, Twu NF, Wu HS. New selective tissue estrogenic activity regulator in menopausal therapy in Taiwan. *Taiwan J Obstet Gynecol* 2005;44:327–31.
39. Chen FP. Hormone therapy and cardiovascular disease. *Taiwan J Obstet Gynecol* 2006;45:287–93.