

# PRENATAL SONOGRAPHIC FEATURES OF FETUSES IN TRISOMY 13 PREGNANCIES (I)

Chih-Ping Chen<sup>1,2,3,4,5,6\*</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, Mackay Memorial Hospital, <sup>2</sup>Department of Medical Research, Mackay Memorial Hospital, Taipei, <sup>3</sup>Department of Biotechnology, Asia University, <sup>4</sup>School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, and <sup>5</sup>Institute of Clinical and Community Health Nursing, and <sup>6</sup>Department of Obstetrics and Gynecology, National Yang-Ming University, Taipei, Taiwan.

## SUMMARY

Prenatal ultrasound is a powerful tool for detecting structural abnormalities in fetuses in trisomy 13 pregnancies. This article provides a comprehensive review of the prenatal sonographic features of trisomy 13, including the major structural abnormalities observed during the first trimester (omphalocele, holoprosencephaly, megacystis and congenital heart defects), the frequencies of second- and third-trimester sonographic features reported in previous studies, and the subtle sonographic findings observed during the second trimester (echogenic intra-cardiac foci, echogenic bowel, single umbilical artery, choroid plexus cysts and intrauterine growth restriction). [*Taiwan J Obstet Gynecol* 2009;48(3):210-217]

**Key Words:** congenital malformations, prenatal diagnosis, trisomy 13, ultrasound

## Introduction

Trisomy 13 syndrome or Patau syndrome is the third most common autosomal chromosomal trisomy after trisomy 21 and trisomy 18. Goldstein and Nielsen [1] reported the incidence of trisomy 13 to be 0.053 per 1,000 live births. In a 13-year incidence study of chromosomal abnormalities in Arhus, Denmark, Nielsen and Wohler [2] found three cases of trisomy 13 among 34,910 newborn children, or an incidence of 0.09 per 1,000 live births, while Wyllie et al [3] found an incidence of 0.077 per 1,000 live births. In a literature survey, Benn and Hsu [4] found three cases of trisomy 13 among 68,159 live births, or an incidence of 0.04 per 1,000. Prenatal ultrasound is a powerful tool for the detection of structural abnormalities in trisomy 13 fetuses. The Table summarizes these structural abnormalities, the markers or subtle findings, and the

placental abnormalities that may present on ultrasound throughout such pregnancies [5]. The placental abnormalities encountered in trisomy 13 pregnancies include reduced placental vascularization and smaller placental volume in the first trimester, and a partial molar appearance and placental mesenchymal dysplasia in the second trimester [6]. To date, at least six studies of fetuses with trisomy 13 have been reported [7-12]. This article provides a comprehensive review of sonographic screening for structural abnormalities in trisomy 13 fetuses during the first trimester and the subtle sonographic findings in the second trimester.

## Sonographic Screening for Structural Abnormalities Associated with Trisomy 13 in the First Trimester

Papageorgiou et al [13] found that 92 of 181 (50.8%) trisomy 13 fetuses who underwent ultrasound examination during the first trimester had omphalocele, holoprosencephaly (HPE) and/or megacystis. Watson et al [12] found that six of eight fetuses with trisomy 13 identified at 10-13 weeks of gestation had abnormal sonographic findings, including cystic hygroma in four



\*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.  
E-mail: cpc\_mmh@yahoo.com  
Accepted: August 12, 2009

**Table.** Prenatal ultrasound features of trisomy 13**Fetus**

- Structural abnormalities
  - First trimester: holoprosencephaly, omphalocele, megacystis, and congenital heart defects
  - Second and third trimesters:
    - Brain: holoprosencephaly, brachycephaly, microcephaly, Dandy-Walker complex, posterior fossa abnormalities, ventriculomegaly, and spina bifida
    - Face: cyclopia, ethmocephaly, cebocephaly, premaxillary agenesis, cleft lip/palate, micrognathia, and hypoplastic nasal bone
    - Neck: cystic hygroma and nuchal edema
    - Heart: ventricular septal defects, atrial septal defects, patent ductus arteriosus, double-outlet right ventricle, hypoplastic left ventricle, mitral or aortic atresia, pulmonary stenosis, and anomalous venous return
    - Chest: diaphragmatic hernia
    - Abdominal wall: omphalocele
    - Urinary tract: renal cystic dysplasia, multicystic kidneys, enlarged and echogenic kidneys, hydronephrosis, and ureteral obstruction and duplication
    - Limbs: postaxial polydactyly of the hands and feet, clenched or overlapping digits, prominent calcaneus, and rocker-bottom feet
- Markers or subtle findings
  - First trimester: increased nuchal translucency thickness, fetal tachycardia, absent nasal bone, abnormal ductus venosus flow, smaller crown-rump length, smaller trunk and head volume, increased frontomaxillary facial angle in the presence of holoprosencephaly, and smaller gestational sac volume
  - Second trimester: echogenic intracardiac foci, echogenic bowel, pyelectasis, single umbilical artery, choroid plexus cysts, mild ventriculomegaly, and intrauterine growth restriction

**Placenta**

- First trimester: reduced placental vascularization and smaller placental volume
- Second trimester: partial molar appearance and placental mesenchymal dysplasia

cases, HPE in one case and irregularly large yolk sac in one case. Bronshtein et al [14] suggested that detailed first-trimester ultrasound examination of fetuses with abnormal nuchal translucency (NT) could shorten the parental decision-making process and spare parents a prolonged period of diagnostic uncertainty and anxiety, particularly when a structural anomaly was clearly diagnosed. In a study of 23 first-trimester fetuses with increased NT thickness  $\geq 3.5$  mm, Bronshtein et al [14] found severe anomalies in eight, mild anomalies in six, trisomy 13 in two, and trisomy 21 in one. In their study, one trisomy 13 fetus with NT thickness of 5.2 mm at 11<sup>+4</sup> weeks of gestation had cleft lip and palate, congenital diaphragmatic hernia, HPE, hypoplastic left heart, micrognathia, and truncus arteriosus. Another trisomy 13 fetus had NT thickness of 6.0 mm at 12 weeks of gestation, and had cleft lip and palate, HPE and proboscis, while one trisomy 21 fetus with NT thickness of 4.4 mm at 13<sup>+3</sup> weeks of gestation had hydronephrosis.

**Omphalocele**

Papageorgiou et al [13] found omphalocele in 28.2% (51/181) of trisomy 13 fetuses in the first trimester. Frequencies of omphalocele in trisomy 13 fetuses in the second and third trimesters of 3.6% (1/28) [11],

11.1% (2/18) [15], 13.3% (2/15) [10], 15.2% (5/33) [8], 22.6% (7/31) [7] and 29.4% (5/17) [16] have been reported. Chromosomal abnormalities have been reported in 10–12% of neonates with omphalocele and in 30% of fetuses with omphalocele [17–22]. When the diagnosis is made in early pregnancy, the incidence of aneuploidy can increase to 61.1% (11/18) at 12–16 weeks of gestation [23] and 66.7% (12/18) at 11–14 weeks of gestation [20]. The chromosomal abnormalities in 11 fetal omphaloceles reported by Blazer et al [23] were trisomy 18 ( $n=5$ ), trisomy 21 ( $n=2$ ), triploidy ( $n=2$ ), trisomy 13 ( $n=1$ ) and 45,X ( $n=1$ ). The reported chromosomal abnormalities in the 12 fetal omphaloceles reported by Snijders et al [20] were trisomy 18 ( $n=10$ ), trisomy 13 ( $n=1$ ) and triploidy ( $n=1$ ). In a meta-analysis of chromosomal abnormalities associated with omphalocele, Chen [24] found that 36.2% (415/1,148) of fetuses with prenatally detected omphalocele had chromosomal abnormalities, including trisomy 18 in 66.7% (277/415), trisomy 13 in 17.3% (72/415) and trisomy 21 in 6.3% (26/415).

**Holoprosencephaly**

Papageorgiou et al [13] found HPE in 26.5% (48/181) of trisomy 13 fetuses in the first trimester. Frequencies of

16.7% (3/18) [15], 17.9% (5/28) [11], 35.5% (11/31) [7], 39.4% (13/33) [8] and 46.7% (7/15) [10] have been reported for HPE in trisomy 13 fetuses in the second and third trimesters. Blaas et al [25] found chromosomal abnormalities in 36.7% (11/30) of fetuses with prenatally detected HPE, and trisomy 13 in 45.5% (5/11). Reported frequencies of chromosomal abnormalities in fetuses with HPE in other studies have included 28.9% (11/38) [26], 33.3% (4/12) [27], 34.0% (17/50) [28], 38.8% (100/258) [29], 41.3% (50/121) [30], 42.9% (6/14) [31] and 57.6% (34/59) [32]. In a meta-analysis of chromosomal abnormalities associated with HPE, Blaas et al [25] suggested that the frequency was 34.7% (92/265) or more. Using modern, high-frequency transvaginal transducers, HPE can be diagnosed in the first trimester [33–48]. Sepulveda et al [41] suggested that failure to identify the “butterfly” sign (a cross-sectional view of the fetal brain, including visualization of both choroid plexuses) was a warning sign for HPE in the first trimester. Kim et al [44] and Timor-Tritsch et al [45] successfully used three-dimensional inversion-rendering mode to evaluate early fetal ventricles in HPE in the first trimester.

### *Megacystis*

Papageorgiou et al [13] found megacystis in 11.6% (21/181) of trisomy 13 fetuses in the first trimester. Fetal megacystis at 10–14 weeks of gestation is defined by a longitudinal bladder diameter of  $\geq 7$  mm [49,50]. Sebire et al [49] detected fetal megacystis in 15 of 24,492 (1/1,633) pregnancies and chromosomal abnormalities in three of the 15 cases (20%). These included trisomy 18 ( $n=1$ ), trisomy 13 ( $n=1$ ) and unbalanced translocation involving chromosomes 14 and 20 ( $n=1$ ). They also found that in the majority of chromosomally normal fetuses with mild or moderate enlargement of the bladder (longitudinal bladder diameter, 8–12 mm), there was spontaneous resolution by 20 weeks of gestation with no obvious adverse effects. They also found progressive obstructive uropathy in all fetuses with severe megacystis (longitudinal bladder diameter,  $\geq 17$  mm). Favre et al [51] reported chromosomal abnormalities in four of 15 fetuses (26.7%) at 11–15 weeks of gestation, including trisomy 13 ( $n=2$ ), trisomy 18 ( $n=1$ ) and trisomy 21 ( $n=1$ ). Liao et al [50] found that 30 of 145 fetuses (20.7%) with megacystis at 10–14 weeks of gestation had chromosomal abnormalities, including trisomy 13 ( $n=17$ ), trisomy 18 ( $n=7$ ), trisomy 21 ( $n=2$ ), triploidy ( $n=1$ ), trisomy 4 ( $n=1$ ), mosaic trisomy 15 ( $n=1$ ), and unbalanced translocation involving chromosomes 14 and 20 ( $n=1$ ). They also expected a significantly higher number of aneuploid fetuses with trisomy 13 and trisomy 18, but not trisomy 21 in cases of megacystis. Liao et al [50] demonstrated that in fetuses

with megacystis and a longitudinal bladder diameter of 7–15 mm, the risk of aneuploidy was 23.6% (26/110), compared with a risk of 11.4% (4/35) in those with a longitudinal bladder diameter of  $> 15$  mm. Liao et al [50] additionally found that increased NT thickness was present in about 75% of aneuploid and about 30% of euploid fetuses with megacystis. About 90% of euploid fetuses with a longitudinal bladder diameter of 7–15 mm had spontaneous resolution, but all fetuses with a longitudinal bladder diameter  $> 15$  mm had progressive obstructive uropathy. Boissier et al [52] reported that three of 12 (25%) fetuses with fetal megacystis in the first trimester had chromosomal abnormalities, including trisomy 18 ( $n=2$ ) and trisomy 21 ( $n=1$ ). Sepulveda [53] concluded that early detection of megacystis should include karyotyping and follow-up scans, and that vesicocentesis should be managed promptly in cases with progressive enlargement of the fetal bladder, while serial vesicocentesis with vesicoamniotic shunting or cystoscopic procedures should be offered as an alternative to termination of pregnancy, if the condition did not resolve.

### *Congenital heart defects (CHD)*

CHD have been found in more than 80% of fetuses with trisomy 13 [54,55]. The commonest CHDs in trisomy 13 fetuses include ventricular septal defects, atrial septal defects, patent ductus arteriosus, overriding aorta, dextroposition, hypoplastic aorta, atretic mitral and/or aortic valves, pulmonary stenosis, and anomalous pulmonary venous return [55]. Frequencies of CHD in trisomy 13 fetuses in the second and third trimesters have been reported as 33.3% (5/15) [42], 45.2% (14/31) [7], 48.5% (16/33) [8], 50.0% (9/18) [15], 53.6% (15/28) [11], and 70.6% (12/17) [16]. Becker and Wegner [56] found that the overall detection rate for major CHD on the 11–13<sup>+</sup>6-week scan was 84.2% (32/38), with 37.5% (3/8) for NT  $< 2.5$  mm and 96.7% (29/30) for NT  $\geq 2.5$  mm. In a study of pregnancies with risk factors for fetal CHD, such as NT  $\geq 4$  mm, a first-degree relative with CHD, or suspicion of CHD or extracardiac abnormality on the 10–14-week scan, Huggon et al [57] found chromosomal abnormalities in 31.5% (70/222) of fetuses with a normal heart on echocardiography (trisomy 21 [ $n=47$ ], trisomy 18 [ $n=9$ ], trisomy 13 [ $n=4$ ], 45,X [ $n=3$ ], and others [ $n=7$ ]). In contrast, chromosomal abnormalities were present in 69.2% (18/26) of fetuses with atrioventricular septal defect (trisomy 21 [ $n=14$ ], trisomy 18 [ $n=1$ ], trisomy 13 [ $n=2$ ], and others [ $n=1$ ]), in 71.4% (5/7) of fetuses with hypoplastic left heart syndrome (trisomy 18 [ $n=1$ ] and 45,X [ $n=4$ ]), in 80.0% (8/10) of fetuses with ventricular septal defect (trisomy 21 [ $n=2$ ], trisomy

18 [ $n=5$ ], and trisomy 13 [ $n=1$ ]), in 92.6% (25/27) of fetuses with disproportion (trisomy 21 [ $n=3$ ], trisomy 18 [ $n=5$ ], trisomy 13 [ $n=4$ ], and 45,X [ $n=13$ ]), and in 94.7% (18/19) of fetuses with isolated tricuspid valve regurgitation (trisomy 21 [ $n=14$ ], trisomy 18 [ $n=2$ ], 45,X [ $n=1$ ], and others [ $n=1$ ]). In the study by Huggon et al [57], seven of the 11 trisomy 13 fetuses (63.6%) had confirmed cardiac abnormalities on echocardiography.

## Echogenic Intracardiac Focus (EIF)

EIF is the subjective sonographic finding of an echogenic spot over the capillary muscle, and its detection depends on the resolution of the ultrasound equipment, the operator's experience, and the fetal position. EIF is a common sonographic finding in the second trimester and is observed in 3–4% of normal fetuses [58,59]. Roberts and Genest [60] reported that mineralization of the cardiac papillary muscle was observed in 15.9% (20/126) of fetuses with trisomy 21 and 39.1% (9/23) of fetuses with trisomy 13, compared with only 2% of normal fetuses. Lehman et al [8] reported that EIF was found in 38.9% (7/18) of fetuses with trisomy 13 at 12–20 weeks of gestation, and Bromley et al [61] found EIF in 18.2% (4/22) of fetuses with trisomy 21, compared with 4.7% (62/1,312) of normal fetuses. However, racial differences exist in the frequency of sonographically identified EIF in second-trimester fetal hearts [62]. Shipp et al [62] found incidences of EIF of 30.4%, 5.9%, 10.5% and 11.1% for Asian, black, white and unknown women, suggesting that EIF is a normal condition that occurs in about one-third of fetuses of Asian women. Frequencies of 3.7% (2/54) [12], 14.3% (4/28) [11], 20% (3/15) [10] and 30.3% (10/33) [8] have been reported in fetuses with trisomy 13.

## Echogenic Bowel

Papp et al [11] found echogenic bowel in 17.9% (5/28) of fetuses with trisomy 13. This condition occurs in approximately 1 per 200 mid-trimester fetuses [63,64]. It may be a normal condition during the second trimester, but has also been associated with intra-amniotic hemorrhage, severe uteroplacental insufficiency, cystic fibrosis, chromosomal abnormalities, and *in utero* infection with cytomegalovirus and toxoplasmosis [63,65]. In a study of karyotyped fetuses with echogenic bowel, after exclusion of fetuses with major anomalies, Al-Kouatly et al [65] found that 3.7% (5/136) had chromosomal abnormalities, including trisomy 21 ( $n=4$ ) and trisomy 18 ( $n=1$ ). Bromley et al [66] reported that echogenic

bowel was found in 12.5% (6/48) of fetuses with trisomy 21, and that the risk of trisomy 21 in fetuses with isolated echogenic bowel was 1.4% (71/5,105). Echogenic bowel is associated with a six- to seven-fold increase in the risk of Down syndrome, or a risk in the range of 1–2% [67,68]. In a study of 50 fetuses with echogenic bowel, Bromley et al [66] found that 16% (8/50) of fetuses had chromosomal abnormalities, including trisomy 21 ( $n=6$ ), trisomy 13 ( $n=1$ ) and Turner syndrome ( $n=1$ ), and that all aneuploid cases had additional anomalies. Snijders et al [63] observed that echogenic bowel was most commonly associated with placental insufficiency and intrauterine growth restriction (IUGR). In their study of 250 fetuses with echogenic bowel, the frequencies of chromosomal abnormalities in fetuses with IUGR, minor abnormalities and major abnormalities were 0% (0/91), 2.5% (3/122) (trisomy 21 [ $n=1$ ], trisomy 18 [ $n=1$ ], and triploidy [ $n=1$ ]), and 35.8% (24/67) (trisomy 21 [ $n=6$ ], trisomy 18 [ $n=6$ ], trisomy 13 [ $n=6$ ], triploidy [ $n=2$ ] and other rearrangements [ $n=4$ ]), respectively.

## Single Umbilical Artery (SUA)

SUA has been found in 0.2–1.9% of deliveries [69–73]. Granese et al [74] reported a prevalence of 0.48% (61/12,672) for SUA on prenatal ultrasound at 16–23 weeks of gestation. SUA can be associated with congenital malformations of all major organ systems and with chromosomal abnormalities [75–80]. Prucka et al [81] reported that SUA existed in 2% (97/4,846) of pathologic specimens and that fetuses with SUA had significantly more chromosomal abnormalities (10.3% vs. 1.0%) and other congenital anomalies (27% vs. 8%). Martínez-Frías et al [82] found that the most commonly associated malformations were bilateral renal agenesis and imperforate anus, followed by unilateral renal agenesis and vertebral defects. SUA has been found in more than 50% of fetuses with trisomy 18 and 10–50% of fetuses with trisomy 13 [67].

In the first trimester, visualization of the umbilical arteries on either side of the bladder and continuous with the umbilical cord insertion to the fetus can be obtained by color flow mapping in an oblique, transverse section of the fetal abdomen including the umbilicus and the fetal bladder [83]. In a study of 717 fetuses immediately before chorionic villus sampling at 11–14 weeks of gestation, Rembouskos et al [83] found SUA in 3.3% (21/634) of fetuses with a normal karyotype, 11.4% (5/44) with trisomy 21, 77.8% (14/18) with trisomy 18 and 9.5% (2/21) with other chromosomal abnormalities. The authors also found that at 11–14

weeks of gestation, the most commonly associated major structural defects were omphalocele and megacystis, and the most common chromosomal abnormality was trisomy 18 in fetuses with SUA.

In a meta-analysis of 936 fetuses with SUA diagnosed in the second and third trimesters, Rembouskos et al [83] found that 79 fetuses (8.4%) had chromosomal abnormalities, including trisomy 18 ( $n=33$ ), trisomy 13 ( $n=15$ ), trisomy 21 ( $n=5$ ), triploidy ( $n=9$ ) and other rearrangements ( $n=17$ ). In a study of fetuses with SUA diagnosed in the second trimester, Lubusky et al [84] found SUA in 4.8% (102/2,147) of cases, including 12.8% (5/39) with trisomy 21, 50% (8/16) with trisomy 18, 25% (1/4) with trisomy 13 and 7.25% (5/69) with other rearrangements. SUA has been reported to occur in 9.3% (5/54) [12], 13.3% (2/15) [10] and 24.2% (8/33) [8] of fetuses with trisomy 13.

## Choroid Plexus Cysts

Choroid plexus cysts (CPCs) occur in approximately 1% of karyotypically normal fetuses and about 50% of fetuses with trisomy 18 [85]. The reported frequency of CPCs in the normal population ranges from 0.3% to 3.6% [86–89]. CPCs are a normal condition during the second trimester and resolve by 26–28 weeks of gestation in more than 90% of cases, with no known pathologic association [63]. Sullivan et al [90] suggested that normal results of maternal serum biochemical tests and an absence of additional fetal anomalies on prenatal ultrasound could reliably exclude any underlying chromosomal abnormality. In a meta-analysis of 1,806 fetuses with CPCs, Snijders et al [63] reported a mean frequency of chromosomal abnormalities of 8%, including trisomy 18 ( $n=121$ ), trisomy 21 ( $n=18$ ) and other rearrangements ( $n=11$ ), with a frequency of 1% for cases with isolated CPCs and 46% for those with multiple anomalies. They also found CPCs in 47% of fetuses with trisomy 18 ( $n=137$ ), 8% with trisomy 21 ( $n=155$ ) and 2% with trisomy 13 ( $n=54$ ). Yoder et al [91] found a likelihood ratio of 13.8 (95% confidence interval, CI, 7.72–25.14;  $p<0.001$ ) for trisomy 18 and 1.87 (95% CI, 0.78–4.46;  $p=0.16$ ) for trisomy 21 in second-trimester fetuses with isolated CPCs. Frequencies of 3.2% (1/31) [7], 6.7% (1/15) [10] and 10.7% (3/28) [11] have been reported for CPCs in fetuses with trisomy 13.

## IUGR

The frequency of chromosomal abnormalities in neonates with small gestational age is  $<10\%$  [92–95].

However, early IUGR is a common prenatal feature of major chromosomal abnormalities, particularly triploidy, trisomy 18 and trisomy 13 [95]. In a study of 458 fetuses with IUGR at 17–39 weeks of gestation, Snijders et al [95] found chromosomal abnormalities in 19.4% (89/458) of cases, including triploidy ( $n=36$ ), trisomy 18 ( $n=32$ ), trisomy 21 ( $n=8$ ), trisomy 13 ( $n=5$ ), trisomy 22 ( $n=1$ ) and other autosomal deletions or duplications ( $n=7$ ). The frequency of chromosomal abnormalities varied with gestational age at presentation (38% at 18–25 weeks, 10% at 26–33 weeks, and 15% at 34–41 weeks), the presence of fetal anomalies (40% with fetal anomalies and 3% without fetal anomalies), amniotic fluid volume (95% with increased volume, 32% with normal volume, 10% with reduced volume, and 5% with absent volume), and Doppler findings (44% with absence of notch [–NT] and presence of end-diastolic frequencies [+EDF], 24% with –NT/–EDF, 12% with +NT/+EDF, and 8% with +NT/–EDF).

Triploid fetuses have early-onset asymmetrical IUGR with macrocephaly and increased head-to-abdominal circumference ratio, whereas fetuses with trisomy 18 and trisomy 13 have symmetrical IUGR before 30 weeks' gestation [63]. Dicke and Crane [96] found that mid-trimester IUGR was evident in 43% of fetuses with trisomy 13 ( $n=9$ ) and 59% of fetuses with trisomy 18 ( $n=22$ ), but not in fetuses with trisomy 21 ( $n=43$ ). Snijders et al [63] found small gestational age in 100% of fetuses with triploidy ( $n=50$ ), 74% with trisomy 18 ( $n=137$ ), 61% with trisomy 13 ( $n=54$ ), 55% with Turner syndrome ( $n=54$ ), and 20% with trisomy 21 ( $n=155$ ). Jauniaux et al [97] reported that 62.5% (10/16) of fetuses with triploidy had fetal crown-rump length below the fifth centile at 10–14 weeks of gestation. Bahado-Singh et al [98] reported that, in the first trimester, fetuses with trisomy 18 and trisomy 13 were growth restricted, with significant shortening of crown-rump length, but that fetuses with trisomy 21 were not associated with early-onset IUGR.

Frequencies of 26.7% (4/15) [10], 48.4% (15/31) [7] and 48.5% (16/33) [8] have been reported for IUGR in fetuses with trisomy 13.

## References

1. Goldstein H, Nielsen KG. Rates and survival of individuals with trisomy 13 and 18: data from a 10-year period in Denmark. *Clin Genet* 1988;34:366–72.
2. Nielsen J, Wohlert M. Chromosome abnormalities found among 34,910 newborn children: results from a 13-year incidence study in Arhus, Denmark. *Hum Genet* 1991;87: 81–3.
3. Wyllie JP, Wright MJ, Burn J, Hunter S. Natural history of trisomy 13. *Arch Dis Child* 1994;71:343–5.



4. Benn PA, Hsu LYF. Prenatal diagnosis of chromosomal abnormalities through amniocentesis. In: Milunsky A, ed. *Genetic Disorders and the Fetus: Diagnosis, Prevention, and Treatment*, 5<sup>th</sup> edition. Baltimore: Johns Hopkins University Press, 2004:214–96.
5. Chen CP, Chien SC. Prenatal sonographic features of trisomy 13. *J Med Ultrasound* 2007;15:58–66.
6. Chen CP. Placental abnormalities and preeclampsia in trisomy 13 pregnancies. *Taiwan J Obstet Gynecol* 2009;48:3–8.
7. Nicolaides KH, Snijders RJM, Gosden CM, Berry C, Campbell S. Ultrasonographically detectable markers of fetal chromosomal abnormalities. *Lancet* 1992;340:704–7.
8. Lehman CD, Nyberg DA, Winter TC 3rd, Kapur RP, Resta RG, Luthy DA. Trisomy 13 syndrome: prenatal US findings in a review of 33 cases. *Radiology* 1995;194:217–22.
9. De Vigan C, Baena N, Cariati E, Clementi M, Stoll C; EUROSCAN Working Group. Contribution of ultrasonographic examination to the prenatal detection of chromosomal abnormalities in 19 centres across Europe. *Ann Genet* 2001;44:209–17.
10. Tongsong T, Sirichotiyakul S, Wanapirak C, Chanprapaph P. Sonographic features of trisomy 13 at midpregnancy. *Int J Gynaecol Obstet* 2002;76:143–8.
11. Papp C, Beke A, Ban Z, Szigeti Z, Toth-Pal E, Papp Z. Prenatal diagnosis of trisomy 13: analysis of 28 cases. *J Ultrasound Med* 2006;25:429–35.
12. Watson WJ, Miller RC, Wax JR, Hansen WF, Yamamura Y, Polzin WJ. Sonographic detection of trisomy 13 in the first and second trimesters of pregnancy. *J Ultrasound Med* 2007;26:1209–14.
13. Papageorgiou AT, Avgidou K, Spencer K, Nix B, Nicolaides KH. Sonographic screening for trisomy 13 at 11 to 13<sup>+</sup>6 weeks of gestation. *Am J Obstet Gynecol* 2006;194:397–401.
14. Bronshtein M, Zimmer EZ, Blazer S. The utility of detailed first trimester ultrasound examination in abnormal fetal nuchal translucency. *Prenat Diagn* 2008;28:1037–41.
15. Picklesimer AH, Moise KJ Jr, Wolfe HM. The impact of gestational age on the sonographic detection of aneuploidy. *Am J Obstet Gynecol* 2005;193:1243–7.
16. Wladimiroff JW, Bhaggoe WR, Kristelijn M, Cohen-Overbeek TE, Den Hollander NS, Brandenburg H, Los FJ. Sonographically determined anomalies and outcome in 170 chromosomally abnormal fetuses. *Prenat Diagn* 1995;15:431–8.
17. Nicolaides KH, Snijders RJ, Cheng HH, Gosden C. Fetal gastro-intestinal and abdominal wall defects: associated malformations and chromosomal abnormalities. *Fetal Diagn Ther* 1992;7:102–15.
18. Calzolari E, Volpato S, Bianchi F, et al. Omphalocele and gastroschisis: a collaborative study of five Italian congenital malformation registries. *Teratology* 1993;47:47–55.
19. Paidas MJ, Crombleholme TM, Robertson FM. Prenatal diagnosis and management of the fetus with an abdominal wall defect. *Semin Perinatol* 1994;18:196–214.
20. Snijders RJM, Brizot ML, Faria M, Nicolaides KH. Fetal exomphalos at 11 to 14 weeks of gestation. *J Ultrasound Med* 1995;14:569–74.
21. Snijders RJM, Sebire NJ, Souka A, Santiago C, Nicolaides KH. Fetal exomphalos and chromosomal defects: relationship to maternal age and gestation. *Ultrasound Obstet Gynecol* 1995;6:250–5.
22. St-Vil D, Shaw KS, Lallier M, Yazbeck S, Di Lorenzo M, Grignon A, Blanchard H. Chromosomal anomalies in newborns with omphalocele. *J Pediatr Surg* 1996;31:831–4.
23. Blazer S, Zimmer EZ, Gover A, Bronshtein M. Fetal omphalocele detected early in pregnancy: associated anomalies and outcomes. *Radiology* 2004;232:191–5.
24. Chen CP. Chromosomal abnormalities associated with omphalocele. *Taiwan J Obstet Gynecol* 2007;46:1–8.
25. Blaas HGK, Eriksson AG, Salvesen KÅ, Isaksen CV, Christensen B, Møllerlökken G, Eik-Nes SH. Brains and faces in holoprosencephaly: pre- and postnatal description of 30 cases. *Ultrasound Obstet Gynecol* 2002;19:24–38.
26. Berry SM, Gosden C, Snijders RJM, Nicolaides KH. Fetal holoprosencephaly: associated malformations and chromosomal defects. *Fetal Diagn Ther* 1990;5:92–9.
27. Parant O, Sarramon MF, Delisle MB, Fournié A. Antenatal diagnosis of holoprosencephaly: a series of twelve cases. *J Gynecol Obstet Biol Reprod (Paris)* 1997;26:687–96. [In French]
28. Whiteford ML, Tolmie JL. Holoprosencephaly in the west of Scotland 1975–1994. *J Med Genet* 1996;33:578–84.
29. Odent S, Le Marec B, Munnich A, Le Merrer M, Bonaiti-Pellié C. Segregation analysis in nonsyndromic holoprosencephaly. *Am J Med Genet* 1998;77:139–43.
30. Croen LA, Shaw GM, Lammer EJ. Holoprosencephaly: epidemiologic and clinical characteristics of a California population. *Am J Med Genet* 1996;64:465–72.
31. Nyberg DA, Mack LA, Bronstein A, Hirsch J, Pagon RA. Holoprosencephaly: prenatal sonographic diagnosis. *AJR Am J Roentgenol* 1987;149:1051–8.
32. Chen CP, Chern SR, Lin CJ, Lee CC, Wang W, Tzen CY. A comparison of maternal age, sex ratio and associated anomalies among numerically aneuploid, structurally aneuploid and euploid holoprosencephaly. *Genet Couns* 2005;16:49–57.
33. Hamada H, Oki A, Tsunoda H, Kubo T. Prenatal diagnosis of holoprosencephaly by transvaginal ultrasonography in the first trimester. *Asia Oceania J Obstet Gynaecol* 1992;18:125–9.
34. Sakala EP, Gaio KL. Fundal uterine leiomyoma obscuring first-trimester transabdominal sonographic diagnosis of fetal holoprosencephaly: a case report. *J Reprod Med* 1993;38:400–2.
35. Gembruch U, Baschat AA, Reusche E, Wallner SJ, Greiwe M. First trimester diagnosis of holoprosencephaly with a Dandy-Walker malformation by transvaginal ultrasonography. *J Ultrasound Med* 1995;14:619–22.
36. van Zalen-Sprock R, van Vugt JM, van der Harten HJ, Nieuwint AW, van Geijn HP. First trimester diagnosis of cyclopia and holoprosencephaly. *J Ultrasound Med* 1995;14:631–3.
37. Tongsong T, Wanapirak C, Chanprapaph P, Sirianguk S. First trimester sonographic diagnosis of holoprosencephaly. *Int J Gynaecol Obstet* 1999;66:165–9.
38. Turner CD, Silva S, Jeanty P. Prenatal diagnosis of alobar holoprosencephaly at 10 weeks of gestation. *Ultrasound Obstet Gynecol* 1999;13:360–2.
39. Wong HS, Lam YH, Tang MHY, Cheung LWK, Ng LKL, Yan KW. First-trimester ultrasound diagnosis of holoprosencephaly: three case reports. *Ultrasound Obstet Gynecol* 1999;13:356–9.
40. Hsu TY, Chang SY, Ou CY, Chen ZH, Tsai WL, Chang MS, Soong YK. First trimester diagnosis of holoprosencephaly and cyclopia with triploidy by transvaginal three-dimensional ultrasonography. *Eur J Obstet Gynecol Reprod Biol* 2001;96:235–7.

41. Sepulveda W, Dezerega V, Be C. First-trimester sonographic diagnosis of holoprosencephaly: value of the "butterfly" sign. *J Ultrasound Med* 2004;23:761-5.
42. Tongsong T, Khunamornpong S, Wanapirak C, Sirichotiyakul S. Prenatal sonographic diagnosis of truncus arteriosus associated with holoprosencephaly. *J Clin Ultrasound* 2005; 33:193-6.
43. Tonni G, Centini G. Three-dimensional first-trimester transvaginal diagnosis of alobar holoprosencephaly associated with omphalocele in a 46,XX fetus. *Am J Perinatol* 2006;23: 67-9.
44. Kim MS, Jeanty P, Turner C, Benoit B. Three-dimensional sonographic evaluations of embryonic brain development. *J Ultrasound Med* 2008;27:119-24.
45. Timor-Tritsch IE, Monteagudo A, Santos R. Three-dimensional inversion rendering in the first- and early second-trimester fetal brain: its use in holoprosencephaly. *Ultrasound Obstet Gynecol* 2008;32:744-50.
46. Tonni G, Ventura A, Centini G, De Felice C. First trimester three-dimensional transvaginal imaging of alobar holoprosencephaly associated with proboscis and hypotelorism (ethmocephaly) in a 46,XX fetus. *Congenit Anom (Kyoto)* 2008;48:51-5.
47. Dane B, Dane C, Aksoy F, Yayla M. Semilobar holoprosencephaly with associated cyclopia and radial aplasia: first trimester diagnosis by means of integrating 2D-3D ultrasound. *Arch Gynecol Obstet* 2009;280:647-51.
48. Solomon BD, Potocki L, Oyer CE, Muenke M. Holoprosencephaly in an 8.5-week triploidy gestation. *Clin Dysmorphol* 2009;18:166-7.
49. Sebire NJ, Von Kaisenberg C, Rubio C, Snijders RJM, Nicolaides KH. Fetal megacystis at 10-14 weeks of gestation. *Ultrasound Obstet Gynecol* 1996;8:387-90.
50. Liao AW, Sebire NJ, Geerts L, Cicero S, Nicolaides KH. Megacystis at 10-14 weeks of gestation: chromosomal defects and outcome according to bladder length. *Ultrasound Obstet Gynecol* 2003;21:338-41.
51. Favre R, Kohler M, Gasser B, Muller F, Nisand I. Early fetal megacystis between 11 and 15 weeks of gestation. *Ultrasound Obstet Gynecol* 1999;14:402-6.
52. Boissier K, Varlet MN, Chauleur C, et al. Early fetal megacystis at first trimester: a six-year retrospective study. *Gynecol Obstet Fertil* 2009;37:115-24. [In French]
53. Sepulveda W. Megacystis in the first trimester. *Prenat Diagn* 2004;24:144-9.
54. Nora JJ, Nora AH. The evolution of specific genetic and environmental counseling in congenital heart diseases. *Circulation* 1978;57:205-13.
55. Jones KL. Trisomy 13 syndrome. In: Jones KL, ed. *Smith's Recognizable Patterns of Human Malformation*. Philadelphia: Elsevier Saunders, 2006:18-21.
56. Becker R, Wegner RD. Detailed screening for fetal anomalies and cardiac defects at the 11-13-week scan. *Ultrasound Obstet Gynecol* 2006;27:613-8.
57. Huggon IC, Ghi T, Cook AC, Zosmer N, Allan LD, Nicolaides KH. Fetal cardiac abnormalities identified prior to 14 weeks' gestation. *Ultrasound Obstet Gynecol* 2002;20:22-9.
58. Schechter AG, Fakhry J, Shapiro LR, Gewitz MH. In utero thickening of the chordae tendinae: a cause of intracardiac echogenic foci. *J Ultrasound Med* 1987;6:691-5.
59. Levy DW, Mintz MC. The left ventricular echogenic focus: a normal finding. *AJR Am J Roentgenol* 1988;150:85-6.
60. Roberts DJ, Genest D. Cardiac histologic pathology characteristic of trisomies 13 and 21. *Hum Pathol* 1992;23:1130-40.
61. Bromley B, Lieberman E, Shipp TD, Richardson M, Benacerraf BR. Significance of an echogenic intracardiac focus in fetuses at high and low risk for aneuploidy. *J Ultrasound Med* 1998; 17:127-31.
62. Shipp TD, Bromley B, Lieberman E, Benacerraf BR. The frequency of the detection of fetal echogenic intracardiac foci with respect to maternal race. *Ultrasound Obstet Gynecol* 2000;15:460-2.
63. Snijders RJM, Farrias M, von Kaisenberg C, Nicolaides KH. Fetal abnormalities. In: Snijders RJM, Nicolaides KH, eds. *Ultrasound Markers for Fetal Chromosomal Defects*. New York: Parthenon Publishing Group, 1996:1-62.
64. Nyberg DA, Souter VL. Chromosomal abnormalities. In: Nyberg DA, McGahan JP, Pretorius DH, Pilu G, eds. *Diagnostic Imaging of Fetal Anomalies*. Philadelphia: Lippincott Williams & Wilkins, 2003:861-906.
65. Al-Kouatly HB, Chasen ST, Streltsoff J, Chervenak FA. The clinical significance of fetal echogenic bowel. *Am J Obstet Gynecol* 2001;185:1035-8.
66. Bromley B, Doubilet P, Frigoletto FD Jr, Krauss C, Estroff JA, Benacerraf BR. Is fetal hyperechoic bowel on second-trimester sonogram an indication for amniocentesis? *Obstet Gynecol* 1994;83:647-51.
67. Nyberg DA, Souter VL, El-Bastawissi A, Young S, Luthardt F, Luthy DA. Isolated sonographic markers for detection of fetal Down syndrome in the second trimester of pregnancy. *J Ultrasound Med* 2001;20:1053-63.
68. Smith-Bindman R, Hosmer W, Feldstein VA, Deeks JJ, Goldberg JD. Second-trimester ultrasound to detect fetuses with Down syndrome: a meta-analysis. *JAMA* 2001; 285:1044-55.
69. Heifetz SA. Single umbilical artery: a statistical analysis of 237 autopsy cases and review of the literature. *Perspect Pediatr Pathol* 1984;8:345-79.
70. Leung AK, Robson WL. Single umbilical artery: a report of 159 cases. *Am J Dis Child* 1989;143:108-11.
71. Lilja M. Infants with single umbilical artery studied in a national registry: general epidemiological characteristics. *Paediatr Perinat Epidemiol* 1991;5:27-36.
72. Jones TB, Sorokin Y, Bhatia R, Zador IE, Bottoms SF. Single umbilical artery: accurate diagnosis? *Am J Obstet Gynecol* 1993;169:538-40.
73. Gornall AS, Kurinczuk JJ, Konje JC. Antenatal detection of a single umbilical artery: does it matter? *Prenat Diagn* 2003; 23:117-23.
74. Granese R, Coco C, Jeanty P. The value of single umbilical artery in the prediction of fetal aneuploidy: findings in 12,672 pregnant women. *Ultrasound Q* 2007;23:117-21.
75. Lenoski EF, Medovy H. Single umbilical artery: incidence, clinical significance and relation to autosomal trisomy. *Can Med Assoc J* 1962;87:1229-31.
76. Monie IW. Genesis of single umbilical artery. *Am J Obstet Gynecol* 1970;108:400-5.
77. Bryan EM, Kohler HG. The missing umbilical artery, I: prospective study based on a maternity unit. *Arch Dis Child* 1974;49:844-52.

78. Byrne J, Blanc WA. Malformations and chromosome anomalies in spontaneously aborted fetuses with single umbilical artery. *Am J Obstet Gynecol* 1985;151:340-2.
79. Chow JS, Benson CB, Doubilet PM. Frequency and nature of structural anomalies in fetuses with single umbilical arteries. *J Ultrasound Med* 1998;17:765-8.
80. Geipel A, Germier U, Welp T, Schwinger E, Gembruch U. Prenatal diagnosis of single umbilical artery: determination of the absent side, associated anomalies, Doppler findings and perinatal outcome. *Ultrasound Obstet Gynecol* 2000;15:114-7.
81. Prucka S, Clemens M, Craven C, McPherson E. Single umbilical artery: what does it mean for the fetus? A case-control analysis of pathologically ascertained cases. *Genet Med* 2004;6:54-7.
82. Martínez-Frías ML, Bermejo E, Rodríguez-Pinilla E, Prieto D; ECEMC Working Group. Does single umbilical artery (SUA) predict any type of congenital defect? Clinical-epidemiological analysis of a large consecutive series of malformed infants. *Am J Med Genet A* 2008;146A:15-25.
83. Rembouskos G, Cicero S, Longo D, Sacchini C, Nicolaides KH. Single umbilical artery at 11-14 weeks' gestation: relation to chromosomal defects. *Ultrasound Obstet Gynecol* 2003;22:567-70.
84. Lubusky M, Dhaifalah I, Prochazka M, Hyjanek J, Mickova I, Vomackova K, Santavy J. Single umbilical artery and its siding in the second trimester of pregnancy: relation to chromosomal defects. *Prenat Diagn* 2007;27:327-31.
85. Snijders RJM, Shawa L, Nicolaides KH. Fetal choroid plexus cysts and trisomy 18: assessment of risk based on ultrasound findings and maternal age. *Prenat Diagn* 1994;14:1119-27.
86. DeRoo TR, Harris RD, Sargent SK, Denholm TA, Crow HC. Fetal choroid plexus cysts: prevalence, clinical significance, and sonographic appearance. *AJR Am J Roentgenol* 1988;151:1179-81.
87. Chan L, Hixson JL, Laifer SA, Marchese SG, Martin JG, Hill LM. A sonographic and karyotypic study of second-trimester fetal choroid plexus cysts. *Obstet Gynecol* 1989;73:703-6.
88. Ostlere SJ, Irving HC, Lilford RJ. A prospective study of the incidence and significance of fetal choroid plexus cysts. *Prenat Diagn* 1989;9:205-11.
89. Chinn DH, Miller EI, Worthy LM, Towers CV. Sonographically detected fetal choroid plexus cysts: frequency and association with aneuploidy. *J Ultrasound Med* 1991;10:255-8.
90. Sullivan A, Giudice T, Vavelidis F, Thiagarajah S. Choroid plexus cysts: is biochemical testing a valuable adjunct to targeted ultrasonography? *Am J Obstet Gynecol* 1999;181:260-5.
91. Yoder PR, Sabbagha RE, Gross SJ, Zelop CM. The second-trimester fetus with isolated choroid plexus cysts: a meta-analysis of risk of trisomies 18 and 21. *Obstet Gynecol* 1999;93:869-72.
92. Chen ATL, Chan YK, Falek A. The effects of chromosomal abnormalities on birth weight in man, II: autosomal defects. *Hum Hered* 1972;22:209-24.
93. Ounsted M, Moar V, Scott WA. Perinatal morbidity and mortality in small-for-dates babies: the relative importance of some maternal factors. *Early Hum Dev* 1981;5:367-75.
94. Khoury MJ, Erickson JD, Cordero JF, McCarthy BJ. Congenital malformations and intrauterine growth retardation: a population study. *Pediatrics* 1988;82:83-90.
95. Snijders RJM, Sherrod C, Gosden CM, Nicolaides KH. Fetal growth retardation: associated malformations and chromosomal abnormalities. *Am J Obstet Gynecol* 1993;168:547-55.
96. Dicke JM, Crane JP. Sonographic recognition of major malformations and aberrant fetal growth in trisomic fetuses. *J Ultrasound Med* 1991;10:433-8.
97. Jauniaux E, Brown R, Snijders RJM, Noble P, Nicolaides KH. Early prenatal diagnosis of triploidy. *Am J Obstet Gynecol* 1997;176:550-4.
98. Bahado-Singh RO, Lynch L, Deren O, Morroti R, Copel JA, Mahoney MJ, Williams J 3rd. First-trimester growth restriction and fetal aneuploidy: The effect of type of aneuploidy and gestational age. *Am J Obstet Gynecol* 1997;176:976-80.