

22Q11.2 MICRODELETION IN A FETUS WITH DOUBLE-OUTLET RIGHT VENTRICLE, PULMONARY STENOSIS AND A VENTRICULAR SEPTAL DEFECT: PRENATAL DIAGNOSIS BY ARRAY COMPARATIVE GENOMIC HYBRIDIZATION

Chih-Ping Chen^{1,2,3,4,5,6*}, Yi-Ning Su⁷, Tung-Yao Chang⁸, Schu-Rern Chern²,
Fuu-Jen Tsai^{4,9,10}, Jonathan Kwei Hwang¹¹, Wayseen Wang^{2,12}

Departments of ¹Obstetrics and Gynecology and ²Medical Research, Mackay Memorial Hospital, Taipei, ³Department of Biotechnology, Asia University, ⁴School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, ⁵Institute of Clinical and Community Health Nursing and ⁶Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, ⁷Department of Medical Genetics, National Taiwan University Hospital, ⁸Taiji Fetal Medicine Center, Taipei, Departments of ⁹Medical Genetics, and ¹⁰Medical Research, China Medical University Hospital, Taichung, Taiwan, ¹¹Loyola Marymount University, Los Angeles, USA, and ¹²Department of Bioengineering, Tatung University, Taipei, Taiwan.

A 30-year-old, primigravid woman was referred to the hospital at 24 weeks' gestation because of fetal cardiac abnormalities detected by sonography. The woman and her husband were nonconsanguineous and healthy, and there was no family history of congenital heart defects or DiGeorge syndrome (DGS). She did not have diabetes mellitus and denied any exposure to teratogenic agents or infectious diseases during this pregnancy. Level II ultrasound revealed a singleton fetus with a fetal biometry equivalent to 24 weeks' gestation, a normal thymus, a normal amount of amniotic fluid, double-outlet right ventricle, pulmonary stenosis, and a ventricular septal defect. Amniocentesis was performed. Bacterial artificial chromosome array comparative genomic hybridization (aCGH) and oligonucleotide aCGH were applied for genetic analysis using cultured amniocytes. Bacterial artificial chromosome aCGH demonstrated a 22q11.21 microdeletion (RP11-690P21 → RP11-1116M14; (Figure 1). Oligonucleotide aCGH demonstrated a 2.6-Mb deletion in 22q11.21 (chromosome 22, 17141308-19792353 bp; Figure 2). G-banded chromosome analysis at the 850-band level revealed a 46,XX karyotype. Fluorescence *in situ*

hybridization (FISH) using dual color DNA probes showed a heterozygous deletion of the DGS/velocardiofacial syndrome (DGS/VCFS) region (Figure 3). FISH chromosome analysis revealed a 46,XX.ish del




Figure 1. Bacterial artificial chromosome array comparative genomic hybridization using CMDX bacterial artificial chromosome array comparative genomic hybridization CA2500 chips (CMDX, Irvine, CA, USA) showing a 22q11.21 microdeletion [arr cgh 22q11.21q11.21 (RP11-690P21 → RP11-1116M14) × 1].



*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail: cpc_mmh@yahoo.com
Accepted: July 17, 2009




Figure 2. Oligonucleotide array comparative genomic hybridization using HumanCytoSNP-12 v1 BeadChips (Illumina, San Diego, CA, USA) showing a 2.6-Mb deletion in 22q11.21 [arr 22q11.21 (17141308-19792353) × 1].




Figure 3. Fluorescence *in situ* hybridization using Vysis LSI DiGeorge/velocardiofacial syndrome region/ARSA dual color DNA probes with a 22q13.3 telomere probe (LSI ARSA; green signal) and a DiGeorge critical region probe at 22q11.2 (LSI DiGeorge/velocardiofacial syndrome; red signal) (all probes by Abbott Laboratories, Abbott Park, IL, USA) showing a normal chromosome 22 (one red signal and one green signal) and a del(22)(q11.2q11.2) chromosome (green signal only) in a metaphase amniocyte.

(22)(q11.2q11.2)(D22S553-) karyotype. Microarray-based chromosome analysis showed arr 22q11.21 (17141308-19792353) × 1.

The 22q11.2 deletion has an estimated incidence of 1 per 4,000 live births [1] and has been detected in the majority of patients with DGS, VCFS and conotruncal anomaly face syndrome, as well as in some patients with Opitz G/BBB syndrome and Cayler cardiofacial syndrome [2–5], all of which are referred to as the 22q11.2 deletion syndrome [5]. The 22q11.2 deletion syndrome is characterized by congenital heart defects, aplasia or hypoplasia of the thymus with immune deficiency, small or absent parathyroid glands with hypocalcemia, palatal abnormalities with speech disorders, and cognitive difficulties [5]. Indications for prenatal testing for the 22q11.2 deletion syndrome include a previous child with DGS/VCFS or a 22q11.2 deletion, an affected parent with DGS/VCFS or a 22q11.2 deletion, and a sonographic finding of conotruncal cardiac defects [5]. Reported prenatal sonographic features associated with a 22q11.2 microdeletion

include conotruncal cardiac malformations, increased nuchal translucency (NT), polyhydramnios, intrauterine growth restriction, extracardiac anomalies, severe pulmonary artery anomalies, and thymic hypoplasia [6–12]. The present case did not present with increased NT, polyhydramnios, thymic hypoplasia or intrauterine growth restriction, but did show cardiac and pulmonary artery anomalies on prenatal ultrasound.

Lautrup et al [13] found that none of 146 fetuses with NT above the 99th centile (>3.5 mm) and a normal karyotype was diagnosed with a 22q11 microdeletion by either multiplex ligation-dependent probe amplification or FISH, and they estimated that the prevalence of a 22q11 microdeletion in these otherwise normal fetuses with increased NT was below 2.7%. In a study of 540 fetuses with prenatally detected cardiac defects and a normal karyotype, Moore et al [14] found that 17 (3.15%) had a 22q11.2 microdeletion diagnosed by FISH. In a study of 27 fetuses with intrauterine growth restriction, congenital heart defects and a normal G-banded karyotype, Chen et al [15] found that three (11.1%) had a 22q11.2 microdeletion and two (7.4%) had a subtelomeric deletion, including monosomy 21q22.3 and monosomy 1p36.3. In a review of 52 patients with DGS, Minier et al [16] found that only six (11.5%) were diagnosed during the prenatal period, 12 (23.1%) were diagnosed by pathologic examination and 34 (65.4%) were diagnosed during infancy. There is marked phenotypic variability among patients with a 22q11.2 deletion [17]. In a study of 30 individuals (19 adults and 11 children) with a 22q11.2 deletion following the diagnosis in their relatives, McDonald-McGinn et al [17] found that 60% (18/30) had no visceral anomalies, and only 32% (6/19) of the adults and 55% (6/11) of the children with a 22q11.2 deletion had major findings.

The 22q11 region is susceptible to chromosomal rearrangements leading to DGS/VCFS, cat-eye syndrome and t(11;22)der(22) syndrome, of which all three breakpoint regions harbor similar low-copy repeat sequences, also known as LCR22s [18,19]. Homologous recombination events between LCR22s during meiosis have been implicated in DGS/VCFS and cat-eye syndrome, and the sites of chromosome breakage on 11q23 and 22q11 in der(22) syndrome occur in the unstable AT-rich palindromic sequences leading to non-homologous recombination mechanisms [19–22]. About 90% of the patients with a 22q11.2 microdeletion have a common approximately 3-Mb deletion, and 7% of the patients have a smaller, nested, approximately 1.5-Mb recurrent deletion, all located in the 22q11.2 proximal deletion region [23,24]. Recently, Ben-Shachar et al [24] identified atypical deletions

located in the 22q11.2 distal region. Patients with an atypical deletion have a 22q11.2 microdeletion distal to the common deletion region and manifest a phenotype distinct from DGS/VCFS [24].

DGS (OMIM 188400), a developmental field defect of the third and fourth pharyngeal pouches, is characterized by hypocalcemia arising from parathyroid hypoplasia, thymic hypoplasia, and outflow tract defects of the heart including tetralogy of Fallot, truncus arteriosus, and interrupted aortic arch. Most cases with DGS result from a deletion of chromosome 22q11.2. However, about 10–15% of patients do not have the standard 22q11.2 deletion but have other genetic causes, such as distal deletions of chromosome 10p and chromosome 4q, unbalanced chromosomal translocations, atypical deletions of 22q11.2, and *UFD1L* haploinsufficiency [25–32]. Although FISH, ultrarapid FISH and multiplex ligation-dependent probe amplification are highly accurate tests for detecting the standard 22q11.2 microdeletion [33–35], routine aCGH has the advantage of detecting uncharacterized chromosomal imbalances and atypical 22q11.2 microdeletions, as well as refining the 22q11.2 deletion breakpoints [36,37] in the event of *in utero* detection of a fetus with congenital heart defects.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-98004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- Devriendt K, Fryns JP, Mortier G, Van Thienen MN, Keymolen K. The annual incidence of DiGeorge/velocardiofacial syndrome. *J Med Genet* 1998;35:789–90.
- Burn J, Takao A, Wilson D. Conotruncal anomaly face syndrome is associated with a deletion within chromosome 22q11. *J Med Genet* 1993;30:822–4.
- McDonald-McGinn DM, Driscoll DA, Bason L, et al. Autosomal dominant “Opitz” GBBB syndrome due to a 22q11.2 deletion. *Am J Med Genet* 1995;59:103–13.
- Bawle EV, Conard J, Van Dyke DL, Czarnecki P, Driscoll DA. Seven new cases of Cayler cardiofacial syndrome with chromosome 22q11.2 deletion, including a familial case. *Am J Med Genet* 1998;79:406–10.
- Driscoll DA. Prenatal diagnosis of the 22q11.2 deletion syndrome. *Genet Med* 2001;3:14–8.
- Chaoui R, Körner H, Bommer C. Prenatal assessment of the thymus in detecting fetuses with del.22q. *Ultrasound Obstet Gynecol* 2000;16(s1):82. [Abstract]

7. Chaoui R, Körner H, Bommer C, Kalache KD. Fetal thymus and the 22q11.2 deletion. *Prenat Diagn* 2002;22:839–40.
8. Chaoui R, Kalache KD, Heling KS, Tennstedt C, Bommer C, Körner H. Absent or hypoplastic thymus on ultrasound: a marker for deletion 22q11.2 in fetal cardiac defects. *Ultrasound Obstet Gynecol* 2002;20:546–52.
9. Boudjemline Y, Fermont L, Le Bidois J, Villain E, Sidi D, Bonnet D. Can we predict 22q11 status of fetuses with tetralogy of Fallot? *Prenat Diagn* 2002;22:231–4.
10. Barrea C, Yoo SJ, Chitayat D, Valsangiacomo E, Winsor E, Smallhorn JF, Hornberger LK. Assessment of the thymus at echocardiography in fetuses at risk for 22q11.2 deletion. *Prenat Diagn* 2003;23:9–15.
11. Volpe P, Marasini M, Caruso G, et al. 22q11 deletions in fetuses with malformations of the outflow tracts or interruption of the aortic arch: impact of additional ultrasound signs. *Prenat Diagn* 2003;23:752–7.
12. Chen CP, Chien SC. Prenatal sonographic features of 22q11.2 microdeletion syndrome. *J Med Ultrasound* 2008;16:123–9.
13. Lautrup CK, Kjaergaard S, Brøndum-Nielsen K, et al. Testing for 22q11 microdeletion in 146 fetuses with nuchal translucency above the 99th percentile and a normal karyotype. *Acta Obstet Gynecol Scand* 2008;87:1252–5.
14. Moore JW, Binder GA, Berry R. Prenatal diagnosis of aneuploidy and deletion 22q11.2 in fetuses with ultrasound detection of cardiac defects. *Am J Obstet Gynecol* 2004;191:2068–73.
15. Chen M, Hwu WL, Kuo SJ, et al. Subtelomeric rearrangements and 22q11.2 deletion syndrome in anomalous growth-restricted fetuses with normal or balanced G-banded karyotype. *Ultrasound Obstet Gynecol* 2006;28:939–43.
16. Minier F, Carles D, Pelluard F, Alberti EM, Stern L, Saura R. DiGeorge syndrome, a review of 52 patients. *Arch Pediatr* 2005;12:254–7. [In French]
17. McDonald-McGinn DM, Tonnesen MK, Laufer-Cahana A, Finucane B, Driscoll DA, Emanuel BS, Zackai EH. Phenotype of the 22q11.2 deletion in individuals identified through an affected relative: cast a wide FISHing net! *Genet Med* 2001;3:23–9.
18. Edelmann L, Pandita RK, Morrow BE. Low-copy repeats mediate the common 3-Mb deletion in patients with velocardio-facial syndrome. *Am J Hum Genet* 1999;64:1076–86.
19. McDermid HE, Morrow BE. Genomic disorders on 22q11. *Am J Hum Genet* 2002;70:1077–88.
20. Edelmann L, Pandita RK, Spiteri E, et al. A common molecular basis for rearrangement disorders on chromosome 22q11. *Hum Mol Genet* 1999;8:1157–67.
21. Edelmann L, Spiteri E, McCain N, et al. A common breakpoint on 11q23 in carriers of the constitutional t(11;22) translocation. *Am J Hum Genet* 1999;65:1608–16.
22. Edelmann L, Spiteri E, Koren K, et al. AT-rich palindromes mediate the constitutional t(11;22) translocation. *Am J Hum Genet* 2001;68:1–13.
23. Carlson C, Sirotkin H, Pandita R, et al. Molecular definition of 22q11 deletions in 151 velo-cardio-facial syndrome patients. *Am J Hum Genet* 1997;61:620–9.
24. Ben-Shachar S, Ou Z, Shaw CA, et al. 22q11.2 distal deletion: a recurrent genomic disorder distinct from DiGeorge syndrome and velocardiofacial syndrome. *Am J Hum Genet* 2008;82:214–21.
25. Daw SCM, Taylor C, Kraman M, et al. A chromosome region of 10p deleted in DiGeorge and velocardiofacial syndromes. *Nat Genet* 1996;13:458–60.
26. Kurahashi H, Nakayama T, Osugi Y, et al. Deletion mapping of 22q11 in CATCH22 syndrome: identification of a second critical region. *Am J Hum Genet* 1996;58:1377–81.
27. Gottlieb S, Driscoll DA, Punnett H, Sellinger B, Emanuel BS, Budarf ML. Characterization of 10p deletions suggests two nonoverlapping regions contribute to the DiGeorge syndrome phenotype. *Am J Hum Genet* 1998;62:495–8.
28. McQuade L, Christodoulou J, Budarf M, Sachdev R, Wilson M, Emanuel B, Colley A. Patient with a 22q11.2 deletion with no overlap of the minimal DiGeorge critical region (MDGCR). *Am J Med Genet* 1999;86:27–33.
29. Rausch A, Pfeiffer RA, Leipold G, Singer H, Tigges M, Hofbeck M. A novel 22q11.2 microdeletion in DiGeorge syndrome. *Am J Hum Genet* 1999;64:659.
30. Saitta S, McGrath JM, Mensch H, Shaikh TH, Zackai EH, Emanuel BS. A 22q11.2 deletion that excludes *UFD1L* and *CDC45L* in a patient with conotruncal and craniofacial defects. *Am J Hum Genet* 1999;65:562–6.
31. Tsai CH, Van Dyke DL, Feldman GL. Child with velocardio-facial syndrome and del(4)(q34.2): another critical region associated with a velocardiofacial syndrome-like phenotype. *Am J Med Genet* 1999;82:336–9.
32. Yamagishi H, Garg V, Matsuoka R, Thomas T, Srivastava D. A molecular pathway revealing a genetic basis for human cardiac and craniofacial defects. *Science* 1999;283:1158–61.
33. Choolani M, Ho SS, Razvi K, et al. FastFISH: technique for ultrarapid fluorescence in situ hybridization on uncultured amniocytes yielding results within 2 h of amniocentesis. *Mol Hum Reprod* 2007;13:355–9.
34. Göktolga U, Gezginç K, Ceyhan ST, Fidan U, Ergün A, Bahçe M, Başer I. Prenatal diagnosis of DiGeorge syndrome. *Taiwan J Obstet Gynecol* 2008;47:441–2.
35. Jalali GR, Vorstman JA, Errami A, Vijzelaar R, Biegel J, Shaikh T, Emanuel BS. Detailed analysis of 22q11.2 with a high density MLPA probe set. *Hum Mutat* 2008;29: 433–40.
36. Bittel DC, Yu S, Newkirk H, Kibiryeva N, Holt A 3rd, Butler MG, Cooley LD. Refining the 22q11.2 deletion breakpoints in DiGeorge syndrome by aCGH. *Cytogenet Genome Res* 2009;124:113–20.
37. Vialard F, Molina Gomes D, Leroy B, et al. Array comparative genomic hybridization in prenatal diagnosis: another experience. *Fetal Diagn Ther* 2009;25:277–84.