

RAPID GENOME-WIDE ANEUPLOIDY DIAGNOSIS USING UNCULTURED AMNIOCYTES AND ARRAY COMPARATIVE GENOMIC HYBRIDIZATION IN PREGNANCY WITH ABNORMAL ULTRASOUND FINDINGS DETECTED IN LATE SECOND AND THIRD TRIMESTERS

Chih-Ping Chen^{1,2,3,4*}, Yi-Ning Su⁵, Fuu-Jen Tsai^{4,6,7}, Schu-Rern Chern², Chin-Yuan Hsu¹, Ming-Chao Huang¹, Wayseen Wang^{2,8}

Departments of ¹Obstetrics and Gynecology and ²Medical Research, Mackay Memorial Hospital, Taipei,

³Department of Biotechnology, Asia University, ⁴School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, ⁵Department of Medical Genetics, National Taiwan University Hospital, Taipei, Departments of

⁶Medical Genetics, and ⁷Medical Research, China Medical University Hospital, Taichung, and

⁸Department of Bioengineering, Tatung University, Taipei, Taiwan.

A 32-year-old, gravida 2, para 1, woman was referred to the hospital at 23 weeks of gestation because of fetal cardiac abnormalities detected by sonography. The woman and her husband were non-consanguineous and healthy, and there was no family history of congenital heart defects. The woman had a cousin who suffered from mental retardation and Down syndrome. Level II ultrasound revealed a singleton fetus with a fetal biometry equivalent to 23 weeks of gestation and a ventricular septal defect. Amniocentesis was performed, and 36 mL of amniotic fluid was aspirated. About 20 mL of amniotic fluid was used for array comparative genomic hybridization (aCGH) using uncultured amniocytes, and 15 mL was used for conventional cytogenetic analysis using cultured amniocytes. Within 3 days, bacterial artificial chromosome (BAC)-based aCGH showed trisomy 21 [arr cgh 21p11.2q22.3 (RP11-430M17 → RP11-1000I21)×3] (Figure 1A). Fluorescence *in situ* hybridization (FISH) analysis of the cultured interphase amniocytes using a combination of BAC probes RP11-135B17 (21q22.3) (47,931,911-48,108,188) (spectrum green) and RP11-450H16 (13q34) (114,940,140-115,099,262) (spectrum red) showed three green




Figure 1. Bacterial artificial chromosome (BAC)-based array comparative genomic hybridization analyses using CMDX (CMDX, Irvine, CA, USA) BAC array comparative genomic hybridization CA3000 chips. (A) Case 1 and (B) Case 2 showed duplications of chromosome 21 consistent with the diagnosis of trisomy 21.



ELSEVIER

*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail: cpc_mmh@yahoo.com

Accepted: September 29, 2009

signals and two red signals, consistent with the diagnosis of trisomy 21 (Figure 2A). Eight days after amniocentesis, conventional cytogenetic analysis revealed a 47,XX,+21 karyotype. Polymorphic DNA marker analysis using quantitative fluorescence polymerase chain reaction (QF-PCR) showed that the trisomy 21 in this fetus was likely to have been caused by paternal meiosis II nondisjunction or postzygotic mitotic nondisjunction (Figure 3).

A 33-year-old, gravida 2, para 1, woman was referred to the hospital at 30 weeks of gestation because




Figure 2. Fluorescence *in situ* hybridization analyses of interphase amniocytes using bacterial artificial chromosome probes RP11-135B17 (21q22.3) (spectrum green) and RP11-450H16 (13q34) (spectrum red). (A) Case 1 and (B) Case 2 showed three green signals and two red signals, consistent with the diagnosis of trisomy 21.




Figure 3. Representative electrophoretogram of quantitative fluorescence polymerase chain reaction assays in Case 1 at short tandem repeat markers for chromosome 21q. Two peaks (250 bp and 266 bp) of unequal fluorescence activity from different parental alleles (maternal:paternal) at a ratio of 1:2 were detected in the cultured fetal tissues using the marker D21S1443. This indicated a homologous duplication of chromosome 21q of paternal origin and was likely to be the result of meiosis II nondisjunction or postzygotic mitotic nondisjunction.

of fetal brain abnormalities detected by sonography. The woman and her husband were non-consanguineous and healthy, and there was no family history of congenital anomalies. Level II ultrasound revealed a singleton fetus with a fetal biometry equivalent to 31 weeks of gestation and bilateral ventriculomegaly with a right ventricular width of 1.7 cm and a left ventricular width of 1.3 cm. Amniocentesis was performed, and 40 mL of amniotic fluid was aspirated. About 20 mL of amniotic fluid was used for aCGH using uncultured amniocytes, and 20 mL was used for conventional cytogenetic analysis using cultured amniocytes. Within 3 days, BAC-based aCGH showed trisomy 21 [arr cgh 21p11.2q22.3 (RP11-1144F24 → RP11-1000I21) × 3] (Figure 1B). FISH (Figure 2B) and QF-PCR (Figure 4) analyses using cultured amniocytes revealed results consistent with trisomy 21. Thirteen days after amniocentesis, conventional cytogenetic analysis revealed a 47,XX,+21 karyotype. The parental DNA was not available. According to polymorphic DNA marker analysis, trisomy 21 in this fetus was likely to have been caused by meiosis I nondisjunction.

We have previously described the prenatal diagnosis of aneuploidy by aCGH using cultured or uncultured amniocytes [1,2]. The current cases further demonstrated that amniocentesis for genome-wide analysis using uncultured amniocytes and aCGH is a useful alternative to cordocentesis for rapid aneuploidy diagnosis in pregnancies with abnormal ultrasound findings detected during the late second and third trimesters.

Amniocentesis is simpler and less invasive than cordocentesis. The fetal loss rate is 0.5% for amniocentesis [3,4], compared with 0.8–2.5% for cordocentesis [5–8]. Cordocentesis is an invasive procedure associated with risks such as fetal loss, fetal bleeding and hemorrhage, fetal bradycardia, fetomaternal transfusion, umbilical cord thrombosis and hemorrhage,




Figure 4. Representative electrophoretogram of quantitative fluorescence polymerase chain reaction assays of Case 2 at short tandem repeat markers for chromosome 21q. Three peaks (116 bp, 124 bp and 132 bp) of equal fluorescence activity at a ratio of 1:1:1 were detected in the cultured amniocytes. This indicated a heterologous duplication of chromosome 21q and was likely to be the result of meiosis I nondisjunction. The parental DNA was not available.

infection, abruptio placentae, and premature labor, even when performed in expert centers [9]. However, cordocentesis is preferable to amniocentesis for conventional cytogenetic analysis, especially in the late second and third trimesters, because it requires less time for cell culture. Fetal karyotypes can be obtained by culture of fetal cord blood lymphocytes from cordocentesis in 2–4 days. However, rapid karyotyping by cordocentesis has been replaced by amniocentesis combined with FISH or QF-PCR [10–15]. FISH and QF-PCR results can be obtained in 24–48 hours, but can detect only specific aneuploidies according to the specific probes or primers used. Recently, aCGH using cultured or uncultured amniocytes has been successfully applied for the prenatal diagnosis of chromosome abnormalities [16–27]. aCGH can detect unbalanced structural and numerical chromosome abnormalities of less than 100 kb, whereas conventional cytogenetic analysis can detect only microscopically visible deletions or duplications of 5–6 Mb at the 500-band level [25]. Tyreman et al [26] found that high-resolution array testing would benefit at least 10% of obstetric patients with abnormal ultrasound findings and a normal karyotype result. aCGH has the advantage of providing a rapid genome-wide study without the need for cell culture. However, aCGH also has the disadvantage of being unable to detect balanced translocations, inversions or polyploidy. Detection of low-level mosaicism depends on the dynamic range of the array used, and several studies have shown that aCGH can detect low-level mosaicism not revealed by conventional cytogenetics [25,28–30]. We, therefore, suggest that in addition to rapid karyotyping by aCGH, conventional cytogenetic analysis is still required to rule out the presence of balanced translocations, inversions and polyploidy.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-98004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- Chen CP, Su YN, Chang TY, Chern SR, Tsai FJ, Hwang JK, Wang W. 22q11.2 microdeletion in a fetus with double-outlet right ventricle, pulmonary stenosis and a ventricular septal defect: prenatal diagnosis by array comparative genomic hybridization. *Taiwan J Obstet Gynecol* 2009;48:437–40.
- Chen CP, Su YN, Tsai FJ, et al. Terminal 2q deletion and distal 15q duplication: prenatal diagnosis by array comparative genomic hybridization using uncultured amniocytes. *Taiwan J Obstet Gynecol* 2009;48:441–5.
- CDC publishes report on chorionic villus sampling and amniocentesis. *Am Fam Physician* 1995;52:1210.
- Centers for Disease Control and Prevention. Chorionic villus sampling and amniocentesis: recommendations for prenatal counseling. *MMWR Recomm Rep* 1995;44:1–12.
- Daffos F, Capella-Pavlovsky M, Forestier F. Fetal blood sampling during pregnancy with use of a needle guided by ultrasound: a study of 606 consecutive cases. *Am J Obstet Gynecol* 1985;153:655–60.
- Nicolaidis P, Nicolini U, Fisk NM, Tannirandorn Y, Nasrat H, Rodeck CH. Fetal blood sampling from the intrahepatic vein for rapid karyotyping in the second and third trimesters. *Br J Radiol* 1991;64:505–9.
- Weiner CP, Okamura K. Diagnostic fetal blood sampling-technique related losses. *Fetal Diagn Ther* 1996;11:169–75.
- Antsaklis A, Daskalakis G, Papantoniou N, Michalas S. Fetal blood sampling—indication-related losses. *Prenat Diagn* 1998;18:934–40.
- Monni G. Prenatal diagnosis by cordocentesis. In: Weiner S, Kurjak A, eds. *Interventional Ultrasound: Progress in Obstetric and Gynecological Sonography Series*. New York: Parthenon Publishing Group, 1999:31–40.
- Verma L, Macdonald F, Leedham P, McConachie M, Dhanjal S, Hulten M. Rapid and simple prenatal DNA diagnosis of Down's syndrome. *Lancet* 1998;352:9–12.
- Levett LJ, Liddle S, Meredith R. A large-scale evaluation of amnio-PCR for the rapid prenatal diagnosis of fetal trisomy. *Ultrasound Obstet Gynecol* 2001;17:115–8.
- Mann K, Fox SP, Abbs SJ, Yau SC, Scriven PN, Docherty Z, Ogilvie CM. Development and implementation of a new rapid aneuploidy diagnostic service within the UK National Health Service and implications for the future of prenatal diagnosis. *Lancet* 2001;358:1057–61.
- Bili C, Divane A, Apeossos A, et al. Prenatal diagnosis of common aneuploidies using quantitative fluorescent PCR. *Prenat Diagn* 2002;22:360–5.
- Cirigliano V, Voglino G, Cañadas MP, et al. Rapid prenatal diagnosis of common chromosome aneuploidies by QF-PCR: assessment on 18,000 consecutive clinical samples. *Mol Hum Reprod* 2004;10:839–46.
- Cirigliano V, Voglino G, Marongiu A, et al. Rapid prenatal diagnosis by QF-PCR: evaluation of 30,000 consecutive clinical samples and future applications. *Ann N Y Acad Sci* 2006;1075:288–98.
- Rickman L, Fiegler H, Carter NP, Bobrow M. Prenatal diagnosis by array-CGH. *Eur J Med Genet* 2005;48:232–40.
- Rickman L, Fiegler H, Shaw-Smith C, et al. Prenatal detection of unbalanced chromosomal rearrangements by array CGH. *J Med Genet* 2006;43:353–61.
- Sahoo T, Cheung SW, Ward P, et al. Prenatal diagnosis of chromosomal abnormalities using array-based comparative genomic hybridization. *Genet Med* 2006;8:719–27.
- van den Veyver IB, Beaudet AL. Comparative genomic hybridization and prenatal diagnosis. *Curr Opin Obstet Gynecol* 2006;18:185–91.

20. Shaffer LG, Bui TH. Molecular cytogenetic and rapid aneuploidy detection methods in prenatal diagnosis. *Am J Med Genet C Semin Med Genet* 2007;145C:87–98.
21. Bi W, Breman AM, Venable SF, et al. Rapid prenatal diagnosis using uncultured amniocytes and oligonucleotide array CGH. *Prenat Diagn* 2008;28:943–9.
22. Cain CC, Saul DO, Oehler E, Blakemore K, Stetten G. Prenatal detection of a subtle unbalanced chromosome rearrangement by karyotyping, FISH and array comparative genomic hybridization. *Fetal Diagn Ther* 2008;24:286–90.
23. Darilek S, Ward P, Pursley A, et al. Pre- and postnatal genetic testing by array-comparative genomic hybridization: genetic counseling perspectives. *Genet Med* 2008;10:13–8.
24. Kashork CD, Theisen A, Shaffer LG. Prenatal diagnosis using array CGH. *Methods Mol Biol* 2008;444:59–69.
25. Breman AM, Bi WM, Cheung SW. Prenatal diagnosis by array-based comparative genomic hybridization in the clinical laboratory setting. *Beijing Da Xue Xue Bao* 2009;41:500–4.
26. Tyreman M, Abbott KM, Willatt LR, Nash R, Lees C, Whittaker J, Simoncic I. High resolution array analysis: diagnosing pregnancies with abnormal ultrasound findings. *J Med Genet* 2009;46:531–41.
27. Vialard F, Molina Gomes D, Leroy B, et al. Array comparative genomic hybridization in prenatal diagnosis: another experience. *Fetal Diagn Ther* 2009;25:277–84.
28. Ballif BC, Rorem EA, Sundin K, et al. Detection of low-level mosaicism by array CGH in routine diagnostic specimens. *Am J Med Genet A* 2006;140:2757–67.
29. Cheung SW, Shaw CA, Scott DA, et al. Microarray-based CGH detects chromosomal mosaicism not revealed by conventional cytogenetics. *Am J Med Genet A* 2007;143:1679–86.
30. Shinawi M, Shao L, Jeng LJ, et al. Low-level mosaicism of trisomy 14: phenotypic and molecular characterization. *Am J Med Genet A* 2008;146:1395–405.