

CHROMOSOME 1P36 DELETION SYNDROME: PRENATAL DIAGNOSIS, MOLECULAR CYTOGENETIC CHARACTERIZATION AND FETAL ULTRASOUND FINDINGS

Chih-Ping Chen^{1,2,3,4,5,6*}, Ming Chen^{7,8,9,10}, Yi-Ning Su¹¹, Chin-Yuan Hsu¹, Fuu-Jen Tsai^{4,12,13},
Schu-Rern Chern², Pei-Chen Wu¹, Chen-Chi Lee¹, Wayseen Wang^{2,14}

Departments of ¹Obstetrics and Gynecology and ²Medical Research, Mackay Memorial Hospital, ⁵Institute of Clinical and Community Health Nursing, ⁶Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, ¹⁰Department of Obstetrics and Gynecology College of Medicine, National Taiwan University, ¹¹Department of Medical Genetics, National Taiwan University Hospital, and ¹⁴Department of Bioengineering, Tatung University, Taipei; ³Department of Biotechnology, Asia University, ⁴School of Chinese Medicine, College of Chinese Medicine, China Medical University, and Departments of ¹²Medical Research and ¹³Medical Genetics, China Medical University Hospital, Taichung; Departments of ⁷Genomic Medicine, ⁸Medical Research and ⁹Obstetrics and Gynecology, Changhua Christian Hospital, Changhua, Taiwan.

SUMMARY

Objective: To present prenatal diagnosis and molecular cytogenetic characterization of *de novo* partial monosomy 1p (1p36.23→pter) and partial trisomy 20p (20p12.1→pter) associated with ventriculomegaly, ventricular septal defect and midface hypoplasia.

Materials, Methods and Results: A 31-year-old, primigravid woman was referred for amniocentesis at 20 gestational weeks because of ventriculomegaly, ventricular septal defect, and midface hypoplasia. Amniocentesis revealed an aberrant derivative chromosome 1, or der(1). Parental karyotypes were normal. Spectral karyotyping analysis revealed that the der(1) contained a segment of chromosome 20 in the distal end of the short arm of chromosome 1. Array comparative genomic hybridization demonstrated an 8.4-Mb distal 1p deletion and a 14-Mb distal 20p duplication. The karyotype was 46,XX,der(1)t(1;20)(p36.23;p12.1)dn. Polymorphic DNA marker analysis determined the paternal origin of the aberrant chromosome. The pregnancy was subsequently terminated. A 462-g malformed female fetus was delivered at 22 gestational weeks with a prominent forehead, midface hypoplasia, a flat nasal bridge, low-set ears, a long philtrum, a pointed chin and micrognathia.

Conclusion: Spectral karyotyping, fluorescence *in situ* hybridization and array comparative genomic hybridization are useful for the prenatal investigation of the nature of a *de novo* aberrant derivative chromosome. Partial monosomy 1p (1p36.23→pter) and partial trisomy 20p (20p12.1→pter) are associated with ventriculomegaly, ventricular septal defect and midface hypoplasia on prenatal ultrasound. Prenatal diagnosis of ventriculomegaly, congenital heart defects and midface hypoplasia should alert clinicians to chromosome 1p36 deletion syndrome and prompt molecular cytogenetic analysis if necessary. [*Taiwan J Obstet Gynecol* 2010;49(4):473–480]

Key Words: chromosome 1, chromosome 1p36 deletion syndrome, chromosome 20, monosomy 1p36, prenatal diagnosis, ultrasound



*Correspondence to: Dr Chih-Ping Chen, Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail: cpc_mmh@yahoo.com
Accepted: September 1, 2010

Introduction

Prenatal diagnosis of a *de novo* unbalanced reciprocal translocation involving chromosomal segments with a subtle difference in banding results in difficulties in

interpretation and genetic counseling, and requires molecular cytogenetic technologies such as spectral karyotyping (SKY), fluorescence *in situ* hybridization (FISH), and array-based comparative genomic hybridization (aCGH) to identify the nature of the aberrant chromosome. We previously reported the utility of SKY and FISH in the identification of a *de novo* aberrant chromosome derived from an unbalanced reciprocal translocation [1]. We additionally report here the pre-natal diagnosis and molecular cytogenetic characterization of *de novo* partial monosomy 1p (1p36.23→pter) and partial trisomy 20p (20p12.1→pter) associated with

ventriculomegaly, a ventricular septal defect (VSD) and midface hypoplasia.

Materials, Methods and Results

A 31-year-old, primigravid woman was referred to the genetic counseling center for amniocentesis at 20 gestational weeks because of abnormal ultrasound findings of ventriculomegaly, VSD and midface hypoplasia (Figure 1). Amniocentesis revealed an aberrant derivative chromosome 1, or der(1) (Figure 2). Chromosome




Figure 1. Prenatal ultrasound at 20 gestational weeks reveals (A) ventriculomegaly, (B) a ventricular septal defect and (C) midface hypoplasia.




Figure 2. G-banded karyotype shows a derivative chromosome 1, or der(1). The proband's karyotype is 46,XX,der(1)t(1;20)(p36.23;p12.1)dn. The arrows indicate the breakpoints.

preparations of blood lymphocytes from the parents revealed normal karyotypes. The derivative chromosome was characterized by SKY using 24-color SKY probes (Applied Spectral Imaging, Carlsbad, CA, USA). SKY analysis revealed that the der(1) contained a segment of chromosome 20 in the distal end of the short arm of chromosome 1 (Figure 3). Repeat amniocentesis was performed at 22 gestational weeks, and 20 mL of aspirated amniotic fluid was applied for aCGH using

uncultured amniocytes. Bacterial artificial chromosome (BAC)-based aCGH using CMDX BAC-based aCGH CA2500 chips (CMDX, Irvine, CA, USA) demonstrated partial monosomy 1p and partial trisomy 20p [arr 1p36.33p36.23 (RP11-668E2→RP11-81J7)×1, 20p13p12.1 (RP11-530N10→RP11-238I2)×3] (Figure 4). Oligonucleotide-based aCGH using HumanCytoSNP-12v1 BeadChips (Illumina, San Diego, CA, USA) further demonstrated an 8.4-Mb distal 1p deletion and a 14-Mb




Figure 3. Spectral karyotyping using 24-color SKY probes shows a der(1) derived from a translocation between chromosomes 1 and 20.




Figure 4. Bacterial artificial chromosome-based array comparative genomic hybridization shows a distal 1p deletion [arr 1p36.33p36.23 (RP11-668E2→RP11-81J7)×1] and a distal 20p duplication [arr 20p13p12.1 (RP11-530N10→RP11-238I2)×3].




Figure 5. Oligonucleotide-based array comparative genomic hybridization shows (A) an 8.4-Mb distal 1p deletion [arr 1p36.33p36.23 (742,429-9,132,849)×1] and (B) a 14-Mb distal 20p duplication [arr 20p13p12.1 (467,697-14,465,220)×3].

distal 20p duplication [arr 1p36.33p36.23 (742,429-9,132,849)×1, 20p13p12.1 (467,697-14,465,220)×3] (Figure 5). The karyotype was 46,XX,der(1)t(1;20)(p36.23;p12.1)dn (Figure 2). Polymorphic DNA marker analysis determined the paternal origin of the aberrant chromosome. The parents opted to terminate the pregnancy. A 462-g malformed female fetus was delivered with a prominent forehead, midface hypoplasia, a flat nasal bridge, low-set ears, a long philtrum, a pointed chin and micrognathia (Figure 6).

Discussion

Chromosome 1p36 deletion syndrome (OMIM 607872), or monosomy 1p36 syndrome, is the most common subtelomeric terminal deletion syndrome, and it is associated with multiple congenital anomalies and mental retardation [2]. Chromosome 1p36 deletion syndrome has an estimated frequency of 1 in 5,000 live births [3] and is characterized by a typical craniofacial dysmorphism of straight eyebrows, deep-set eyes, midface




Figure 6. Anterior and lateral views of the craniofacial appearance of the proband.

hypoplasia, a broad nasal bridge, a long philtrum, a pointed chin, large anterior fontanel, microbrachycephaly, epicanthal folds, low-set ears, brachydactyly, camptodactyly, short feet, developmental delay, mental retardation, hypotonia, seizures, structural brain abnormalities, congenital heart defects, eye problems, hearing loss and skeletal, renal and genital abnormalities [4,5].

The present case prenatally manifested ventriculomegaly, VSD, and midface hypoplasia during second-trimester ultrasound. In a study of 60 patients with chromosome 1p36 deletion syndrome, Battaglia et al [5] found midface hypoplasia in 100% (60/60), brain abnormalities in 88% (43/49) and congenital heart defects in 71% (34/48) of patients. In their study, the reported congenital heart defects included atrial septal defect (28%), VSD (23%), patent ductus arteriosus (12%), valvular anomalies (20.5%), tetralogy of Fallot (7.7%), coarctation of the aorta (5.1%), infundibular stenosis of the right ventricle (2%) and Ebstein's anomaly (2%), and brain magnetic resonance imaging abnormalities included enlargement of the lateral ventricles (37%), cortical atrophy (20%), enlargement of the subarachnoid spaces (22%), diffuse brain atrophy (10%), and enlargement of the frontotemporal opercula (4%).

In chromosomal abnormalities observed in chromosome 1p36 deletion syndrome, 52% of the rearrangements are terminal deletions, 29% are interstitial deletions, 12% are complex rearrangements including more than one deletion or deletions with duplications, triplications, insertions and/or inversions, and only 7% are derivative chromosomes derived from unbalanced translocations [4,6]. The present case had a karyotype of 46,XX,der(1)t(1;20)(p36.23;p12.1)dn. To our knowledge, this case is the fourth reported case of monosomy 1p36 associated with an unbalanced translocation between chromosome 1p and 20p. Howard-Peebles and Black [7] first reported the diagnosis of monosomy

1p36 and a karyotype of 46,XY,der(1)t(1;20)(p36;p13) in a fetus conceived by a carrier mother who had a previous aneuploid child with 46,XX,der(1)t(1;20)(p36;p13) and multiple anomalies. The male fetus prenatally manifested hydrocephalus and complex congenital heart defects during prenatal ultrasound at 16 gestational weeks, and postnatally manifested truncus arteriosus, VSD, agenesis of the corpus callosum and dilated ventricles. The female aneuploid child had microcephaly, microphthalmia, optic nerve colobomas, facial dysmorphism, agenesis of the corpus callosum, hydrocephalus, tetralogy of Fallot and neonatal death. Campeau et al [8] additionally reported prenatal diagnosis of monosomy 1p36 and a karyotype of 46,XX,der(1)t(1;20)(p36.1;p12.2) by amniocentesis in a fetus conceived by a 30-year-old carrier mother because of severe fetal ventriculomegaly on prenatal ultrasound at 21 gestational weeks. Parental cytogenetic analysis showed that the mother's karyotype was 46,XX,t(1;20)(p36.1;p12.2). The phenotype of partial trisomy 20p includes a low birth weight, occipital flattening, a round face, prominent cheeks, upslanting palpebral fissures, hypertelorism, epicanthal folds, a short nose, coarse hair, congenital heart defects, vertebral anomalies and clinodactyly [9]. Although some of the clinical features observed in our patient could be due to the effect of partial trisomy 20p in addition to monosomy 1p36, partial trisomy 20p has been shown to be associated with no structural brain malformations [9–10].

To date, at least 10 cases of prenatally detected chromosome 1p36 deletion syndrome have been reported [8,11–19]. Chromosome 1p36 deletion syndrome may prenatally manifest elevated maternal serum α -fetoprotein levels, congenital heart defects and cerebral malformations (Table). Our case additionally shows that midface hypoplasia can be a cardinal prenatal sonographic feature of chromosome 1p36 deletion syndrome. Heilstedt et al [16] reported increased maternal serum

Table. Clinical findings of cases with prenatally detected or undetected chromosome 1p36 deletion syndrome

Author (year)	Maternal age (yr)	Indication for prenatal genetic diagnosis	Chromosomal aberration	Inheritance	Outcome
Howard-Peebles & Black [7] (1994)	NA	Abnormal ultrasound: hydrocephalus and complex congenital heart defects at 16 wk	Prenatal diagnosis: 46,XY Postnatal diagnosis: 46,XY,der(1)t(1;20)(p36;p13)mat	Maternal	Termination: truncus arteriosus, VSD, ACC and dilated ventricles
Chen et al [11] (1997)	33	Elevated MSAFP, abnormal ultrasound: polyhydramnios, hydronephrosis, cerebellar and ventricular enlargement	46,XY,der(1)t(1;4)(p36.13;q33)mat	Maternal	Delivered with postnatal growth retardation, microcephaly, plagiocephaly and minor anomalies; seizures began at 13 weeks old
Kulharya et al [12] (1997)	NA	Abnormal ultrasound: absence of cerebellar vermis, thick nuchal folds, bilateral hydronephrosis, ascites, club feet, arthrogryposis and a single umbilical artery at 23.5 wk	46,XY,der(1)t(1;6)(p36.3;q22.2)	<i>De novo</i>	Delivery at 37 wk, facial dysmorphism, ASD, PDA, ambiguous genitalia, hypospadias, hydronephrosis, rudimentary postaxial polydactyly of the left hand, equinovarus deformity of the feet, neonatal death
Robbins-Furman et al [13] (1997)	NA	Elevated MSAFP	46,XX,del(1)(p36.3)	<i>De novo</i>	Oligohydramnios; delivery at 37 wk with dysmorphic features; neonatal seizures began at 9 weeks old
Faivre et al [14] (1999)	34	Abnormal ultrasound: ventriculomegaly, hypertelorism and Ebstein's anomaly at 24 wk	46,XY,del(1)(p36)	<i>De novo</i>	Termination at 28 wk, hydrops fetalis, ventriculomegaly, pleural effusion, Ebstein's anomaly
Cavani et al [15] (2003) Case III, 3	33	Abnormal ultrasound: IUGR and anomaly of the corpus callosum at 31 wk	46,XY,der(1)t(1;6)(p36.3;q25.2)mat	Maternal	Delivered at 34 wk, IUGR, macrocephaly, lobar holoprosencephaly, right optic nerve coloboma, dilated third and lateral ventricles, VSD, neonatal death
Case III, 5	NA	Previous aneuploid child The same mother as Case III, 3	46,XX,der(1)t(1;6)(p36.3;q25.2)mat	Maternal	Termination, ventriculomegaly, diffuse edema, pulmonary segmentation defects
Case III, 6	NA	Previous aneuploid child The same mother as Case III, 3	46,XY,der(1)t(1;6)(p36.3;q25.2)mat	Maternal	Termination

Reddy and Yang [17] (2003) Case 1	15	Abnormal ultrasound: hydrocephalus and IUGR at 30.3 wk	Prenatal diagnosis: 46,XX Postnatal diagnosis: 46,XX,der(1)t(1;9)(p36.3;p13)[88]/ 46,XX[12]	<i>De novo</i>	Delivered with PDA, unilateral optic nerve hypoplasia, pes calcaneovalgus, dysmorphic features; developmental delay at 6 months of age Termination at 20 wk
	29	Abnormal ultrasound: bilateral choroid plexus cysts, VSD, rocker-bottom feet and hydrocephalus at 18 wk	46,XY,der(1)t(1;21)(p36.3;q22.1)	<i>De novo</i>	
	29	Abnormal ultrasound: ventriculomegaly, a small choroid plexus cyst and nuchal edema at 21 wk	46,XY,del(1)(p36)	<i>De novo</i>	Delivery at 40 wk, dysmorphic features, hypertonia, neonatal seizures; delayed development, enlargement of the third and lateral ventricles, small periventricular cysts, delayed myelination at 5 months of age
Campeau et al [8] (2008) Case 1	31	Abnormal ultrasound: ventriculomegaly, a single umbilical artery and short femurs at 17 wk; IUGR, ventriculomegaly, club foot, VSD, asymmetric ventricles and tortuous aortic arch at 31 wk Abnormal ultrasound: severe ventriculomegaly at 21 wk	Prenatal diagnosis: 46,XX Postnatal diagnosis: 46,XX,del(1)(p36.3)	<i>De novo</i>	Delivered at 34 wk, cleft palate, ASD, VSD, PDA, dysmorphic features, hypotonia; hydrocephalus and colpocephaly on MRI at 2 years old; global developmental delay at 2.5 years old Termination at 21 ⁺ 5 wk, facial dysmorphism, clinodactyly, flexion contraction of lower limbs, misaligned toes, hydrocephalus, cerebellar hypoplasia, focal polymicrogyria
	30		46,XX,der(1)t(1;20)(p36.1;p12.2)mat	Maternal	
Vialard et al [20] (2009) Case 39	NA	Abnormal ultrasound: congenital heart defects and cerebral malformation	Prenatal diagnosis: 46,XY Postnatal diagnosis: 46,XY,del(1)(p36.22)	NA	Ultrasound at 27 wk revealed partial corpus callosum, vermian hypoplasia, aortic coarctation, two intraventricular communications, right heart dilation, tricuspid valve dysgenesis
	31	Abnormal ultrasound: ventriculomegaly, VSD and midface hypoplasia at 20 wk	46,XX,der(1)t(1;20)(p36.23;p12.1)	<i>De novo</i>	Termination at 22 wk, facial dysmorphism

NA = not available; MSAFP = maternal serum α -fetoprotein; VSD = ventricular septal defect; ACC = agenesis of corpus callosum; ASD = atrial septal defect; PDA = patent ductus arteriosus; IUGR = intrauterine growth restriction; MRI = magnetic resonance imaging.

α -fetoprotein levels in four of five prenatally diagnosed cases of monosomy 1p36 and suggested that careful attention should be paid to chromosome 1p36 in case of prenatal cytogenetic testing for increased maternal serum α -fetoprotein levels. The Table also shows that terminal deletion 1p36 can be missed at prenatal diagnosis because of a lower level of resolution [7,8,17,20]. In this regard, aCGH using cultured or uncultured amniocytes may be needed to correctly identify subtle chromosome aberrations [21,22]. We suggest that prenatal diagnosis of ventriculomegaly, congenital heart defects and midface hypoplasia should alert clinicians to the possibility of chromosome 1p36 deletion syndrome and prompt molecular cytogenetic analysis if necessary.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-99004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- Chen CP, Chen M, Su YN, et al. Prenatal diagnosis and molecular cytogenetic characterization of *de novo* partial trisomy 7p (7p15.3→pter) and partial monosomy 13q (13q33.3→qter) associated with Dandy-Walker malformation, abnormal skull development and microcephaly. *Taiwan J Obstet Gynecol* 2010;49:320–6.
- Shapira SK, McCaskill C, Northrup H, et al. Chromosome 1p36 deletions: The clinical phenotype and molecular characterization of a common newly delineated syndrome. *Am J Hum Genet* 1997;61:642–50.
- Shaffer LG, Lupski JR. Molecular mechanisms for constitutional chromosomal rearrangements in humans. *Annu Rev Genet* 2000;34:297–329.
- Battaglia A, Shaffer LG. 1p36 deletion syndrome. In: Pagon RA, Bird TC, Dolan CR, Stephens K, eds. *Gene Reviews* University of Washington, Seattle, 1993–2008. Available at: <http://www.ncbi.nlm.nih.gov/bookshelf/br.fcgi?book=gene&part=del1p36>. [Date accessed: August 9, 2010]
- Battaglia A, Hoyme HE, Dallapiccola B, et al. Further delineation of deletion 1p36 syndrome in 60 patients: A recognizable phenotype and common cause of developmental delay and mental retardation. *Pediatrics* 2008;121:404–10.
- Gajicka M, Mackay KL, Shaffer LG. Monosomy 1p36 deletion syndrome. *Am J Med Genet Part C Semin Med Genet* 2007;145C:346–56.
- Howard-Peebles PN, Black SH. FISH identification of a deletion 1p36 from a half-cryptic maternal translocation (March of Dimes 25th Clinical Genetics Conference and American College of Medical Genetics 1st Annual Meeting Abstract). *Am J Med Genet* 1994;52:381. [Abstract C-14]
- Campeau PM, Ah Mew N, Cartier L, Mackay KL, Shaffer LG, Der Kaloustian VM, Thomas MA. Prenatal diagnosis of monosomy 1p36: a focus on brain abnormalities and a review of the literature. *Am J Med Genet* 2008;146A:3062–9.
- Oppenheimer S, Dignan P, Soukup S. Partial trisomy 20p: familial occurrence. *Am J Med Genet* 2000;95:316–9.
- Thomas MA, Duncan AM, Bardin C, Der Kaloustian VM. Lissencephaly with der(17)t(17;20)(p13.3;p12.2)mat. *Am J Med Genet* 2004;124A:292–5.
- Chen Z, Grebe TA, Guan XY, et al. Maternal balanced translocation leading to partial duplication of 4q and partial deletion of 1p in a son: Cytogenetic and FISH studies using band-specific painting probes generated by chromosome microdissection. *Am J Med Genet* 1997;71:160–6.
- Kulharya AS, Carlin ME, Stettler RW, Huslig M, Kukolich MK, Garcia-Heras J. Prenatal diagnosis of a *de novo* trisomy 6q22.2→6qter and monosomy 1pter→1p36.3. Case report with a 2-year follow-up and a brief review of other prenatal cases of partial trisomy 6q. *Clin Genet* 1997;51:115–7.
- Robbins-Furman P, Elder FFB, Mastrobattista JM, Northrup H, Shapira SK. Prenatal diagnosis of deletion 1p36 syndrome. *Am J Hum Genet* 1997;61:A139.
- Faivre L, Morichon-Delvallez N, Viot G, et al. Prenatal detection of a 1p36 deletion in a fetus with multiple malformations and a review of the literature. *Prenat Diagn* 1999;19:49–53.
- Cavani S, Perfumo C, Faravelli F, et al. Cryptic 1p36.3/6q25.2 translocation in three generations ascertained through a foetus with IUGR and cerebral malformations. *Prenat Diagn* 2003;23:819–23.
- Heilstedt HA, Ballif BC, Howard LA, Kashork CD, Shaffer LG. Population data suggest that deletions of 1p36 are a relatively common chromosome abnormality. *Clin Genet* 2003;64:310–6.
- Reddy KS, Yang X. Submicroscopic terminal deletion of 1p36.3 and Xp23 hidden in complex chromosome rearrangements: Independent mechanism of telomere restitution on the two chromatids. *Am J Med Genet* 2003;117A:261–7.
- Hsieh LJ, Hsieh TC, Yeh GP, Lin MI, Chen M, Wang BB. Prenatal diagnosis of a fetus affected with Down syndrome and deletion 1p36 syndrome by fluorescence *in situ* hybridization and spectral karyotyping. *Fetal Diagn Ther* 2004;19:356–60.
- Lissauer D, Larkins SA, Sharif S, MacPherson L, Rhodes C, Kilby MD. Prenatal diagnosis and prenatal imaging features of fetal monosomy 1p36. *Prenat Diagn* 2007;27:874–8.
- Vialard F, Molina Gomes D, Leroy B, et al. Array comparative genomic hybridization in prenatal diagnosis: another experience. *Fetal Diagn Ther* 2009;25:277–84.
- Chen CP, Su YN, Chang TY, Chern SR, Tsai FJ, Hwang JK, Wang W. 22q11.2 microdeletion in a fetus with double-outlet right ventricle, pulmonary stenosis and a ventricular septal defect: prenatal diagnosis by array comparative genomic hybridization. *Taiwan J Obstet Gynecol* 2009;48:437–40.
- Chen CP, Su YN, Tsai FJ, et al. Terminal 2q deletion and distal 15q duplication: prenatal diagnosis by array comparative genomic hybridization using uncultured amniocytes. *Taiwan J Obstet Gynecol* 2009;48:441–5.