

Short Communication

Mosaic deletion-duplication syndrome of chromosome 3: Prenatal molecular cytogenetic diagnosis using cultured and uncultured amniocytes and association with fetoplacental discrepancy

Chih-Ping Chen^{a,b,c,d,e,f,*}, Yi-Ning Su^g, Chin-Yuan Hsu^a, Schu-Rern Chern^b, Chen-Chi Lee^a, Yu-Ting Chen^b, Wen-Lin Chen^a, Wayseen Wang^{b,h}

^aDepartment of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^bDepartment of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^cDepartment of Biotechnology, Asia University, Taichung, Taiwan

^dSchool of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^eInstitute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^fDepartment of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^gDepartment of Medical Genetics, National Taiwan University Hospital, Taipei, Taiwan

^hDepartment of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 15 August 2011

Abstract

Objective: To present prenatal molecular cytogenetic diagnosis of mosaicism for terminal 3p deletion and distal 3q duplication using cultured and uncultured amniocytes, and the association with fetoplacental discrepancy.

Materials, Methods, and Results: A 35-year-old primigravid woman was referred for genetic counseling at 21 weeks of gestation because of 20% (5/25 colonies) mosaicism for add(3)(p26) detected by amniocentesis. Repeated amniocenteses were performed. Array comparative genomic hybridization (aCGH) and interphase fluorescence *in situ* hybridization (FISH) were applied in the uncultured amniocytes. aCGH analysis detected 0.15-Mb microdeletion of 3p26.3 with *CHL1* haploinsufficiency and a 49.42-Mb duplication of 3q24-q29 in the uncultured amniocytes. Interphase FISH analysis revealed 27.3% mosaicism (12/44 cells) in the uncultured amniocytes. Metaphase FISH analysis revealed 23.3% mosaicism (7/30 cells) in the cultured amniocytes. Conventional cytogenetic analysis showed a karyotype of 46,XX,der(3)(qter → q24::p26.3 → qter)[10]/46,XX[20] (33% mosaicism). Subsequent fetal blood sampling showed a karyotype of 46,XX,der(3)(qter → q24::p26.3 → qter)[5]/46,XX[35] (12.5% mosaicism). The parents elected to terminate the pregnancy, and a malformed fetus was delivered at 24 weeks of gestation with characteristic facial dysmorphism and clinodactyly of the hands. Cytogenetic analysis of the extra-embryonic tissues revealed the results of 46,XX (40 cells) in placenta, 25% mosaicism (10/40 cells) in amniotic membrane and 50% mosaicism (20/40 cells) in umbilical cord.

Conclusion: Our presentation highlights the utility of molecular cytogenetic technologies in prenatal diagnosis of rare mosaic chromosome rearrangements and provides evidence for fetoplacental discrepancy under such circumstances.

Copyright © 2011, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: 3p deletion; 3q duplication; *CHL1*; Deletion-duplication syndrome of chromosome 3; Fetoplacental discrepancy; Mosaicism

Introduction

The deletion-duplication syndrome of chromosome 3 consists of partial 3p deletion and partial 3q duplication. Both 3p deletion syndrome and 3q duplication syndrome are rare but clinically recognizable chromosome abnormalities.

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

Clinical reports with deletion-duplication syndrome of chromosome 3 resulting from familial pericentric inversions of chromosome 3 have been well described. Most cases resulted from a pericentric inversion with breakpoints at 3p and 3q in the parents. These cases include $\text{dup}(3)(\text{q21} \rightarrow \text{qter})$ & $\text{del}(3)(\text{pter} \rightarrow \text{p25})$ [1–5], $\text{dup}(3)(\text{q23} \rightarrow \text{qter})$ & $\text{del}(3)(\text{pter} \rightarrow \text{p25})$ [6–9] and $\text{dup}(3)(\text{q25} \rightarrow \text{qter})$ & $\text{del}(3)(\text{pter} \rightarrow \text{p25})$ [10–12].

Prabhakara et al. [13] first reported prenatal diagnosis of deletion-duplication syndrome of chromosome 3 in a fetus with lumbosacral meningocele, a karyotype of $46,\text{XY},\text{add}(3)(\text{p26})$ and a familial pericentric inversion of $46,\text{XY},\text{inv}(3)(\text{p26q21})$ in the father. In that case, molecular cytogenetic investigations revealed $\text{dup}(3)(\text{q21} \rightarrow \text{qter})$ & $\text{del}(3)(\text{pter} \rightarrow \text{p26.1})$ in the fetus. Prenatal diagnosis of mosaic deletion-duplication syndrome of chromosome 3 has not previously been described. Here, we present such a case in which a cytogenetic discrepancy between the fetus and the placenta was noted. Our presentation highlights the utility of high-resolution molecular cytogenetic technologies in prenatal diagnosis of mosaic chromosome rearrangements using both cultured and uncultured amniocytes.

Materials, methods and results

A 35-year-old primigravid woman was referred for genetic counseling at 21 weeks of gestation because of an abnormal

amniocentesis result of mosaicism for a derivative chromosome 3 of $\text{add}(3)(\text{p26})$. She had undergone amniocentesis at 20 weeks of gestation in an obstetric clinic because of advanced maternal age. Amniocentesis revealed a karyotype of $46,\text{XX},\text{add}(3)(\text{p26})[5 \text{ colonies}]/46,\text{XX}[20 \text{ colonies}]$. Among 25 colonies of cultured amniocytes, five colonies had a derivative chromosome 3 [$\text{der}(3)$], of which additional maternal was attached to band 3p26, whereas the other 20 colonies had a normal karyotype. Repeated amniocentesis was performed at 21 weeks of gestation. Oligonucleotide-based array comparative genomic hybridization (aCGH) using CytoChip Oligo Array (BlueGnome, Cambridge, UK) and interphase fluorescence *in situ* hybridization (FISH) were applied in the uncultured amniocytes. The aCGH analysis rapidly detected gene dosage decrease at 3p26.3 and gene dosage increase at 3q24–q29 in the uncultured amniocytes (Fig. 1A). There were a 0.15-Mb microdeletion of 3p26.3 (134,741–287,068 bp) and a 49.42-Mb duplication of 3q24–q29 (149,900,377–199,321,417 bp) (Fig. 1B) [UCSC genome browser on March 2006 (NCBI 36/hg 18) assembly].

The interphase FISH analysis using chromosome 3p- and 3q-specific bacterial artificial chromosome (BAC) clone probes RP11-210M17 (3p26.3: 129,717–287,700 bp) (spectrum red) and RP11-338O10 (3q29: 195,287,280–195,356,374 bp) (spectrum green) revealed 27.3% (12/44) mosaicism for the $\text{der}(3)$. Among 44 interphase uncultured amniocytes analyzed, 12 cells (27.3%) showed three green signals and one red signal




Fig. 1. (A) Oligonucleotide-based array comparative genomic hybridization (aCGH) using CytoChip Oligo Array (BlueGnome, Cambridge, UK) in uncultured amniocytes shows gene dosage decrease at 3p26.3 and gene dosage increase at 3q24–q29 (arrows). (B) aCGH shows a 0.15-Mb microdeletion of 3p26.3 (134,741–287,068 bp) and a 49.42-Mb duplication of 3q24–q29 (149,900,377–199,321,417 bp).




Fig. 2. Interphase fluorescence *in situ* hybridization (FISH) analysis using bacterial artificial chromosome (BAC) clone probes RP11-210M17 (3p26.3: 129,717–287,700 bp) (spectrum red) and RP11-338O10 (3q29: 195,287,280–195,356,374 bp) (spectrum green) shows (A) three green signals and one red signal in the abnormal uncultured amniocytes with a derivative chromosome 3 [der(3)], and (B) two red signals and two green signals in the normal uncultured amniocytes.




Fig. 3. Metaphase FISH analysis of cultured amniocytes using BAC clone probes RP11-210M17 and RP11-338O10 shows (A) an abnormal amniocyte with a normal chromosome 3 containing one green signal and one red signal, and a der(3) containing two green signals, and (B) a normal amniocyte with two normal chromosomes 3.

consistent with the diagnosis of 3q duplication and 3p deletion, whereas the other 32 cells were normal (Fig. 2). The metaphase FISH analysis of cultured amniocytes using BAC clone probes RP11-210M17 and RP11-338O10 revealed 23.3% (7/30)




Fig. 4. Metaphase FISH analysis of the abnormal cultured amniocytes using BAC clone probes RP11-306H5 (spectrum green) (3p26.3, encompassing *CHL1* gene), RP11-129K1 (spectrum red) (3p26.3, encompassing *CNTN4* gene) and RP11-338O10 (spectrum yellow) (3q29) shows a normal chromosome 3 containing one green signal, one red signal and one yellow signal, and a der(3) containing two yellow signals and one red signal. The result indicates *CHL1* haploinsufficiency but presence of the *CNTN4* gene in the der(3).

mosaicism for the derivative chromosome 3 (Fig. 3). The metaphase FISH analysis of the abnormal cultured amniocytes using BAC clone probes RP11-306H5 (3p26.3: 325,397–326,005 bp) (spectrum green) encompassing *CHL1* gene, RP11-129K1 (3p26.3: 2,215,034–2,377,366 bp) (spectrum red) encompassing *CNTN4* gene and RP11-338O10 (3q29) showed *CHL1* haploinsufficiency but presence of the *CNTN4* gene in the der(3) (Fig. 4). Among 30 metaphase cultured amniocytes analyzed, seven cells (23.3%) showed 3p deletion and 3q duplication, whereas the other 23 cells were normal. Conventional cytogenetic analysis showed a karyotype of 46,XX,der(3)(qter → q24::p26.3 → qter)[10 colonies]/46,XX[20 colonies] (33% mosaicism) (Fig. 5).

Prenatal ultrasound findings were unremarkable. Subsequent fetal blood sampling showed a karyotype of 46,XX,der(3)(qter → q24::p26.3 → qter)[5 cells]/46,XX[35 cells] (12.5% mosaicism). The parents elected to terminate the pregnancy, and a 616-g female fetus was delivered at 24 weeks of gestation with a triangular face, hypertelorism, depressed nasal bridge, anteverted nostrils, long philtrum, down-turned mouth, low-set ears and clinodactyly of the hands (Fig. 6). Cytogenetic analysis of the extraembryonic tissues revealed the results of 46,XX[40 cells] in placenta, 46,XX,der(3)(qter → q24::p26.3 → qter)[10 cells]/46,XX[30 cells] (25% mosaicism) in amniotic membrane and 46,XX,der(3)(qter → q24::p26.3 → qter)[20 cells]/46,XX[20 cells] (50%



Fig. 5. (A) A karyotype of 46,XX. (B) A karyotype of 46,XX,der(3)(qter → q24::p26.3 → qter).




Fig. 6. (A) and (B) Craniofacial appearance of the fetus at birth.

mosaicism) in umbilical cord. Polymorphic DNA marker analysis excluded uniparental disomy 3.

Discussion

Prabhakara et al. [13] first reported prenatal diagnosis of a malformed fetus with deletion-duplication of chromosome 3 arising from meiotic recombination of a familial pericentric inversion. In a literature review of patients with partial 3q duplication and partial 3p deletion, Steinbach et al. [12] suggested that deletion-duplication of chromosome 3 is a result of meiotic recombination. The presence of a normal cell line in our case is consistent with the postzygotic origin of the recombinant chromosome 3 in the fetus, and provides evidence that deletion-duplication of chromosome 3 can present in a mosaic form and arise from a postzygotic mitotic event.

The present case demonstrates a fetoplacental discrepancy in mosaic deletion-duplication syndrome of chromosome 3, and shows a limitation of using placenta as a diagnostic tool for prenatal diagnosis of mosaic chromosome rearrangements. Our case reinforces the notion that amniocentesis offers a more reliable diagnosis of chromosome aberrations compared to chorionic villus sampling.

The present case shows that oligonucleotide-based aCGH using CytoChip Oligo Array is able to detect genomic imbalance in uncultured amniocytes with 27.3% (12/44) mosaicism for terminal 3p deletion and distal 3q duplication. We previously demonstrated that BAC-based aCGH using CMDX BAC aCGH 2500 Chips was not able to detect genomic imbalance in uncultured amniocytes with 18% (9/50) mosaicism for trisomy 9 [14], and aCGH using CMDX BAC aCGH 3000 Chips and Oligo HD Scan (CMDX, Irvine, CA, USA) was not able to detect genomic imbalance in uncultured amniocytes with 25% (5/20) mosaicism for trisomy 8 [15]. We previously also demonstrated that CytoChip Oligo Array (BlueGnome, Cambridge, UK) was able to detect genomic imbalance in uncultured amniocytes with 12% (6/50) mosaicism for trisomy 2 [16]. It is likely that variation in the ability

of detection of low-level mosaicism in uncultured amniocytes exists in different commercial kits of aCGH.

The 3p deletion syndrome is a rare contiguous gene syndrome involving chromosome 3p25-p26 deletion and is characterized by low birth weight, psychomotor delay, growth retardation and characteristic facial dysmorphism of micro- and brachycephaly, flat occiput, triangular face, hypertelorism, epicanthic folds, ptosis, broad and short nose, long philtrum, down-turned mouth, micrognathia, low-set ears, hypertrichosis and synophrys [17–21]. Fernandez et al. [22] suggested that *CNTN4* disruption is associated with 3p deletion syndrome phenotype and abnormal central nervous system development. Dijkhuizen et al. [23] suggested that loss of *CNTN4* and *CRBN* contributes to mental retardation in 3p deletion syndrome. There was no loss of the genes of *CNTN4* and *CRBN* in this case. The 3p deletion in this case encompassed only a 0.15-Mb region of terminal 3p26.3 overlapping part of the *CHL1* gene (213,650–426,097 bp). The *CHL1* gene encodes cell adhesion molecule L1-like (CALL or CAL1) protein which is highly expressed in central and peripheral nervous system [24,25]. Frints et al. [25] and Shrimpton et al. [26] suggested that interruption or loss of the *CHL1* gene may cause cognitive deficit. Pohjola et al. [27] recently reported a family with a 1.1-Mb terminal 3p deletion encompassing only the *CHL1* gene and was associated with nonspecific features, mild learning difficulty, microcephaly and growth retardation. They suggested that a small terminal deletion comprising only the *CHL1* gene may not be related to dysmorphic features and profound mental retardation but may cause only a mild mental deficit.

The 3q duplication syndrome is a well-described chromosome disorder that has the clinical overlap with Cornelia de Lange syndrome, such as developmental delay, failure to thrive, low anterior hairline, prominent eyelashes, depressed nasal bridge, anteverted nares, long prominent philtrum, micrognathia, rhizomelic shortening of the limbs and genital hypoplasia; and other features such as normal birth weight, bushy eyebrows, ocular hypertelorism, up-slanting palpebral fissures, epicanthic folds, a broad nose and normal lips [28].

Partial duplication of 3q can be associated with major congenital malformations, such as omphalocele [6,29–34], cystic hygroma [35,36], holoprosencephaly [37], spina bifida or sacral dimple [3], lumbosacral meningocele [13], Dandy-Walker malformation [38], and cerebellar hypoplasia and agenesis of the corpus callosum [39]. The minimal critical region for most of the clinical features of 3q duplication syndrome has been narrowed down to 3q26.3–q29 [40–45]. The 3q duplication in this case encompassed a 49.42-Mb duplication of 3q24–q29 overlapping the critical region of 3q duplication syndrome, and was associated with characteristic craniofacial dysmorphism of 3q duplication syndrome.

In conclusion, by using FISH and aCGH in cultured and uncultured amniocytes, we are able to precisely define a mosaic chromosome rearrangement in a fetus with concomitant 3p deletion syndrome and 3q duplication syndrome. Our presentation highlights the utility of molecular cytogenetic technologies in prenatally detected rare mosaic chromosome rearrangements and provides evidence for fetoplacental discrepancy under such circumstances.

Acknowledgments

This work was supported by research grants NSC-97-2314-B-195-006-MY3 and NSC-99-2628-B-195-001-MY3 from the National Science Council, and MMH-E-100-04 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Falek A, Schmidt R, Jervis GA. Familial de Lange syndrome with chromosome abnormalities. *Pediatrics*. 1966;37:92–101.
- [2] Boué J, Hirschhorn K, Lucas M, Gautier M, Moszer M, Bach C. Aneusomies de recombinaison. Conséquence d'une inversion pericentrique d'un chromosome 3 paternel. *Ann Pediatr*. 1974;21:567–73.
- [3] Allderdice PW, Browne N, Murphy DP. Chromosome 3 duplication q21 leads to qter deletion p25 leads to pter syndrome in children of carriers of a pericentric inversion inv(3) (p25q21). *Am J Hum Genet*. 1975;27:699–718.
- [4] Patil S, Hanson J, Zellweger H. Duplication-deletion of chromosome 3 syndrome. *Am J Hum Genet*. 1978;30:A89.
- [5] Kawashima H, Maruyama S. A case of chromosome 3 duplication q deletion p syndrome born to the mother with a pericentric inversion, inv(3)(p25q21). *Jinrui Idengaku Zasshi*. 1979;24:9–12.
- [6] Mulcahy MT, Pemberton PJ, Sprague P. Trisomy 3q: two clinically similar but cytogenetically different cases. *Ann Génét*. 1979;22:217–20.
- [7] Sutherland GR, Mulley JC, Goldblatt E. A large kindred with an inv(3)(p25q23): clinical, cytogenetic and genetic marker studies. *Ann Génét*. 1981;24:202–5.
- [8] Preus M, Vekemans M, Kaplan P. Diagnosis of chromosome 3 duplication q23 → qter, deletion p25 → pter in a patient with the C (trigonocephaly) syndrome. *Am J Med Genet*. 1986;23:935–43.
- [9] Polívková Z, Kucerová M, Dolanská M, Hresová M, Kofer J. Similarity between the syndrome of partial trisomy of the long arm of chromosome 3 and the Cornelia de Lange syndrome. *Cesk Pediatr*. 1988;43:723–6 [Czech].
- [10] Fineman RM, Hecht F, Ablow RC, Howard RO, Breg WR. Chromosome 3 duplication q/deletion p syndrome. *Pediatrics*. 1978;61:611–8.
- [11] Pope IS, Thuline HC, Aronson MM, Bozarth B, Greene AE, Coriell LL. Duplication of a segment of chromosome 3 in a subject with multiple congenital anomalies and a 47, XYY father, inversion of chromosomes 3 and 9 in the mother, and inversion of chromosome 9 in a brother. *Cytogenet Cell Genet*. 1979;24:127–8.
- [12] Steinbach P, Adkins Jr WN, Caspar H, Dumars KW, Gebauer J, Gilbert EF, et al. The dup(3q) syndrome: report of eight cases and review of the literature. *Am J Med Genet*. 1981;10:159–77.
- [13] Prabhakara K, Bruno DL, Padman P, Prasad S, Sudheer Kumar R, et al. Prenatal detection of deletion-duplication of chromosome 3 arising from meiotic recombination of a familial pericentric inversion. *Prenat Diagn*. 2008;28:466–8.
- [14] Chen C-P, Lin H-M, Su Y-N, Chern S-R, Tsai F-J, Wu P-C, et al. Mosaic trisomy 9: prenatal diagnosis and molecular genetic analyses. *Taiwan J Obstet Gynecol*. 2010;49:341–50.
- [15] Chen C-P, Chen M, Pan Y-J, Su Y-N, Chern S-R, Tsai F-J, et al. Prenatal diagnosis of mosaic trisomy 8: clinical report and literature review. *Taiwan J Obstet Gynecol*. 2011;50:331–8.
- [16] Chen C-P, Su Y-N, Lin S-Y, Chern S-R, Chen Y-T, Lee M-S, et al. Prenatal diagnosis of mosaic trisomy 2: discrepancy between molecular cytogenetic analyses of uncultured amniocytes and karyotyping of cultured amniocytes in a pregnancy with severe fetal intrauterine growth restriction. *Taiwan J Obstet Gynecol*. 2011;50:390–3.
- [17] Verjaal M, De Nef MB. A patient with a partial deletion of the short arm of chromosome 3. *Am J Dis Child*. 1978;132:43–5.
- [18] Narahara K, Kikkawa K, Murakami M, Hiramoto K, Namba H, Tsuji K, et al. Loss of the 3p25.3 band is critical in the manifestation of del(3p) syndrome: karyotype-phenotype correlation in cases with deficiency of the distal portion of the short arm of chromosome 3. *Am J Med Genet*. 1990;35:269–73.
- [19] Kariya S, Aoji K, Akagi H, Fukushima K, Chikumoto E, Ogawa T, et al. A terminal deletion of the short arm of chromosome 3: karyotype 46, XY, del(3)(p25-pter); a case report and literature review. *Int J Pediatr Otorhinolaryngol*. 2000;56:71–8.
- [20] Cargile CB, Goh DL-M, Goodman BK, Chen X-N, Korenberg JR, Semenza GL, et al. Molecular cytogenetic characterization of a subtle interstitial del(3)(p25.3p26.2) in a patient with deletion 3p syndrome. *Am J Med Genet*. 2002;109:133–8.
- [21] Chen C-P, Lin S-P, Ho C-S, Chern S-R, Lee C-C, Chen W-L, et al. Distal 3p monosomy associated with epilepsy in a boy. *Genet Couns*. 2005;16:429–32.
- [22] Fernandez T, Morgan T, Davis N, Klin A, Morris A, Farhi A, et al. Disruption of *contactin 4* (*CNTN4*) results in developmental delay and other features of 3p deletion syndrome. *Am J Hum Genet*. 2004;74:1286–93.
- [23] Dijkhuizen T, van Essen T, van der Vlies P, Verheij JBG, Sikkema-Raddatz B, van der Veen AY, et al. FISH and array-CGH analysis of a complex chromosome 3 aberration suggests that loss of *CNTN4* and *CRBN* contributes to mental retardation in 3pter deletions. *Am J Med Genet*. 2006;140A:2482–7.
- [24] Wei M-H, Karavanova I, Ivanov SV, Popescu NC, Keck CL, Pack S, et al. In silico-initiated cloning and molecular characterization of a novel human member of the *LI* gene family of neural cell adhesion molecules. *Hum Genet*. 1998;103:355–64.
- [25] Frints SGM, Marynen P, Hartmann D, Fryns J-P, Steyaert J, Schachner M, et al. CALL interrupted in a patient with non-specific mental retardation: Gene dosage-dependent alteration of murine brain development and behavior. *Hum Mol Genet*. 2003;12:1463–74.
- [26] Shrimpton AE, Jensen KA, Hoo JJ. Karyotype-phenotype analysis and molecular delineation of a 3p26 deletion/8q24.3 duplication case with a virtually normal phenotype and mild cognitive deficit. *Am J Med Genet*. 2006;140A:388–91.
- [27] Pohjola P, de Leeuw N, Penttinen M, Kääriäinen H. Terminal 3p deletions in two families — correlation between molecular karyotype and phenotype. *Am J Med Genet*. 2010;152A:441–6.
- [28] Deardorff MA, Clark DM, Krantz ID. Cornelia de Lange Syndrome. In: Pagon RA, Bird TD, Dolan CR, Stephens K, editors. *GeneReviews* [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2005 Sep 16 [updated 2006 Aug 14].
- [29] Chen C-P, Liu F-F, Jan S-W, Chen C-P, Lan C-C. Partial duplication of 3q and distal deletion of 11q in a stillbirth with an omphalocele

- containing the liver, short limbs, and intrauterine growth retardation. *J Med Genet* 1996;33:615–7.
- [30] Chen C-P, Lee C-C, Chuang C-Y, Town D-D, Lee M-S, Chen M-H. Recurrent omphalocele with partial trisomy 3q and partial monosomy 11q. *Clin Genet* 1997;52:196–8.
- [31] Chen C-P. Inconsistency of omphalocele contents in three consecutive siblings with partial trisomy 3q and partial monosomy 11q. *Prenat Diagn* 1999;19:591.
- [32] Chen C-P. Chromosomal abnormalities associated with omphalocele. *Taiwan J Obstet Gynecol* 2007;46:1–8.
- [33] Cinti R, Botta G, Asnaghi V, Del Monaco A, Salvego M, Silengo M. *De novo* partial duplication of 3q and distal deletion of 20p in a 15-week abortus with omphalocele. *Fetal Diagn Ther* 2000;15:61–2.
- [34] Yatsenko SA, Mendoza-Londono R, Belmont JW, Shaffer LG. Omphalocele in trisomy 3q: further delineation of phenotype. *Clin Genet* 2003;64:404–13.
- [35] Chen C-P. Prenatal diagnosis of partial trisomy 3q(3q22 → qter) and monosomy 6q(6q25.3 → qter) in a fetus with sonographic findings of cystic hygroma colli and unilateral pleural effusion. *Prenat Diagn* 2001;21:73.
- [36] Pires A, Ramos L, Venâncio M, Rei AI, Castedo S, Saraiva J. Prenatal foetal diagnosis of partial trisomy 3q and monosomy 13p due to a maternal balanced rearrangement. *Prenat Diagn* 2005;25:292–5.
- [37] Chen C-P, Liu F-F, Jan S-W, Lin C-L, Lan C-C. Prenatal diagnosis of terminal deletion 7q and partial trisomy 3p in fetuses with holoprosencephaly. *Clin Genet* 1996;50:321–6.
- [38] de Azevedo Moreira LM, Neri FB, de Quadros Uzêda S, de Carvalho AFL, Santana GC, Souza FR, et al. Multiple congenital malformations including severe eye anomalies and abnormal cerebellar development with Dandy-Walker malformation in a girl with partial trisomy 3q. *Ophthalmic Genet* 2005;26:37–43.
- [39] Roberts N, Elmslie F, Rich P, Johnson R, Bhide A. Prenatal cranial ultrasound features of duplication chromosome 3q(21-24). *Prenat Diagn* 2006;26:293–5.
- [40] Rizzu P, Baldini A. Subchromosomal band interval mapping and ordering of DNA markers in the region 3q26.3-q27 involved in the dup(3q) syndrome. *Genomics* 1994;24:580–2.
- [41] Aqua MS, Rizzu P, Lindsay EA, Shaffer LG, Zackai EH, Overhauser J, et al. Duplication 3q syndrome: molecular delineation of the critical region. *Am J Med Genet* 1995;55:33–7.
- [42] Faas BHW, de Vries BBA, van Es-van Gaal J, Merckx G, Draaisma JMT, Smeets DFCM. A new case of dup(3q) syndrome due to a pure duplication of 3qter. *Clin Genet* 2002;62:315–20.
- [43] Battaglia A, Novelli A, Ceccarini C, Carey JC. Familial complex 3q;10q rearrangement unraveled by subtelomeric FISH analysis. *Am J Med Genet* 2006;140A:144–50.
- [44] Grossmann V, Müller D, Muller W, Fresser F, Erdel M, Janecke AR, et al. Essentially "pure" trisomy 3q27 → qter: further delineation of the partial trisomy 3q phenotype. *Am J Med Genet* 2009;149A:2522–6.
- [45] Shanske AL, Leonard J, Nahum O, Coppock DL, Levy B. Delineation of the breakpoints of pure duplication 3q due to a *de novo* duplication event using SOMA. *Am J Med Genet* 2010;152A:3185–8.