

Research Letter

Magnetic resonance imaging demonstration of sirenomelia in one fetus of a dizygotic twin pregnancy conceived by intracytoplasmic sperm injection, *in vitro* fertilization and embryo transfer

Chih-Ping Chen^{a,b,c,d,e,f,*}, Chin-Yuan Hsu^a, Maw-Shuan Lee^g, Yu-Peng Liu^{h,i}, Fuu-Jen Tsai^{d,j},
Pei-Chen Wu^a, Schu-Rern Chern^b, Wayseen Wang^{b,k}

^a Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^c Department of Biotechnology, Asia University, Taichung, Taiwan

^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^g Institute of Medicine, Chung Shan Medical University, Taichung, Taiwan

^h Department of Radiology, Mackay Memorial Hospital Hsinchu Branch, Hsinchu, Taiwan

ⁱ Mackay Medicine, Nursing and Management College, Taipei, Taiwan

^j Departments of Medical Genetics, and Medical Research, China Medical University Hospital, Taichung, Taiwan

^k Department of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 25 October 2010

A 32-year-old, primigravid woman presented with a twin pregnancy at 21 weeks of gestation for evaluation of oligohydramnios in one co-twin. The woman and her husband were healthy and non-consanguineous. There was no family history of congenital malformations, and the woman did not have diabetes mellitus. The woman had suffered from bilateral tubal occlusion and primary infertility. This was her first pregnancy that was conceived by intracytoplasmic sperm injection (ICSI), *in vitro* fertilization (IVF) and embryonic transfer (ET). Four embryos had been implanted and two survived. Level II ultrasound at 21 weeks of gestation revealed a normal co-twin and an abnormal co-twin with marked oligohydramnios, and absent kidneys and urinary tracts. Magnetic resonance imaging (MRI) evaluation of the fetuses at 31 weeks of gestation revealed a normal co-twin and an anomalous co-twin with oligohydramnios, pulmonary hypoplasia, a small stomach, ill-defined small bowel loops, absent bladder, renal agenesis and a single leg consistent with the diagnosis of sirenomelia (Fig. 1). The pregnancy was uneventful until 35 weeks of gestation when premature rupture of the membranes occurred. A 2128-g normal female co-twin and a 1576-g abnormal co-twin with a single lower extremity without the

foot, an imperforate anus, absence of external genitalia and a single umbilical artery were delivered by cesarean section because of malpresentation. Postnatal X-ray examination of the abnormal infant showed sirenomelia with hypoplastic pelvis, absence of the right, lower limb and the left foot, and abnormal lumbosacral spine (Fig. 2). Cytogenetic analysis of the cord blood of the abnormal co-twin revealed a karyotype of 46,XX. A molecular zygosity test of the twins confirmed dizygotic twinning (Fig. 3). The abnormal co-twin died soon after birth (Fig. 4). The normal co-twin was doing well at the age of 4 months.

Sirenomelia is characterized by a complete or incomplete fusion of the lower extremities, imperforate anus and absent external genitalia, and can be associated with anomalies such as Potter syndrome, limb–body wall complex, pentology of Cantrell, esophageal atresia, hydrocephalus, holoprosencephaly, neural tube defects and VACTERL (vertebral segmentation defects, anal atresia/stenosis, cardiac malformation, tracheo-esophageal fistula and/or esophageal atresia, and renal and limb anomalies) association [1,2]. Sirenomelia has an overall incidence of 1.5–4.2/100,000 births and may be associated with maternal diabetes and monozygotic twinning [1,2]. About 2% of the cases with sirenomelia are associated with maternal diabetes, and there is a 100~150-fold increase in the incidence of sirenomelia in monozygotic twinning over that in singleton pregnancy and dizygotic twinning [1,2]. Recently, sirenomelia has been suggested as a primary defect

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).




Fig. 1. Magnetic resonance imaging (MRI) at 31 weeks of gestation demonstrating a fetus with oligohydramnios, pulmonary hypoplasia, a small stomach, ill-defined bowel loops, absent bladder, agenesis of kidneys and a single leg consistent with the diagnosis of sirenomelia.

of blastogenesis affecting multiple primordial fields originating from the caudal mesenchyme [3–6].

We previously described limb–body wall complex in one fetus of a dizygotic twin pregnancy conceived by IVT-ET [7]. In this report, we additionally describe sirenomelia in one fetus of a dizygotic twin pregnancy conceived by ICSI and IVT-ET. Monozygotic twinning is well known to be associated with early embryonic structural developmental defects such as




Fig. 2. Postnatal X-ray showing sirenomelia with hypoplastic pelvis, absent right lower limb, absent left foot and an abnormal lumbosacral spine with sacrococcygeal dysgenesis.

D13S790




Fig. 3. Molecular zygosity test of the twins shows dizygotic twinning.

sirenomelia, holoprosencephaly, anencephaly, extrophy of the cloaca malformation and VATERR (vertebral defects, anal atresia, tracheo-esophageal fistula, esophageal atresia, and radial and renal dysplasia) association [8]. The majority of monozygotic twins with early embryonic structural developmental defects are discordant for anomalies, and only 5–20% of the cases are concordant [9]. About 15% of the cases with sirenomelia are reported to be associated with twin pregnancies, most often monozygotic twinning [10]. However, the present case was a dizygotic twin pregnancy. Dizygotic twin pregnancy has been shown to be associated with structural abnormalities in one fetus [11,12]. Sirenomelia has been reported in dizygotic twin pregnancy [13–16]. Sirenomelia has also been reported in the product of ICSI [17].



Fig. 4. The abnormal fetus with sirenomelia.

Sirenomelia in our case was likely to be related to developmental defects of the fetus. This case provides additional evidence for the occurrence of sirenomelia in multiple pregnancies after ICSI and IVF-ET with no influence of monozygosity and invasive prenatal diagnostic procedures of the pregnant woman.

Acknowledgments

This work was supported by research grants NSC-96-2314-B-195-008-MY3 and NSC-97-2314-B-195-006-MY3 from the National Science Council, and MMH-E-99004 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Chen C-P, Shih S-L, Liu F-F, Jan S-W. Cebocephaly, alobar holoprosencephaly, spina bifida, and sirenomelia in a stillbirth. *J Med Genet* 1997;34:252–5.
- [2] Chen C-P, Shih S-L, Jan S-W, Lin Y-N. Sirenomelia with an uncommon osseous fusion associated with a neural tube defect. *Pediatr Radiol* 1998;28:293–6.
- [3] Dueterhoeft SM, Ernst LM, Siebert JR, Kapur RP. Five cases of caudal regression with an aberrant abdominal umbilical artery: further support for a caudal regression-sirenomelia spectrum. *Am J Med Genet* 2007;143A:3175–84.
- [4] Rougemont A-L, Bouron-Dal Soglio D, Déilets V, Jovanovic M, Perreault G, Laurier Oligny L, et al. Caudal dysgenesis, sirenomelia, and situs inversus totalis: a primitive defect in blastogenesis. *Am J Med Genet* 2008;146A:1470–6.
- [5] Castori M, Silvestri E, Cappellacci S, Binni F, Sforzolini GS, Grammatico P. Sirenomelia and VACTERL association in the offspring of a woman with diabetes. *Am J Med Genet* 2010;152A:1803–7.
- [6] Thottungal AD, Charles AK, Dickinson JE, Bower C. Caudal dysgenesis and sirenomelia-single centre experience suggests common pathogenic basis. *Am J Med Genet* 2010;152A:2578–87.
- [7] Chen C-P, Lee M-S, Tsai F-J, Huang M-C, Chern S-R, Wang W. Limb-body wall complex in one fetus of a dizygotic twin pregnancy obtained by egg donation, *in vitro* fertilization and embryo transfer: prenatal diagnosis and literature review. *Taiwan J Obstet Gynecol* 2009;48:446–50.
- [8] Schinzel AAGL, Smith DW, Miller JR. Monozygotic twinning and structural defects. *J Pediatr* 1979;95:921–30.
- [9] Jones KL. Monozygotic twinning and structural defects – General. In: Jones KL, editor. *Smith's recognizable patterns of human malformation*. Philadelphia: Elsevier Saunders; 2006. p. 748–51.
- [10] Di Lorenzo M, Brandt ML, Veilleux A. Sirenomelia in an identical twin: a case report. *J Pediatr Surg* 1991;26:1334–6.
- [11] Chen C-P, Liu Y-P, Chang T-Y, Tsai F-J, Chen C-Y, Wu P-C, et al. Prenatal diagnosis of persistent cloaca with hydrometrocolpos and ascites by magnetic resonance imaging in one fetus of a dizygotic twin pregnancy. *Taiwan J Obstet Gynecol* 2010;49:385–6.
- [12] Chen C-P, Su Y-N, Chang T-Y, Liu Y-P, Tsai F-J, Chen M-R, et al. Prenatal diagnosis of rhabdomyomas and cerebral tuberos sclerosis by magnetic resonance imaging in one fetus of a dizygotic twin pregnancy associated with a frameshift mutation in the *TSC2* gene. *Taiwan J Obstet Gynecol* 2010;49:387–9.
- [13] Schwaibold H, Oehler U, Helpap B, Bohm N. Sirenomelia and anencephaly in one of dizygotic twins. *Teratology* 1986;34:243–7.
- [14] Managoli S, Chaturvedi P, Vilhekar KY, Iyenger J, Dutta V, Gagane N. Sirenomelia dipus in a dizygotic twin. *Indian J Pediatr* 2002;69:1093–5.
- [15] Assimakopoulos E, Athanasiadis A, Zafrakas M, Dragoumis K, Bontis J. Caudal regression syndrome and sirenomelia in only one twin in two diabetic pregnancies. *Clin Exp Obstet Gynecol* 2004;31:151–3.
- [16] Nisenblat V, Leibovitz Z, Paz B, Shapiro I, Degani S, Ohel G. Dizygotic twin pregnancy discordant for sirenomelia. *J Ultrasound Med* 2007;26:97–103.
- [17] Bakhtar O, Benirschke K, Masliah E. Sirenomelia of an intracytoplasmic sperm injection conceptus: a case report and review of mechanism. *Pediatr Dev Pathol* 2006;9:245–53.