

Case Report

Rapid detection of *de novo* P253R mutation in *FGFR2* using uncultured amniocytes in a pregnancy affected by polyhydramnios, Blake's pouch cyst, and Apert syndrome

Chih-Ping Chen^{a,b,c,d,e,f,g,*}, Yi-Ning Su^h, Tung-Yao Changⁱ, Schu-Rern Chern^b,
Chen-Yu Chen^a, Jun-Wei Su^{a,j}, Wayseen Wang^{b,k}

^aDepartment of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^bDepartment of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^cDepartment of Medicine, Mackay Medical College, New Taipei City, Taiwan

^dDepartment of Biotechnology, Asia University, Taichung, Taiwan

^eSchool of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^fInstitute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^gDepartment of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^hDepartment of Obstetrics and Gynecology, School of Medicine, Taipei Medical University, Taipei, Taiwan

ⁱTaiji Fetal Medicine Center, Taipei, Taiwan

^jDepartment of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan

^kDepartment of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 22 January 2013

Abstract

Objective: To present prenatal ultrasound and molecular genetic diagnosis of Apert syndrome.

Case Report: A 30-year-old, gravida 3, para 2 woman was referred for genetic counseling at 32 weeks of gestation because of polyhydramnios and craniofacial and digital abnormalities in the fetus. She had undergone amniocentesis at 18 weeks of gestation because of maternal anxiety. Results of amniocentesis revealed a karyotype of 46,XX. A prenatal ultrasound at 32 weeks of gestation revealed a female fetus with a fetal biometry equivalent to 32 weeks, polyhydramnios with an increased amniotic fluid index of 26.1 cm, frontal bossing, midface hypoplasia, hypertelorism, Blake's pouch cyst with an apparent posterior fossa cyst in communication with the fourth ventricle on axial images, digital fusion, and bilateral syndactyly of the hands and feet. A DNA testing for the *FGFR2* gene was immediately performed using uncultured amniocytes obtained by repeated amniocentesis, which revealed a heterozygous c.758C>G, CCT>CGT transversion leading to a p.Pro253Arg (P253R) mutation in the *FGFR2* gene. Subsequently, a diagnosis of Apert syndrome was made. Molecular analysis of the *FGFR2* gene in the parents did not reveal such a mutation. The fetus postnatally manifested frontal bossing, midface hypoplasia, and bilateral syndactyly of the hands (mitten hands) and feet.

Conclusion: Prenatal diagnosis of polyhydramnios, frontal bossing, and midface hypoplasia associated with brain and digital abnormalities should include a differential diagnosis of Apert syndrome. A molecular analysis of *FGFR2* using uncultured amniocytes is useful for rapid confirmation of Apert syndrome at prenatal diagnosis.

Copyright © 2013, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: Apert syndrome; Blake's pouch cyst; *FGFR2*; P253R; prenatal diagnosis

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Number 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

Introduction

Apert syndrome [Online Mendelian Inheritance in Man (OMIM): #101200] is characterized by acrocephaly, craniosynostosis, midface hypoplasia, and syndactyly of the hands and feet with the bony structures exhibiting a tendency to fuse. The disorder is caused by mutations in the fibroblast growth factor receptor 2 (*FGFR2*) gene (OMIM: 176943), which is located at 10q26.13. Most cases (98%) of Apert syndrome are sporadic, and only a few are inherited in an autosomal dominant pattern [1]. The S252W (p.Ser252Trp, c.755C>G, TCG>TGG) and P253R (p.Pro253Arg, c.758C>G, CCT>CGT) mutations account for over 98% of the cases with Apert syndrome with approximately two-thirds of the cases being caused by the S252W mutation and the remaining one-third of the cases being caused by the P253R mutation [2–5]. Here, we report our experience of a rapid detection of *de novo* P253R mutation in the *FGFR2* gene using uncultured amniocytes in a pregnancy affected by polyhydramnios, Blake's pouch cyst, and Apert syndrome.

Case report

A 30-year-old, gravida 3, para 2 woman was referred for genetic counseling at 32 weeks of gestation because of polyhydramnios and craniofacial and digital abnormalities in the fetus. Her husband was 31 years old. She and her husband were both healthy and unrelated, and there was no family history of congenital malformations. She had undergone amniocentesis at 18 weeks of gestation because of maternal anxiety. Amniocentesis revealed a karyotype of 46,XX. A prenatal ultrasound at 32 weeks of gestation revealed a female fetus with a fetal biometry equivalent to 32 weeks, polyhydramnios with an increased amniotic fluid index of 26.1 cm, frontal bossing, midface hypoplasia, hypertelorism, Blake's pouch cyst with an apparent posterior fossa cyst in communication with the fourth ventricle on axial images, digital fusion, and bilateral syndactyly of the hands and feet (Fig. 1). A DNA testing for the *FGFR2* gene was immediately performed using uncultured amniocytes obtained by repeated amniocentesis, which revealed a heterozygous c.758C>G, CCT>CGT transversion leading to a p.Pro253Arg (P253R) mutation in the *FGFR2* gene (Fig. 2). Subsequently, a diagnosis of Apert syndrome was made. However, a molecular analysis of the *FGFR2* gene in the parents did not reveal such a mutation. The woman decided to discontinue the pregnancy, and a 2372-g female baby was delivered with frontal bossing, midface hypoplasia, and bilateral syndactyly of the hands (mitten hands) and feet (Fig. 3).

Discussion

The peculiar aspect of this case is the association of the P253R mutation in the *FGFR2* gene with severe syndactyly of the hands and feet, polyhydramnios, and Blake's pouch cyst. A genotype–phenotype analysis in Apert syndrome has

suggested two recurrent mutations of S252W and P253R on the severity of craniofacial abnormalities and syndactyly, respectively [6,7]. Slaney et al [6] found that syndactyly in the hands and feet were more severe in patients with the P253R mutation, whereas cleft palate was significantly more frequent in patients with the S252W mutation. von Gernet et al [7] found higher frequency of profound midface retrusion and severe malocclusion in patients with the S252W mutation and higher frequency of profound severity of syndactyly in patients with the P253R mutation.

Polyhydramnios may develop in the third trimester in pregnancies with Apert syndrome [8,9]. Chen et al [9] previously reported polyhydramnios in a pregnancy affected with the S252W mutation, in addition to upper airway obstruction and gastroesophageal reflux. The present case shows that a fetus affected by the P253R mutation and a Blake's pouch cyst may also develop polyhydramnios in the third trimester. The Blake's pouch is an inferior protrusion of the fourth ventricle into the retrocerebellar subarachnoid space resulting from a finger-like expansion of the posterior membranous area [10]. Paladini et al [11] proposed that the diagnostic criteria of a Blake's pouch cyst should include (1) normal anatomy and size of the vermis; (2) mild/moderate anticlockwise rotation of the vermis; and (3) normal size of the cisterna magna. Paladini et al [11] found that a Blake's pouch cyst in the fetus is associated with an apparently increased risk of congenital heart disease and trisomy 21.

Polyhydramnios associated with Apert syndrome has been suggested to be caused by decreased fetal swallowing related to abnormalities in the central nervous system (CNS) [12]. The present case was associated with posterior fossa anomaly. The CNS anomalies associated with Apert syndrome have been well documented [13–21]. For example, Ferreira et al [17] reported brain anomalies in cases with Apert syndrome including ventriculomegaly (48.5%), hydrocephalus (9%), gyral abnormalities, and other anomalies (e.g., agenesis or hypogenesis of the corpus callosum, posterior fossa anomalies) (21%). Quintero-Rivera et al [19] reported brain anomalies in 30 cases with Apert syndrome including nonprogressive ventriculomegaly (76%), hydrocephalus (13%), completely absent septum pellucidum (17%), partially absent septum pellucidum (23%), deficiency of septal leaflets (10%), agenesis of the corpus callosum (7%), deficiency of the corpus callosum (3%), and thinning of the corpus callosum (13%).

Using the sophisticated sonographic observations of syndactyly, midface hypoplasia, and abnormal cranial shape, prenatal sonographic diagnosis of Apert syndrome can be achieved in the second trimester [21–23]. However, as presented in this case, most prenatally detected cases of Apert syndrome were reported in the third trimester because the craniofacial and digital abnormalities and polyhydramnios associated with Apert syndrome are more obvious only in the late stages of gestation upon performing a prenatal ultrasound [24]. Recently, a noninvasive prenatal diagnosis of Apert syndrome using fetal DNA from maternal plasma by




Fig. 1. Prenatal ultrasound at 32 weeks of gestation shows (A) and (B) a Blake's pouch cyst, (C) and (D) frontal bossing and midface hypoplasia, (E) hypertelorism, (F) and (G) digital fusion and syndactyly of the hands, and (H) and (I) digital fusion and syndactyly of the foot.




Fig. 2. DNA testing shows a c.758C>G, CCT>CGT, or P253R mutation in the fetus. WT = wild type.

polymerase chain reaction-based techniques has been performed [25,26]. However, at present, rapid detection of the *FGFR2* mutation in suspected Apert syndrome cases on pre-natal ultrasound using common mutation sequencing on uncultured amniocytes is easier and less expensive than noninvasive prenatal diagnosis using high-throughput next generation DNA sequencing. Therefore, we suggest performing a molecular analysis of the *FGFR2* gene using uncultured amniocytes for rapid confirmation of Apert syndrome at pre-natal diagnosis.

Acknowledgments

This work was supported by research grants from the National Science Council (Grant Nos. NSC-99-2628-B-195-001-MY3 and NSC-101-2314-B-195-011-MY3) and the Mackay Memorial Hospital, Taipei, Taiwan (Grant No. MMH-E-102-04).



Fig. 3. The fetus at birth. (A) Craniofacial appearance, (B) the right hand, (C) the left hand, (D) the left foot, and (E) the right foot.

References

- [1] Tolarova MM, Harris JA, Ordway DE, Vargervik K. Birth prevalence, mutation rate, sex ratio, parents' age, and ethnicity in Apert syndrome. *Am J Med Genet* 1997;72:394–8.
- [2] Park WJ, Theda C, Maestri NE, Meyers GA, Fryburg JS, Dufresne C, et al. Analysis of phenotypic features and *FGFR2* mutations in Apert syndrome. *Am J Hum Genet* 1995;57:321–8.
- [3] Moloney DM, Slaney SF, Oldridge M, Wall SA, Sahlin P, Stenman G, et al. Exclusive paternal origin of new mutations in Apert syndrome. *Nat Genet* 1996;13:48–53.
- [4] Lajeunie E, Cameron R, El Ghouzi V, de Parseval N, Journeau P, Gonzales M, et al. Clinical variability in patients with Apert's syndrome. *J Neurosurg* 1999;90:443–7.
- [5] Oldridge M, Zackai EH, McDonald-McGinn DM, Iseki S, Morriss-Kay GM, Twigg SR, et al. *De novo* alu-element insertions in *FGFR2* identify a distinct pathological basis for Apert syndrome. *Am J Hum Genet* 1999;64:446–61.
- [6] Slaney SF, Oldridge M, Hurst JA, Morriss-Kay GM, Hall CM, Poole MD, et al. Differential effects of *FGFR2* mutations on syndactyly and cleft palate in Apert syndrome. *Am J Hum Genet* 1996;58:923–32.
- [7] von Gernot S, Golla A, Ehrenfels Y, Schuffenhauer S, Fairley JD. Genotype-phenotype analysis in Apert syndrome suggests opposite effects of the two recurrent mutations on syndactyly and outcome of craniofacial surgery. *Clin Genet* 2000;57:137–9.
- [8] Filkins K, Russo JF, Boehmer S, Camous M, Przylepa KA, Jiang W, et al. Prenatal ultrasonographic and molecular diagnosis of Apert syndrome. *Prenat Diagn* 1997;17:1081–4.
- [9] Chen CP, Lin SP, Su YN, Chen CY, Tsai FJ, Liu YP, et al. Apert syndrome associated with upper airway obstruction and gastroesophageal reflux inducing polyhydramnios in the third trimester. *Taiwan J Obstet Gynecol* 2010;49:230–4.
- [10] Shekdar K. Posterior fossa malformations. *Semin Ultrasound CT MR* 2011;32:228–41.
- [11] Paladini D, Quarantelli M, Pastore G, Sorrentino M, Sglavo G, Nappi C. Abnormal or delayed development of the posterior membranous area of the brain: anatomy, ultrasound diagnosis, natural history and outcome of Blake's pouch cyst in the fetus. *Ultrasound Obstet Gynecol* 2012;39:279–87.
- [12] Nyberg DA, Jeanty P, Glass I. Syndromes and multiple anomaly conditions. In: Nyberg DA, McGahan JP, Pretorius DH, Pilu G, editors. *Diagnostic imaging of fetal anomalies*. Philadelphia: Lippincott Williams & Wilkins; 2003. p. 133–220.
- [13] Cohen Jr MM, Kreiborg S. The central nervous system in the Apert syndrome. *Am J Med Genet* 1990;35:36–45.
- [14] Cinalli G, Renier D, Sebag G, Sainte-Rose C, Arnaud E, Pierre-Kahn A. Chronic tonsillar herniation in Crouzon's and Apert's syndromes: the role of premature synostosis of the lambdoid suture. *J Neurosurg* 1995;83:575–82.
- [15] Cinalli G, Sainte-Rose C, Kollar EM, Zerah M, Brunelle F, Chumas P, et al. Hydrocephalus and craniosynostosis. *J Neurosurg* 1998;88:209–14.
- [16] Renier D, Arnaud E, Cinalli G, Sebag G, Zerah M, Marchac D. Prognosis for mental function in Apert's syndrome. *J Neurosurg* 1996;85:66–72.
- [17] Ferreira JC, Carter SM, Bernstein PS, Jabs EW, Glickstein JS, Marion RW, et al. Second-trimester molecular prenatal diagnosis of sporadic Apert syndrome following suspicious ultrasound findings. *Ultrasound Obstet Gynecol* 1999;14:426–30.
- [18] Yacubian-Fernandes A, Palhares A, Giglio A, Gabarra RC, Zanini S, Portela L, et al. Apert syndrome: analysis of associated brain malformations and conformational changes determined by surgical treatment. *J Neurosurg* 2004;31:116–22.
- [19] Quintero-Rivera F, Robson CD, Reiss RE, Levine D, Benson CB, Mulliken JB, et al. Intracranial anomalies detected by imaging studies in 30 patients with Apert syndrome. *Am J Med Genet A* 2006;140:1337–8.
- [20] Quintero-Rivera F, Robson CD, Reiss RE, Levine D, Benson CB, Mulliken JB, et al. Apert syndrome: what prenatal radiographic findings should prompt its consideration? *Prenat Diagn* 2006;26:966–72.
- [21] David AL, Turnbull C, Scott R, Freeman J, Bilardo CM, van Maarle M, et al. Diagnosis of Apert syndrome in the second-trimester using 2D and 3D ultrasound. *Prenat Diagn* 2007;27:629–32.
- [22] Skidmore DL, Pai AP, Toi A, Steele L, Chitayat D. Prenatal diagnosis of Apert syndrome: report of two cases. *Prenat Diagn* 2003;23:1009–13.
- [23] Athanasiadis AP, Zafrafas M, Polychronou P, Florentin-Arar L, Papasozomenou P, Norbury G, et al. Apert Syndrome: the current role of prenatal ultrasound and genetic analysis in diagnosis and counselling. *Fetal Diagn Ther* 2008;24:495–8.
- [24] Chen CP, Su YN, Hsu CY, Ling PY, Tsai FJ, Chern SR, et al. Second-trimester molecular prenatal diagnosis of sporadic Apert syndrome following sonographic findings of mild ventriculomegaly and clenched hands mimicking trisomy 18. *Taiwan J Obstet Gynecol* 2010;49:129–32.
- [25] Raymond FL, Whittaker J, Jenkins L, Lench N, Chitty LS. Molecular prenatal diagnosis: the impact of modern technologies. *Prenat Diagn* 2010;30:674–81.
- [26] Au PK, Kwok YK, Leung KY, Tang LY, Tang MH, Lau ET. Detection of the S252W mutation in fibroblast growth factor receptor 2 (FGFR2) in fetal DNA from maternal plasma in a pregnancy affected by Apert syndrome. *Prenat Diagn* 2011;31:218–20.