

Research Letter

Prenatal diagnosis of *de novo* monosomy 7q33-qter associated with hydrops fetalis, semilobar holoprosencephaly, and premaxillary dysgenesis

Pi-Lin Sung^{a,b,c}, Eong-Eong Cheng^{a,b}, Yann-Jang Chen^d, Schu-Rern Chern^e, Chung-Yu Shih^{a,b},
Chia-Ming Chang^{a,b}, Peng-Hui Wang^{a,b}, Ming-Shyen Yen^{a,b}, Chi-Ying F. Huang^{c,f},
Chih-Ping Chen^{b,e,g,h,i,j,*}

^a Department of Obstetrics and Gynecology, Taipei Veterans General Hospital, Taipei, Taiwan

^b Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University School of Medicine, Taipei, Taiwan

^c Institute of Clinical Medicine, National Yang-Ming University, Taipei, Taiwan

^d Department of Life Sciences and Institute of Genome Sciences, National Yang-Ming University, Taipei, Taiwan

^e Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^f Institute of Biopharmaceutical Sciences, National Yang-Ming University, Taipei, Taiwan

^g Department of Biotechnology, Asia University, Taipei, Taiwan

^h School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

ⁱ Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^j Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

Accepted 4 May 2013

A 30-year-old nulligravida woman was referred to Taipei Veteran General Hospital for genetic amniocentesis at 21 weeks of gestation due to multiple abnormalities detected in routine prenatal ultrasound since 20 weeks of gestation. The woman and her 30-year-old husband were healthy, and there was no family history of congenital malformations. Down syndrome risk was less than 1/3000 at her 18-week gestation.

A detailed high-resolution ultrasound examination revealed semilobar holoprosencephaly, premaxillary dysgenesis (bilateral cleft lip/cleft palate), massive left pleural effusion, and ascites (Fig. 1A–D). Routine cytogenetics with GTG-banding (G-banding obtained by trypsin and Giemsa) documented the presence of a 46,XX,del(7)(pter → q32): *dn* (Fig. 2A) in all the 20 metaphases examined (from 13 colonies) because both parents' chromosome examinations revealed normal findings. The other parameters of the fetus were as follows: estimated body weight, 576 g (>97th percentile); biparietal diameter, 45.5 mm (<2nd percentile); femoral length, 35.3 mm (20th percentile), and abdominal circumference, 211 mm² (>98th percentile). After genetic counseling with parents, termination of pregnancy was performed at 22 weeks of gestation and a 510-g fetus with bilateral cleft lip and palate, hydrops fetalis, flat/broad nasal bridge, and upslanting palpebral fissures was

delivered (Fig. 3A–C). Further investigation of abnormality was suggested; oral informed consent was obtained from the parents, but they refused autopsy. The tissue from the fetal skin was sent for further molecular studies in order to determine the breakpoints. Genome-wide oligonucleotide-based array comparative genomic hybridization (array CGH) (SurePrint H3 Human CGH Microarray Kit 60k; Agilent Technologies, Santa Clara, CA, USA) demonstrated a 23.25-Mb deletion at 7q33 to qter (135353693–158602499) × 1 (Fig. 2B). Fluorescent in situ hybridization (FISH) studies on cultured fetal skin cells indicated the deletion of terminal long arm of chromosome 7 (Fig. 2C). The parental origin of the deletion on terminal 7q was analyzed with quantitative fluorescent polymerase chain reaction using the cultured fetal cell and the polymorphic small tandem repeat markers specific for chromosome 7q (Fig. 4). After studying with the microsatellite markers D7S1804, D7S3070, and D7S1823 at 7q32.3, 7q36.1, and 7q36.2, respectively, only D7S3070 and D7S1823 inherited from maternal allele were present. The breakpoint was close to 7q32.3, and the deletion was of paternal origin.

We have presented a case of prenatal diagnosis and molecular characterization of a *de novo* terminal deletion of 7q33-qter. To date, more than 35 cases of partial monosomy 7q32-qter have been reported; the syndrome of 7q was further delineated by Bernstein et al in 1980 [1], which usually arose *de novo*. The variable clinical manifestations of 7q terminal deletion syndrome include growth restriction, developmental

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.
E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).




Fig. 1. Prenatal ultrasound: (A) Premaxillary dysgenesis: bilateral cleft lip; (B) semilobar holoprosencephaly; (C) hydrops fetalis: ascites and pleural effusion; and (D) fetal facial structure by three-dimensional ultrasound confirmation.

retardation, microcephaly, holoprosencephaly, bilateral cleft palate and/or lip, ocular abnormalities, abnormal palm/sore crease, caudal dysgenesis, dental abnormalities, and congenital heart defect [1–10]. The present case with hydrops fetalis, including ascites, pleural effusion, and pericardial effusion, is very rare. Finley et al [7] described a female fetus at 27 weeks of gestation that had intrauterine growth restriction, microcephaly, single semilunar valve, a ventricular septal defect (VSD), a large atrial septal defect, a single outflow tract and truncus arteriosus, and ascites, but no pericardial or pleural effusion. The cytogenetic result yielded a 46,XX,del(7q)(q32-qter). Fetal ascites had resolved by 32 weeks. After delivery, the small for gestational age (SGA) infant had symmetric growth restriction and complex heart defects with type III truncus arteriosus, VSD, atrial septal defect, a right aortic arch, and a persistent left superior vena cava entering the coronary artery. The infant had episodes of bradycardia and hypoxia, and died at the age of 8 weeks. Finley et al [7] suggested that fetal ascites were due to cardiac defect. Congenital heart defect has been reported in up to 20% of 7q deletion. Tiller et al [11] reported a preterm male infant with deletion of 7q34-qter that had hypoplasia of the main pulmonary artery, absent pulmonary valve, VSD, and anomalous right pulmonary artery, and Chen et al [12] reported a prenatal case with deletion of 7q35-qter that showed microcephaly and tetralogy of Fallot characterized by an overriding aorta, a

small VSD, and a small pulmonary trunk. In the present case, no cardiac defect was detected in prenatal ultrasound that could have caused hydrops fetalis; however, cardiac defect could be assessed if autopsy was allowed by parents.

About 87% of patients with *de novo* terminal 7q deletions are associated with holoprosencephaly and 45% with caudal deficiency sequence [13–15]. The 7q terminal region is believed to contain genes that “play a critical role in differentiation of midline mesoderm at both ends of the developing notocord” [16]. Schwartz et al [9] reported a newborn infant with 46,XX,del(7)(pter → q32:) that had cebocephaly with holoprosencephaly. McMorrow et al [17] described a male infant with deletion of 7q32-qter having hydrocephalus with a prominent forehead, horizontal palpebral fissures; bilateral anophthalmos; elongated, narrow nose; a flat nasal bridge; agenesis of the nasal septum; a completely cleft palate; a small mouth; a wide nasal philtrum; low-set, deformed ears; a webbed neck; bilaterally undescended testes; and a left club foot. Chuang et al [18] reported two fetus with der(7)t(1;7)(q32;q32)pat inherited from the father with double translocations, 46,XY, t(1;7)(q32;q32), t(14;15)(q32.1;q26.3) that had alobar holoprosencephaly and cleft palate. Several genes involved in normal brain development were mapped in the region of 7q36, such as sonic hedgehog gene (*SHH*), or *HPE3* [19], *En2* [20,21], and *HTR5A* [22]. Haploinsufficiency of these genes in the present patient severely affected brain function and facial




Fig. 2. (A) G-banded chromosome 7 and del(7q33). (B) Array CGH analysis with the Human CGH Microarray Kit 8 × 60 k array showed 23.25-Mb terminal deletion at 7q33(135353693–158602499) × 1, which was indicated by a red arrow. (C) FISH metaphase from amniotic fluid cells hybridized with a 7 centromere probe (PR11-41F22; red) and loss of signal of 7q telomere probe (RP11-477A19; green) on chromosome 7.

structure, as reported previously. The gene for caudal development, *HLXB9* [16,23], was also mapped in the deleted region, but perinatal investigations on sacral vertebrae did not reveal remarkable abnormalities. Haploinsufficiency of *HLXB9* in our patient seems not to cause severe consequences of sacral development.

The breakpoint and the parental origin of the deletion on terminal 7q were analyzed with quantitative fluorescent polymerase chain reaction using the skin cells and the polymorphic small tandem repeat markers specific for

chromosome 7q (Fig. 3). With the microsatellite marker D7S1804, two alleles inherited from the parents were seen in the proband, but with the markers D7S3070 and D7S1823, only one maternal allele was detected. The deletion was of paternal origin. Numerous studies of the parental origin of syndromes associated with cytogenetically detectable deletions have been published, including the 4p deletion associated with Wolf–Hirschhorn syndrome [24], deletion of 5p associated with the cri-du-chat syndrome [25], deletions of 18q [26], and terminal deletion associated with the 9p deletion




Fig. 3. (A) Anterior view of the face: bilateral cleft lip; (B) lateral view of the face; and (C) whole-body view: hydrops fetalis.




Fig. 4. According to the electropherograms on amplified PCR products using chromosome 7 specific STRs and genomic DNAs, the proband lost paternal markers at 7q36.1 and 7q36.2, which indicated the terminal deletion was inherited from paternal side. PCR = polymerase chain reaction; STR = small tandem repeat.

syndrome [27]. All have been shown to be paternal in origin in the majority of patients studied. Thomas et al [28] summarized parental origin and, wherever possible, the chromosomal origin of 115 *de novo* unbalanced structural chromosome abnormalities detectable by light microscopy. Simple terminal deletions were approximately 70% paternal and 30% maternal in origin in the 39 cases with terminal deletion studied. For chromosome 7q, two cases were reported: Case 112 with $\text{del}(7)(\text{q}36.1)$ was of paternal origin and Case 115 with $\text{del}(7)(\text{q}36.2)$ was of maternal origin. Chen et al [12] reported $\text{del}(7)(\text{q}35)$ in a female fetus, which was of maternal origin. The terminal deletion of chromosome 7q does not appear to be derived from paternal origin predominantly.

In summary, a great variety of phenotypes can be detected in terminal 7q deletion syndrome. The present case provided

evidence that monosomy of 7q33-qter is associated with hydrops fetalis, and congenital cardiac defect should be examined carefully.

Acknowledgments

This study was supported partially by grants from Yen Tjing Ling Medical Foundation (CI-100-34) and Taipei Veterans General Hospital (V101B-036).

References

- [1] Bernstein R, Dawson B, Morcom G, Wagner J, Jenkins T. Two unrelated children with distal long arm deletion of chromosome 7: clinical features, cytogenetic and gene marker studies. *Clin Genet* 1980;17:228–37.

- [2] de Grouchy J, Turleau C. Tentative localization of a Hageman (Factor XII) locus on 7q, probably the 7q35 band. *Humangenetik* 1974;24:197–200.
- [3] Harris EL, Wappner RS, Palmer CG, Hall B, Dinno N, Seashore MR, et al. 7q deletion syndrome (7q32 leads to 7qter). *Clin Genet* 1977;12:233–8.
- [4] Kousseff BG, Hsu LY, Paciuc S, Hirschhorn K. A partial long arm deletion of chromosome 7:46, XY, del(7)(q32). *J Med Genet* 1977;14:144–7.
- [5] Biederman B, Bowen P. Balanced t(8;9)(q12;q33)pat carrier with phenotypic abnormalities attributable to a *de novo* terminal deletion of the long arm of chromosome 7. *Hum Genet* 1978;41:101–7.
- [6] Schrandt-Stumpel C, Schrandt J, Fryns JP, Hamers G. Caudal deficiency sequence in 7q terminal deletion. *Am J Med Genet* 1988;30:757–61.
- [7] Finley BE, Seguin JH, Bennett TL, Ardinger R, Burlbaw J, Levitch L, et al. Terminal deletion of 7q presenting *in utero* with a truncus arteriosus and nonimmune hydrops. *Am J Med Genet* 1993;47:221–2.
- [8] Friedrich U, Osterballe O, Stenbjerg S, Jorgensen J. A girl with karyotype 46, XX, del(7)(pter leads to q32:). *Hum Genet* 1979;51:231–5.
- [9] Schwartz S, Meekins J, Panny SR, Sun CC, Cohen MM. Brief clinical report: cebocephaly–holoprosencephaly in a newborn girl with a terminal 7q deletion [46, XX, del(7)(pter leads to q32:)]. *Am J Med Genet* 1983;15:141–4.
- [10] Young RS, Weaver DD, Kukulich MK, Heerema NA, Palmer CG, Kawira EL, et al. Terminal and interstitial deletions of the long arm of chromosome 7: a review with five new cases. *Am J Med Genet* 1984;17:437–50.
- [11] Tiller GE, Watson MS, Duncan LM, Dowton SB. Congenital heart defect in a patient with deletion of chromosome 7q. *Am J Med Genet* 1988;29:283–7.
- [12] Chen CP, Chern SR, Chang TY, Tzen CY, Lee CC, Chen WL, et al. Prenatal diagnosis of *de novo* terminal deletion of chromosome 7q. *Prenat Diagn* 2003;23:375–9.
- [13] Frinits SG, Schrandt-Stumpel CT, Schoenmakers EF, Engelen JJ, Reekers AB, Van den Neucker AM, et al. Strong variable clinical presentation in 3 patients with 7q terminal deletion. *Genet Couns* 1998;9:5–14.
- [14] Rodriguez L, Cuadrado Perez I, Herrera Montes J, Lorente Jareno ML, Lopez Grondona F, Martinez-Frias ML. Terminal deletion of the chromosome 7(q36–qter) in an infant with sacral agenesis and anterior myelomeningocele. *Am J Med Genet* 2002;110:73–7.
- [15] Su PH, Chen JY, Chen SJ, Tsao TF, Lai YJ. Sacral dysgenesis associated with terminal deletion of chromosome 7 (q36–qter). *Pediatr Neonatol* 2008;49:189–92.
- [16] Lynch SA, Bond PM, Copp AJ, Kirwan WO, Nour S, Balling R, et al. A gene for autosomal dominant sacral agenesis maps to the holoprosencephaly region at 7q36. *Nat Genet* 1995;11:93–5.
- [17] McMorrow LE, Toth IR, Gluckson MM, Leff A, Wolman SR. A lethal presentation of *de novo* deletion 7q. *J Med Genet* 1987;24:629–31.
- [18] Chuang L, Kuo PL, Yang HB, Chien CH, Chen PY, Chang CH, et al. Prenatal diagnosis of holoprosencephaly in two fetuses with der (7) t(1;7)(q32;q32)pat inherited from the father with double translocations. *Prenat Diagn* 2003;23:134–7.
- [19] Roessler E, Belloni E, Gaudenz K, Jay P, Berta P, Scherer SW, et al. Mutations in the human Sonic Hedgehog gene cause holoprosencephaly. *Nat Genet* 1996;14:357–60.
- [20] Joyner AL. Engrailed, *Wnt* and *Pax* genes regulate midbrain–hindbrain development. *Trends Genet* 1996;12:15–20.
- [21] Song DL, Chalepakakis G, Gruss P, Joyner AL. Two Pax-binding sites are required for early embryonic brain expression of an Engrailed-2 transgene. *Development* 1996;122:627–35.
- [22] Schanen NC, Scherer SW, Tsui LC, Francke U. Assignment of the 5-hydroxytryptamine (serotonin) receptor 5A gene (HTR5A) to human chromosome band 7q36.1. *Cytogenet Cell Genet* 1996;72:187–8.
- [23] Ross AJ, Ruiz-Perez V, Wang Y, Hagan DM, Scherer S, Lynch SA, et al. A homeobox gene, *HLXB9*, is the major locus for dominantly inherited sacral agenesis. *Nat Genet* 1998;20:358–61.
- [24] Quarrell OW, Snell RG, Curtis MA, Roberts SH, Harper PS, Shaw DJ. Paternal origin of the chromosomal deletion resulting in Wolf–Hirschhorn syndrome. *J Med Genet* 1991;28:256–9.
- [25] Overhauser J, McMahon J, Oberlender S, Carlin ME, Niebuhr E, Wasmuth JJ, et al. Parental origin of chromosome 5 deletions in the cri-du-chat syndrome. *Am J Med Genet* 1990;37:83–6.
- [26] Cody JD, Pierce JF, Brkanac Z, Plaetke R, Ghidoni PD, Kaye CI, et al. Preferential loss of the paternal alleles in the 18q-syndrome. *Am J Med Genet* 1997;69:280–6.
- [27] Micale MA, Haren JM, Conroy JM, Crowe CA, Schwartz S. Parental origin of *de novo* chromosome 9 deletions in del(9p) syndrome. *Am J Med Genet* 1995;57:79–81.
- [28] Thomas NS, Durkie M, Van Zyl B, Sanford R, Potts G, Youings S, et al. Parental and chromosomal origin of unbalanced *de novo* structural chromosome abnormalities in man. *Hum Genet* 2006;119:444–50.