

Case Report

VACTERL association with hydrocephalus in a fetus conceived by *in vitro* fertilization and embryo transfer

Chih-Ping Chen^{a,b,c,d,e,f,g,*}, Tung-Yao Chang^h, Yi-Yung Chen^a, Schu-Rern Chern^b, Jun-Wei Su^{a,i},
Wayseen Wang^{b,j}

^a Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^c Department of Medicine, Mackay Medical College, New Taipei City, Taiwan

^d Department of Biotechnology, Asia University, Taichung, Taiwan

^e School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^f Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^g Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^h Taiji Fetal Medicine Center, Taipei, Taiwan

ⁱ Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan

^j Department of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 4 May 2013

Abstract

Objective: We present a case of VACTERL association with hydrocephalus (VACTERL-H) in a fetus conceived by *in vitro* fertilization (IVF) and embryo transfer (ET) and review the literature.

Case report: A 35-year-old woman presented with multiple fetal anomalies at 22 weeks of gestation. She and her husband were non-consanguineous and there was no family history of congenital malformations. This was her second pregnancy conceived via IVF-ET. Two embryos had been implanted and only one survived. She underwent chorionic villus sampling at 17 weeks of gestation because of oligohydramnios and advanced maternal age. Cytogenetic analysis revealed a karyotype of 46,XY, and array comparative genomic hybridization analysis revealed no genomic imbalance. Prenatal ultrasound at 21 weeks of gestation revealed a singleton with fetal biometry equivalent to 18 weeks, ventriculomegaly, a small cerebellum, and a ventricular septal defect. Level II ultrasound showed a single umbilical artery, scoliosis, a right club hand, radial aplasia, and renal agenesis. The parents elected to terminate the pregnancy at 22 weeks of gestation, and a fetus was delivered with bilateral arthrogryposis, right radial aplasia, a club hand and thumb aplasia, hypoplasia of the left thumb, scoliosis, and an imperforate anus. The clinical findings were consistent with the diagnosis of VACTERL-H. Molecular analysis of *PTEN*, *FANCB*, and *HOXD13* genes revealed no mutation.

Conclusion: Prenatal diagnosis of radial ray defects in fetuses conceived by assisted reproductive technology should include a differential diagnosis of VACTERL association with anorectal malformation. VACTERL-H may occur in pregnancy after IVF-ET.

Copyright © 2013, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

Keywords: assisted reproductive technology; hydrocephalus; *in vitro* fertilization; prenatal diagnosis; VACTERL association

Introduction

VACTERL association (OMIM 192350) occurs in 1/10,000–1/40,000 infants and is an acronym to describe the non-random association of at least three of the following core abnormalities: vertebral defects (V), anal atresia (A), cardiac defects (C), tracheo-esophageal fistula (TE), renal anomalies

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).




Fig. 1. (A) X-Ray of the fetus at birth. (B) Radial ray defects of the right upper limb and absence of the right thumb. (C) Left thumb hypoplasia. (D) Imperforate anus.

(R), and limb defects (L) [1]. Approximately 90% of VACTERL association cases occur sporadically, whereas 10% of cases involve familial inheritance [2]. Genetic factors associated with VACTERL association include mitochondrial dysfunction, respiratory chain deficiency, chromosomal deletions or duplications, and mutations in *HOXD13*, *ZIC3*, *PTEN*, *FANCB*, and *FOXF1* genes [3–24]. Here we describe our experience of prenatal diagnosis of VACTERL association with hydrocephalus (VACTERL-H) in a fetus conceived by *in vitro* fertilization (IVF) and embryo transfer (ET) and review the literature.

Case report

A 35-year-old woman (gravida 2, para 0) presented with multiple fetal anomalies at 22 weeks of gestation. The woman's husband was 35 years old. She and her husband were non-consanguineous and there was no family history of congenital malformations. The woman did not have diabetes mellitus during this pregnancy. She had experienced one spontaneous abortion and had suffered from infertility. This was her second pregnancy conceived via IVF-ET. Two embryos had been implanted and only one survived. She underwent chorionic villus sampling at 17 weeks of gestation because of oligohydramnios and advanced maternal age. Cytogenetic analysis

revealed a karyotype of 46,XY, and an array comparative genomic hybridization analysis revealed no genomic imbalance. Prenatal ultrasound at 21 weeks of gestation revealed a singleton with fetal biometry equivalent to 18 weeks, ventriculomegaly, a small cerebellum, and a ventricular septal defect (VSD). Level II ultrasound showed a single umbilical artery, scoliosis, a right club hand, right radial aplasia, and right renal agenesis. The parents elected to terminate the pregnancy at 22 weeks of gestation, and a 342-g malformed fetus was delivered with bilateral arthrogryposis, right radial aplasia, a right club hand, aplasia of the right thumb, hypoplasia of the left thumb, scoliosis, and an imperforate anus (Fig. 1). The clinical findings were consistent with the diagnosis of VACTERL-H. Molecular analysis of *PTEN*, *FANCB*, and *HOXD13* genes revealed no mutation.

Discussion

With the advent of fetal ultrasonography, prenatal diagnosis of VACTERL association is possible in the second trimester [25–28]. The present case exhibited multiple abnormalities such as scoliosis, an imperforate anus, VSD, unilateral renal agenesis and radial ray defects, and hydrocephalus. In patients with VACTERL association, the frequency of the core abnormalities is 60–80% for vertebral anomalies, 55–90% for

imperforate anus/anal atresia, 25% for genitourinary anomalies, 40–80% for cardiac malformations, 50–80% for tracheo-esophageal fistula, 50–80% for renal anomalies, and 40–50% for limb malformations [1]. The present case had radial ray defects with radial aplasia, thumb aplasia/hypoplasia, and bilateral arthrogryposis of the hands. Radial ray defects include partial or complete absence of the radius and the first carpal and metacarpal bones and the two phalanges of the thumb, and occur in 1:30,000 live births [28,29]. Prenatal diagnosis of radial ray defects should prompt detailed fetal anatomy scans to assess growth restriction, abnormal digits, vertebral defects, anal atresia, tracheo-esophageal fistula, cardiac abnormalities, facial anomaly, thumb abnormality, and renal anomalies, including differential diagnosis of VACTERL association, trisomy 18, trisomy 13, Cornelia de Lange syndrome, Holt–Oram syndrome, Robert syndrome, Fanconi anemia, thrombocytopenia-absent-radius (TAR), acrofacial dysostosis, and Baller–Gerold syndrome [28].

The present case had VACTERL association and hydrocephalus but no history of familial inheritance, no *PTEN* or *HOXD13* mutations, no chromosomal abnormality, no genomic imbalance, and no heterotaxy. Chromosomal abnormalities associated with VACTERL association include distal 13q deletion [7], 5q11.2 deletion [16], a complex supernumerary ring chromosome 18 [18], microdeletion at 16q24.1 [15], microduplication at 22q11.21 [22], and microduplications at 1q41, 2q37.3, and 8q24.3 [24]. Cases with VACTERL-H (OMIM 276950) have been reported [30–38]. VACTERL-H has been described in association with autosomal recessive inheritance [30,31,35], X-linked VACTERL-H with *FANCB* mutations and Fanconi anemia [36,38], autosomal recessive inheritance (*FANCD1/BRCA2*) with Fanconi anemia [37], X-linked VACTERL association with or without hydrocephalus (VACTERL X; OMIM 314390), and autosomal dominant heterozygous *PTEN* mutation [6]. Reardon et al [6] reported a H61D mutation in the *PTEN* gene in a child with VACTERL association including ventriculomegaly, macrocephaly, tracheo-esophageal fistula, and bilateral radial hand anomalies. *PTEN* (OMIM 601728) is a tumor-suppressor gene and *PTEN* mutations have been associated with tumor-predisposing syndrome and PTEN hamartoma syndrome. Garcia-Barceló et al reported a heterozygous *de novo* 21-bp deletion in a polyalanine tract in the *HOXD13* gene in a 17-year-old girl with VACTERL association [14]. *HOXD13* (OMIM 142989) is a homeobox gene that plays a role in limb development. VACTERL X is associated with X-linked visceral heterotaxy-1 (OMIM 306955), which is caused by mutation in or deletion of the *ZIC3* gene. Wessels et al [19] reported VACTERL association and X-linked heterotaxy but no hydrocephalus in a male infant with polyalanine expansion in the *ZIC3* gene. Chung et al [21] reported VACTERL-H and X-linked heterotaxy in a four-generation family in which affected males had a 1.3-Mb deletion of Xq26.3 including the *ZIC3* gene. *ZIC3* (OMIM 300265) encodes a zinc-finger transcription factor that functions in the earliest stages of left–right body axis formation. Loss-of-function *FANCB* mutations result in X-linked VACTERL-H and Fanconi anemia complementation group B (OMIM

300514) in affected males [36,38]. *FANCB* (OMIM 300515) is mapped to Xp22.2 and encodes a Fanconi anemia protein. Affected males with *FANCB* mutations are predicted to manifest major phenotypic signs of ventriculomegaly, bilateral absent thumbs and radii, vertebral defects, renal agenesis, and growth retardation, and less frequent signs of brain, pituitary, eye, and ear malformations, esophageal, duodenal, and anal atresia, tracheo-esophageal fistula, lung segmentation defects, and small genitalia, in addition to Fanconi anemia. In patients with microdeletions of the FOX gene cluster (*FOXF1*, *MTHFSD*, *FOXC2*, and *FOXL1*) at 16q24.1, VACTERL association has been observed with vertebral anomalies, esophageal, duodenal and anal atresia, congenital heart defects, and urinary tract malformations, in addition to alveolar capillary dysplasia with misalignment of pulmonary veins (OMIM 265380) [15,17]. *FOXF1* (OMIM 601089) encodes forkhead-related activator 1, which is a transcription factor. *FOXF1* mutations are responsible for alveolar capillary dysplasia and multiple congenital anomalies [15].

The peculiar aspect of the present case is the association of VACTERL-H with assisted reproductive technology (ART). Sunagawa et al [39] reported concomitant occurrence of VACTERL association in dichorionic twin fetuses conceived by intracytoplasmic sperm injection (ICSI) and ET. In a German case–control study, Zwink et al [40] found a significantly increased risk of anorectal malformations after IVF [odds ratio (OR) 10.9; 95% confidence interval (CI) 6.2–19] and after ICSI (OR 7.5; 95% CI 4.6–12.2), and OR of 4.9, 11.9, and 7.9 for isolated anorectal malformations, anorectal malformations with associated anomalies, and anorectal malformations with VATER/VACTERL association, respectively. It is well known that monozygotic twinning is associated with early embryonic structural developmental defects such as sirenomelia, holoprosencephaly, anencephaly, exstrophy of the cloaca, and VACTERL association [41,42]. Recent investigations have shown increased congenital anomalies after ART [40,43–49]. We previously reported sirenomelia, limb-body wall complex, Cantrell syndrome, and anencephaly in pregnancies conceived via IVF-ET [50–52]. This case provides evidence of the occurrence of VACTERL-H in pregnancy after IVF-ET.

In conclusion, prenatal diagnosis of radial ray defects in fetuses conceived by ART should include a differential diagnosis of VACTERL association with anorectal malformation. VACTERL-H may occur in pregnancy after IVF-ET.

Acknowledgments

This work was supported by research grants NSC-99-2628-B-195-001-MY3 and NSC-101-2314-B-195-011-MY3 from the National Science Council, Taiwan and MMH-E-102-04 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Solomon BD. VACTERL/VATER association. *Orphanet J Rare Dis* 2011;6:56.

- [2] Solomon BD, Pineda-Alvarez DE, Raam MS, Cummings DA. Evidence for inheritance in patients with VACTERL association. *Hum Genet* 2010;127:731–3.
- [3] Cormier-Daire V, Rustin P, Rötting A, Chrétien D, Le Merrer M, Belli D, et al. Craniofacial anomalies and malformations in respiratory chain deficiency. *Am J Med Genet* 1996;66:457–63.
- [4] Damian MS, Seibel P, Schachenmayr W, Reichmann H, Dorndorf W. VACTERL with the mitochondrial np 3243 point mutation. *Am J Med Genet* 1996;62:398–403.
- [5] Gebbia M, Ferrero GB, Pilia G, Bassi MT, Aylsworth A, Penman-Splitt M, et al. X-linked situs abnormalities result from mutations in ZIC3. *Nat Genet* 1997;17:305–8.
- [6] Reardon W, Zhou XP, Eng C. A novel germline mutation of the PTEN gene in a patient with macrocephaly, ventricular dilatation, and features of VATER association. *J Med Genet* 2001;38:820–3.
- [7] Walsh LE, Vance GH, Weaver DD. Distal 13q Deletion syndrome and the VACTERL association: case report, literature review, and possible implications. *Am J Med Genet* 2001;98:137–44.
- [8] von Kleist-Retzow J-C, Cormier-Daire V, Viot G, Goldenberg A, Mardach B, Amiel J, et al. Antenatal manifestations of mitochondrial respiratory chain deficiency. *J Pediatr* 2003;143:208–12.
- [9] Ware SM, Peng J, Zhu L, Fernbach S, Colicos S, Casey B, et al. Identification and functional analysis of ZIC3 mutations in heterotaxy and related congenital heart defects. *Am J Hum Genet* 2004;74:93–105.
- [10] Thauvin-Robinet C, Faivre L, Huet F, Journeau P, Glorion C, Rustin P, et al. Another observation with VATER association and a complex IV respiratory chain deficiency. *Eur J Med Genet* 2006;49:71–7.
- [11] Felix JF, Tibboel D, de Klein A. Chromosomal anomalies in the aetiology of oesophageal atresia and tracheo-oesophageal fistula. *Eur J Med Genet* 2007;50:163–75.
- [12] Tüzel E, Samli H, Kuru I, Türkmen S, Demir Y, Maralcan G, et al. Association of hypospadias with hypoplastic synpolydactyly and role of HOXD13 gene mutations. *Urology* 2007;70:161–4.
- [13] Zhao X, Sun M, Zhao J, Leyva JA, Zhu H, Yang W, et al. Mutations in HOXD13 underlie syndactyly type V and a novel brachydactyly–syndactyly syndrome. *Am J Hum Genet* 2007;80:361–71.
- [14] Garcia-Barceló M-M, Wong KK, Lui VC-H, Yuan Z-W, So M-T, Ngan ES-W, et al. Identification of a HOXD13 mutation in a VACTERL patient. *Am J Med Genet* 2008;146A:3181–5.
- [15] Stankiewicz P, Sen P, Bhatt SS, Storer M, Xia Z, Bejjani BA, et al. Genomic and genic deletions of the FOX gene cluster on 16q24.1 and inactivating mutations of FOXF1 cause alveolar capillary dysplasia and other malformations. *Am J Hum Genet* 2009;84:780–91.
- [16] de Jong EM, Douben H, Eussen BH, Felix JF, Wessels MW, Poddighe PJ, et al. 5q11.2 deletion in a patient with tracheal agenesis. *Eur J Hum Genet* 2010;18:1265–8.
- [17] Shaw-Smith C. Genetic factors in esophageal atresia, tracheo-esophageal fistula and the VACTERL association: roles for FOXF1 and the 16q24.1 FOX transcription factor gene cluster, and review of the literature. *Eur J Med Genet* 2010;53:6–13.
- [18] van der Veken LT, Dieleman MM, Douben H, van de Brug JC, van de Graaf R, Hoogeboom AJ, et al. Low grade mosaic for a complex supernumerary ring chromosome 18 in an adult patient with multiple congenital anomalies. *Mol Cytogenet* 2010;3:13.
- [19] Wessels MW, Kuchinka B, Heydanus R, Smit BJ, Dooijes D, de Krijger RR, et al. Polyalanine expansion in the ZIC3 gene leading to X-linked heterotaxy with VACTERL association: a new polyalanine disorder? *J Med Genet* 2010;47:351–5.
- [20] Yu S, Shao L, Kilbride H, Zwick DL. Haploinsufficiencies of FOXF1 and FOXC2 genes associated with lethal alveolar capillary dysplasia and congenital heart disease. *Am J Med Genet* 2010;152A:1257–62.
- [21] Chung B, Shaffer LG, Keating S, Johnson J, Casey B, Chitayat D. From VACTERL-H to heterotaxy: variable expressivity of ZIC3-related disorders. *Am J Med Genet* 2011;155A:1123–8.
- [22] Schramm C, Draaken M, Bartels E, Boemers TM, Aretz S, Brockschmidt FF, et al. De novo microduplication at 22q11.21 in a patient with VACTERL association. *Eur J Med Genet* 2011;54:9–13.
- [23] Solomon BD, Patel A, Cheung SW, Pineda-Alvarez DE. VACTERL association and mitochondrial dysfunction. *Birth Defects Res A Clin Mol Teratol* 2011;91:192–4.
- [24] Hilger A, Schramm C, Pennimpede T, Wittler L, Dworschak GC, Bartels E, et al. De novo microduplications at 1q41, 2q37.3, and 8q24.3 in patients with VATER/VACTERL association. *Eur J Hum Genet* 2013. <http://dx.doi.org/10.1038/ejhg.2013.58>.
- [25] Chen C-P, Tzen C-Y, Shih S-L, Yeh L-F, Wang W. A fetus with VACTERL association, unilateral aplasia of humerus and partial hemihypoplasia. *Clin Dysmorphol* 2000;9:147–8.
- [26] Chen C-P, Shih J-C, Chang J-H, Lin Y-H, Wang W. Prenatal diagnosis of right pulmonary agenesis associated with VACTERL sequence. *Prenat Diagn* 2003;23:515–8.
- [27] Chen C-P, Shih J-C, Huang M-C, Liu Y-P, Su J-W, Chern S-R, et al. Prenatal ultrasound demonstration of scoliosis, absence of one rib, a radial club hand, congenital heart defects and absent stomach in a fetus with VACTERL association. *Taiwan J Obstet Gynecol* 2012;51:139–42.
- [28] Kennelly MM, Moran P. A clinical algorithm of prenatal diagnosis of radial ray defects with two and three dimensional ultrasound. *Prenat Diagn* 2007;27:730–7.
- [29] Sofer S, Bar-Ziv J, Abeliovich D. Radial ray aplasia and renal anomalies in a father and son. A new syndrome. *Am J Med Genet* 1983;14:151–7.
- [30] Sujansky E, Leonard B. VACTERL association with hydrocephalus – a new recessive syndrome? *Am J Hum Genet* 1983;35:119A.
- [31] Briard ML, Le Merrer M, Plauchu H, Dodinval P, Lambotte C, Moraine C, et al. Association of VACTERL and hydrocephalus: a new familial entity. *Ann Génét* 1984;27:220–3 (in French).
- [32] Evans JA, Stranc LC, Kaplan P, Hunter AGW. VACTERL with hydrocephalus: further delineation of the syndrome(s). *Am J Med Genet* 1989;34:177–82.
- [33] Iafolla AK, McConkie-Rosell A, Chen YT. VATER and hydrocephalus: distinct syndrome? *Am J Med Genet* 1991;38:46–51.
- [34] Porteous MEM, Cross I, Burn J. VACTERL with hydrocephalus: one end of the Fanconi anemia spectrum of anomalies? *Am J Med Genet* 1992;43:1032–4.
- [35] Corsello G, Giuffrè L. VACTERL with hydrocephalus: a further case with probable autosomal recessive inheritance. *Am J Med Genet* 1994;49:137–8.
- [36] Holden ST, Cox JJ, Kesterton I, Thomas NS, Carr C, Woods CG. Fanconi anaemia complementation group B presenting as X linked VACTERL with hydrocephalus syndrome. *J Med Genet* 2006;43:750–4.
- [37] Alter BP, Rosenberg PS, Brody LC. Clinical and molecular features associated with biallelic mutations in FANCD1/BRCA2. *J Med Genet* 2007;44:1–9.
- [38] McCauley J, Masand N, McGowan R, Rajagopalan S, Hunter A, Michaud JL, et al. X-linked VACTERL with hydrocephalus syndrome: further delineation of the phenotype caused by FANCB mutations. *Am J Med Genet* 2011;155A:2370–80.
- [39] Sunagawa S, Kikuchi A, Yoshida S, Miyashita S, Takagi K, Kawame H, et al. Dichorionic twin fetuses with VACTERL association. *J Obstet Gynaecol Res* 2007;33:570–3.
- [40] Zwink N, Jenetzky E, Hirsch K, Reifferscheid P, Schmiedeke E, Schmidt D, et al. Assisted reproductive techniques and risk of exstrophy–epispadias complex: a German case–control study. *J Urol* 2013;189:1524–9.
- [41] Schinzel AAGL, Smith DW, Miller JR. Monozygotic twinning and structural defects. *J Pediatr* 1979;95:921–30.
- [42] Camacho AF, Schneider A, Dorsainville D, Schutzman DL. Monozygotic twins discordant for VACTERL association. *Prenat Diagn* 2008;28:366–8.
- [43] Reefhuis J, Honein MA, Schieve LA, Correa A, Hobbs CA, Rasmussen SA. Assisted reproductive technology and major structural birth defects in the United States. *Hum Reprod* 2009;24:360–6.
- [44] Bahtiyar MO, Campbell K, Dulay AT, Kontic-Vucinic O, Weeks BP, Friedman AH, et al. Is the rate of congenital heart defects detected by fetal echocardiography among pregnancies conceived by in vitro fertilization really increased? A case–historical control study. *J Ultrasound Med* 2010;29:917–22.

- [45] Källén B, Finnström O, Lindam A, Nilsson E, Nygren KG, Otterblad PO. Congenital malformations in infants born after in vitro fertilization in Sweden. *Birth Defects Res A Clin Mol Teratol* 2010;88:137–43.
- [46] Sagot P, Bechoua S, Ferdynus C, Facy A, Flamm X, Gouyon JB, et al. Similarly increased congenital anomaly rates after intrauterine insemination and IVF technologies: a retrospective cohort study. *Hum Reprod* 2012;27:902–9.
- [47] Davies MJ, Moore VM, Willson KJ, Van Essen P, Priest K, Scott H, et al. Reproductive technologies and the risk of birth defects. *N Engl J Med* 2012;366:1803–13.
- [48] Hansen M, Kurinczuk JJ, de Klerk N, Burton P, Bower C. Assisted reproductive technology and major birth defects in Western Australia. *Obstet Gynecol* 2012;120:852–63.
- [49] Wen J, Jiang J, Ding C, Dai J, Liu Y, Xia Y, et al. Birth defects in children conceived by in vitro fertilization and intracytoplasmic sperm injection: a meta-analysis. *Fertil Steril* 2012;97:1331–7.
- [50] Chen C-P, Lee M-S, Tsai F-J, Huang M-C, Chern S-R, Wang W. Limb-body wall complex in one fetus of a dizygotic twin pregnancy obtained by egg donation, in vitro fertilization and embryo transfer: prenatal diagnosis and literature review. *Taiwan J Obstet Gynecol* 2009;48:446–50.
- [51] Chen C-P, Hsu C-Y, Lee M-S, Liu Y-P, Tsai F-J, Wu P-C, et al. Magnetic resonance imaging demonstration of sirenomelia in one fetus of a dizygotic twin pregnancy conceived by intracytoplasmic sperm injection, in vitro fertilization and embryo transfer. *Taiwan J Obstet Gynecol* 2011;50:561–3.
- [52] Chen C-P, Huang M-C, Chern S-R, Wu P-S, Su J-W, Wang W. Discordant anencephaly and Cantrell syndrome in monozygotic twins conceived by ICSI and IVF-ET. *Taiwan J Obstet Gynecol* 2013;52:297–9.