



## Review Article

## Noninvasive prenatal diagnosis

Wei-Lun Cheng<sup>a</sup>, Ching-Hua Hsiao<sup>b, c</sup>, Hua-Wei Tseng<sup>a</sup>, Tai-Ping Lee<sup>a, d, \*</sup><sup>a</sup> Department of Biotechnology and Laboratory Science in Medicine, National Yang-Ming University, Taipei, Taiwan, ROC<sup>b</sup> Department of Obstetrics and Gynecology, Taipei City Hospital, Women and Children Campus, Taipei, Taiwan, ROC<sup>c</sup> Institute of Biomedical Engineering, National Yang-Ming University, Taipei, Taiwan, ROC<sup>d</sup> Department of Otorhinolaryngology, National Yang-Ming University, Taipei, Taiwan, ROC

## ARTICLE INFO

## Article history:

Accepted 27 May 2015

## Keywords:

prenatal examination  
noninvasive prenatal diagnosis  
microfluidic system

## ABSTRACT

Prenatal examination plays an important role in present medical diagnosis. It provides information on fetal health status as well as the diagnosis of fetal treatment feasibility. The diagnosis can provide peace of mind for the perspective mother. Timely pregnancy termination diagnosis can also be determined if required. Amniocentesis and chorionic villus sampling are two widely used invasive prenatal diagnostic procedures. To obtain complete fetal genetic information and avoid endangering the fetus, noninvasive prenatal diagnosis has become the vital goal of prenatal diagnosis. However, the development of a high-efficiency separation technology is required to obtain the scarce fetal cells from maternal circulation. In recent years, the rapid development of microfluidic systems has provided an effective method for fetal cell separation. Advantages such as rapid analysis of small samples, low cost, and various designs, greatly enhance the efficiency and convenience of using microfluidic systems for cell separation. In addition, microfluidic disks can be fully automated for high throughput of rare cell selection from blood samples. Therefore, the development of microfluidic applications in noninvasive prenatal diagnosis is unlimited. Copyright © 2015, Taiwan Association of Obstetrics & Gynecology. Published by Elsevier Taiwan LLC. All rights reserved.

## Introduction

Prenatal diagnosis is an important medical technology for nearly two centuries. The procedure can notify parents of hereditary diseases in the fetus such as Down syndrome, sickle cell anemia, Edwards's syndrome, cystic fibrosis, and Duchenne muscular dystrophy. These diseases may cause neonate stunted growth, intellectual disability, physical disability, and death. To date, expecting parents can select various methods to confirm fetal health. However, different diagnosis methods may be presented with different levels of risks, where invasive prenatal diagnosis procedures may induce miscarriage risks. Therefore, the development of safe and highly valuable prenatal diagnostic techniques is greatly sought after for scientists around the world.

The first prenatal diagnosis can be traced back to as early as the 20<sup>th</sup> century, after Wilhelm Röntgen's [1] discovery of X-rays. Although X-rays can be utilized to observe fetus appearance, it

provides no genetic diagnosis. In 1966, Steele and Breg [2] separated fetal chromosomes from amniotic fluid for chromosome analysis which has laid the foundation of amniocentesis in modern medicine. Later, an Italian biologist, Simoni et al [3], performed the first trimester chorionic villus sampling (CVS) and risk assessment to validate this method as a reliable prenatal diagnosis tool [3,4]. These two methods are regarded as a model for prenatal diagnosis; however, their invasiveness may lead to a risk of miscarriage. Due to the potential risks of invasive prenatal genetic diagnosis, different noninvasive prenatal diagnosis (NIPD) techniques are actively being developed.

## Invasive prenatal diagnosis

Although invasive prenatal diagnosis is dangerous, there exists no effective replacement among today's technologies. Table 1 indicates the most direct methods of siphoning fetal samples from the mother and their associated risk of miscarriage [5–9].

## Amniocentesis

Amniocentesis was used to treat pregnant women with polyhydramnios in 1880. In 1966, American physicians Steele and Breg

\* Corresponding author. Department of Otorhinolaryngology and Biotechnology and Laboratory Science in Medicine, National Yang-Ming University, Number 155, Section 2, Lie-Nong Street, Taipei 112, Taiwan, ROC.

E-mail addresses: [long0way@msn.com](mailto:long0way@msn.com), [tplee@ym.edu.tw](mailto:tplee@ym.edu.tw) (T.-P. Lee).

**Table 1**  
Comparison of invasive prenatal diagnosis.

	Execution time (wk)	Sampling location	Diagnosis	Risk of miscarriage (%)	Refs
Chorionic villous sampling	10–12	Chorionic villus	Chromosomal abnormalities	0.5–2	[5]
Amniocentesis	14–16	Amniotic sac	Chromosomal abnormalities/neural tube defects	0.06–1.3	[6,7]
Fetal blood sampling	≥17	Fetal umbilical cord	Chromosomal abnormalities/metabolic disorders/fetal infections	2–3	[8,9]

FISH = fluorescence in situ hybridization; PCR = polymerase chain reaction.

[2] successfully cultured amniotic fluid cells for chromosome analysis. Presently, it is the most widely used invasive prenatal diagnosis because of its high accuracy and low risk characteristics. Previous research has suggested that spontaneous abortion percentage after amniocentesis is 1.7% [10], which was slightly higher than the control group that underwent only ultrasonic examinations. The report in 2000 also suggested that the fetus miscarriage risk of amniocentesis is ~0.6–0.68% [11]. This method is applicable for women at 16–18 weeks of pregnancy, and ~200–300 mL of amniotic fluid in the uterus of pregnant women is required. Amniotic fluid contains  $\sim 2-3 \times 10^5$  cells per 10 mL, and these cells are produced by fetal movements in the amniotic sac due to swallowing, urination, and physical movements [12]. Amniocentesis is useful for the diagnoses of many single-gene diseases and congenital defects such as Down syndrome, thalassemia, Adrenoleukodystrophy, and Huntington's Disease.

#### Chorionic villous sampling

The advantage of CVS is that it is suitable for early screening for women at around 10–13 weeks of pregnancy and in special cases it can be performed as early as 8 weeks [13]. This approach extracts tissue and fetal placental chorionic cells transcervically or transabdominally, where the transabdominal method has been confirmed to be safer than the transcervical method in a previous report [14]. The miscarriage risk of CVS is higher than amniocentesis, which is ~0.5–2%. The fetal diseases that can be identified by CVS are similar to those of the amniocentesis, which include chromosomal abnormalities and genetic defects [5].

#### Percutaneous umbilical cord blood sampling

Cordocentesis, also known as percutaneous umbilical cord blood sampling (PUBS), draws blood samples directly from the fetal umbilical vein with a sampling needle. This method is suitable for women at the second trimester pregnancy stage, generally performed after 17 weeks of pregnancy [15]. The late sampling time of PUB than amniocentesis and CVS is because the early fetal umbilical vein is fragile and not suitable for puncture. Studies that investigated umbilical cord puncture procedure and fetal damage found

no obvious pathological symptoms a week after PUBS [16]. Such direct sampling method has certain risks associated with gestational age, operating procedures, and sampling frequency of needle piercing, with different outcomes [8]. A statistical analysis has indicated that PUBS causes ~2–3% of miscarriages [9]. However, PUBS can be used to diagnose fetal chromosomal abnormalities, infections, and metabolic disorders that cannot be determined with CVS or amniocentesis.

#### NIPT

Currently, many scientists have been developing NIPT techniques for collecting fetal samples in order to reduce the risk of miscarriage from invasive procedures. Table 2 summarizes each technique and its disadvantages [17–28].

#### Ultrasonography

Ultrasonography is generally considered as a safe method of image rendering. It is one of the most common NIPT technologies, where images are formed when the echoes of ultrasound that penetrated the uterus tissue is received. The sound waves are reflected to the receiving probe with various interfaces, and after concussion, the signals are transformed to electrical signals to render two-dimensional, three-dimensional, or higher images. Ultrasound is often used for examining fetal congenital disability or abnormal development, by observing fetus nuchal translucency and nose bone development, which can predict the possibility of a fetus suffering from Down syndrome. However, in clinical practice, Down syndrome diagnosed with ultrasonography has a high false positives rate of ~5% [17]. Although ultrasonography examination cannot detect most of the inherited diseases or genetic defects, it is still an indispensable NIPT tool.

#### First and second trimester screening

Second trimester maternal serum screening includes double, triple, and quadruple tests. It is one of the most commonly chosen examinations. This method determines specific protein concentrations in maternal serum to calculate the likelihood and risk

**Table 2**  
Comparison of noninvasive prenatal testing.

	Execution time (wk)	Target	Diagnosis	Detection rate for Down's syndrome (%)	Disadvantage	Ref.
Ultrasonography	<20	(1) Nuchal translucency (2) Nose bone	Down's syndrome Deformity	80–90	High false positive (4.5–6.0%)	[17–19]
Triple test	16	(1) $\alpha$ -fetoprotein (2) Oestriol (3) hCG	Down's syndrome Neural tube defects	60–70	No definitive diagnosis	[20,21]
Cell-free fetal DNA (mRNA) in maternal blood	>12	(1) Fetal mRNA (2) Fetal DNA	Single-gene disorders Aneuploidy	>99	Large sample Cost effectiveness	[22–25]
Fetal cell in maternal blood	4–14	(1) Fetal lymphocytes (2) Trophoblasts (3) Nucleated red blood cells	The disease can be detected by PCR or FISH	75	Very rare	[26–28]

FISH = fluorescence in situ hybridization; PCR = polymerase chain reaction.

assessment of trisomy 21. Maternal serum screening utilizes indicators such as alpha-fetoprotein, oestriol, and human chorionic gonadotropin [29,30]. The execution time is during the second trimester of pregnancy, which is equivalent to ~15–20 weeks. The attractive advantage of maternal serum screening is that the serum is taken directly from the mother, with no invasion into the uterus. However, previous studies have indicated that this method of testing for Down syndrome is associated with a 5% false positive rate and the sensitivity is ~57–60% [31,32]. The current most effective noninvasive screening for trisomy 21 is a combination of maternal age, measurement of fetal nuchal translucency, maternal serum free  $\beta$ -human chorionic gonadotropin, and pregnancy-associated plasma protein-A concentration at 11<sup>+0</sup> weeks to 13<sup>+6</sup> weeks of gestation. This combinatory method provided around 90% detection rate of trisomy 21 with the false-positive rate of 5% [33]. In different strategies, all of the high risk assessments are recommended to be performed in combination with amniocentesis or CVS for definitive confirmation.

#### *Cell-free fetal DNA/mRNA in maternal blood*

In 1996, two teams observed cell-free DNA in patient serums for small-cell lung cancer and head and neck cancer [34,35]. The identification of cell-free DNA in serum has prompted the development of other related diagnostic technologies. Lo et al [36], was successful in identifying a male fetus cell-free DNA from the mother's serum and plasma. Three years later, fetus cell-free mRNA was also found from the mother's plasma [37].

A previous study has used real-time polymerase chain reaction (PCR) to quantify the cell-free DNA found in maternal plasma, and observed 3–6% of Y-chromosomal sequences [38]. This result suggests that maternal plasma contains a certain amount of fetal cell-free DNA for prenatal analysis. Generally cell-free DNA is commonly used for sex determination, fetal Rhesus D blood group identification, and the diagnosis of single-gene disorders such as achondroplasia and  $\beta$ -thalassemia [39–42]. However, due to the fragmented nature of cell-free DNA, the identification of fetus DNA is often difficult, where the distinctions are often made by fetal DNA markers such as male fetus Y chromosomal markers, single-nucleotide polymorphisms, and unique fetal DNA methylation markers [43]. Because of fragment information, cell-free DNA analysis has limitations for identifying fetal aneuploidy diseases such as Down syndrome, Edward's syndrome, and Patau syndrome [38,44]. Next generation sequencing can sequence a large amount of DNA sequences in a short time with upgraded computing speed and decoding algorithms. With massive parallel sequencing and powerful bioinformatics processing, maternal fetal cell-free DNA diagnosis of fetal aneuploidy has been demonstrated recently [45], with a 99% detection rate of Down syndrome and false positive rates of <1% [46–49]. In August 2011, a technology called Noninvasive Fetal Trisomy NIPD of Down syndrome was made available in a private prenatal diagnosis center in Hong Kong [45]. This method uses GC-correlation methodology with massively parallel sequencing for large-scale sequencing of maternal serum cell-free DNA [50]. The DNA fragments were obtained through computer analysis of sequencing results, and the analysis of aneuploidy was performed by comparing the DNA fragments to those found with normal levels in pregnant women in the control sample pool [23]. However, the acceptance of such exorbitant novel technology from the general public is very low.

#### *Fetal cell in maternal blood*

The first observation of fetal cells in maternal blood circulation was made by Christian Georg Schmorl, a German pathologist, in

1893. He had found multi-nucleated syncytial trophoblasts in lung tissue of a pregnant woman that died from eclampsia [51]. This discovery indicated that fetal cells are drawn into the mother's body through some mechanisms and thereby enabling NIPD by separation of fetal cells from maternal blood. Compared with the maternal fetal free DNA/mRNA, maternal circulating fetal cells have complete genetic information in their nucleus or cytoplasm that can be useful for advanced complex genetic diagnosis. This is especially useful when identifying fetal genetic defects that the mother or father also possess, since the fetal cell is not easily mistaken as a parent native cell. However, the frequency of fetal cells is very low. Therefore, effective isolation techniques for fetal cells have been the subject of constant research and development. The types of fetal cells in maternal blood can be broadly classified as trophoblasts, fetal lymphocytes, fetal nucleated red blood cells (fnRBCs), and hematopoietic stem cells (HMCs) [52]. Due to the different characteristics of each type of cell, these cells may have a different difficulty of separation and can affect the accuracy of the analyses.

#### *Trophoblasts*

Trophoblasts have been isolated as a target in earlier cell fetus research with specific antitrophoblast antibody H315 [53–55]. However, a later report has reported that this antibody is not sufficient to separate trophoblasts for prenatal diagnosis [56]. Furthermore, trophoblasts should not appear in normal pregnant women. Some reports have indicated that ~1% of placental cells with placental mosaicism can lead to discrepancies in the true state of cell nucleus type [57,58], which can greatly affect the accuracy of trophoblast based diagnosis. Therefore, NIPD based on trophoblasts is limited.

#### *Fetal lymphocytes*

Previous studies have found cells with XY chromosomes in pregnant women from conceived male fetuses [59,60]. These cells are fetal lymphocytes that have crossed the placenta into maternal circulation. Some groups have used fetal lymphocytes for fetal sex prediction [59,61], and have indicated by microscopic observation that one in every 1000–5000 mother lymphocytes is a Y-chromosome lymphocyte [62]. In addition, fetal lymphocytes can be amplified with *in vitro* culture [43], where the rare cell population can be amplified for additional analysis. The fatal flaw of using fetal lymphocytes for NIPT is that fetal cells can retain in the mother for 27 years [63]. Therefore, the test results are not reliable for second pregnancies.

#### *HMCs*

Fetal cells are very rare; therefore many researchers have increased the number of target cells to include other potentially useful fetal materials. In recent years, some groups included CD34<sup>+</sup> fetal stem cells to the target list [64]. CD34<sup>+</sup> HMCs have the ability for *in vitro* culture like fetal lymphocytes, however, maternal and fetal CD34<sup>+</sup> HMCs are not easily distinguishable. The HMC culture, while easily enlarged, the separation of fetal cell from maternal cell contamination is an important issue. It is also worth noting that residual CD34<sup>+</sup> HMCs are also found in women after pregnancy [52].

#### *FnRBCs*

The NIPD suitable target cells should have several properties such as a short half-life, appears in the early stages of pregnancy, and have specific cell markers. The fnRBCs are a good candidate with all three properties [52], which are also the most studied and isolated fetal cells from maternal blood [64–71]. The biggest drawback of this type of cell is that they are very rare and some of

the nucleated red blood cell (RBC) is maternal [72]. Previous reports have pointed out that each milliliter of maternal blood contain one to two fnRBCs [67,68], while another group has also reported hundreds of fnRBCs/mm of maternal blood [65]. The fnRBCs are approximately between 9  $\mu$ m and 13  $\mu$ m in diameter, which is bigger than adult RBCs (6–8  $\mu$ m) [73] and more similar to white blood cells (8–15  $\mu$ m). The density of fnRBCs ranges between 1.077 g/mL and 1.130 g/mL [74], similar to those of the adult RBCs (1.090–1.110 g/mL). Nucleated RBCs are RBCs that have not denucleated or are mid-erythroblasts in erythropoiesis, where the erythroblast discharges the cell nuclear and becomes a reticulocyte before entering into circulation. The discharged cell nucleus is called a pyrenocyte and are scavenged by microphages [75].

The inter- and extracellular cell markers of fnRBCs change following the development process. For example, burst-forming unit erythroid during early hematopoiesis express CD71 in bone marrow, but CD71 disappears when cells differentiate into mature RBCs. Hemoglobin appears later, gradually formed in the late erythroblast. CD45 has not been observed in full development process [76].

The biggest difference between fnRBCs and adult RBCs is the hemoglobin structure. Generally, a globin chain constructs of hemoglobin tetramers in RBCs and nRBCs changes at different stages of fetal development. The currently known fetal hemoglobin contains  $\zeta_2\epsilon_2$ ,  $\zeta_2\gamma_2$ ,  $\alpha_2\epsilon_2$ , and  $\alpha_2\gamma_2$  [73], unlike adult RBCs tetramers of  $\alpha_2\beta_2$ . The hemoglobin in fetal hematopoiesis is also called embryonic hemoglobin, and  $\zeta$  and  $\epsilon$  globin chain will be replaced by adult hemoglobin at ~9–18 weeks of gestation [77,78]. Hematopoiesis remains after the fetus's birth, and is transferred to the bone marrow. The  $\gamma$  globin chain will not completely disappear, but the vast majority of hemoglobin tetramers will be of  $\alpha_2\beta_2$  [77,78]. Therefore,  $\epsilon$  and  $\gamma$  globin chain can be used to distinguish between maternal nRBCs and fnRBCs in maternal blood; however,  $\gamma$  globin chain may appear in general adult patients with beta-thalassemia [79], and  $\epsilon$  globin chain disappears in early pregnancy. These specific markers are used by many researchers for isolating fetal cells [79–81].

### FnRBCs separation from maternal blood

There are many methods such as a magnetic-activated cell sorting (MACS), charge flow separation, fluorescence-activated cell sorting (FACS), density gradient centrifugation (DGC), and immunomagnetic beads for separating fetal cells from maternal blood. The purity and recovery rate are different for each of the methods. For most experiments, a combination of different separation

technology is often used to ensure sample purity. However, the increase in separation concatenation may decrease the recovery rate. However, poor separation methods will lead to low sample purity. Therefore, the design for optimal separation for the experiments will be the most important part of cell separation. Table 3 summarizes the performance of each isolation technique of fetal cells [26,64,65,68–70,82–84].

In fnRBCs separation, most research groups separate peripheral blood mononuclear cell (PBMC) from maternal blood, and remove RBCs using DGC [64,65,67,69,85]. The DGC separation buffer contains Ficoll, Percoll, and Histopaque (Sigma). Histopaque is Ficoll, with the addition of sodium diatrizoate products that enhance stratification in blood samples. The DGC separation buffers are divided into different densities, some of the commonly densities are 1.077 g/mL, 1.083 g/mL, and 1.119 g/mL. Previous studies have isolated fnRBCs from Ficoll DGC separated RBCs [64,67]. Another study has also determined that buffers with a higher density (1.119 g/mL) are better for isolating RBCs than low density buffers (1.090 g/mL) [68]. In addition, cell loss in Ficoll DGC and Percoll DGC with  $1.7 \times 10^6$  CD45-mononuclear cells, observed after implementation of Ficoll DGC or Percoll DGC, the ratio of loss was 14.6% and 41.2%, respectively [86]. These results indicate that different DGC separation buffers in different cell types will have different recovery rates.

After extracting PBMC from maternal blood using DGC, the fnRBCs groups are further separated with fetal specific cell surface markers. The most common screening methods are MACS, FACS, and immunomagnetic beads. The removal the CD45<sup>+</sup> cells prior to separation can effectively enhance the results of all purification techniques [87]. The CD45<sup>+</sup> cell population can be used as fnRBCs specific cell marker for positive selection. CD71 is the most common fnRBCs surface marker [64,88,89] that is also known as transferrin receptor or p90. It is the first cell marker to be used to separate fnRBCs [90]. There are also some groups that have chosen glycophorin A as the main cell marker for separation [69]. However, glycophorin A may appear on mature RBCs [76], so unless the previous step has the ability to remove the RBC cleanly, otherwise it will seriously affect the purity of the result. The fnRBCs can also be selected by i-antigen or  $\gamma$ -hemoglobin [64,68]. The i-antigen is originally expressed on the fnRBCs, which is gradually replaced by I-antigen after birth [91]. However, adult RBCs will still express a small amount of i-antigen [92], which may interfere with the separation results. In terms of specificity, the fetal specific hemoglobin is the best choice for separation. However, hemoglobin is an intracellular cell marker where immunofluorescence and cell permeability and fixation will lead to cell loss. In addition, using

**Table 3**  
Isolation of fetal cells.

Group	Year	Target cells	Method	Efficiency	Refs
Ponnusamy et al	2008	FNRCs	Percoll 1118 DGC/MACS Depletion CD45/GPA positive selection/selection lysis	3.0 cells/20mL blood (confirmed with $\epsilon$ -Hb)	[69]
Samura et al	2000	NRBCs	Histopaque 1119 or 1090 DGC/MACS Depletion CD45/flow sorting $\gamma$ -Hb <sup>+</sup>	10.67–21.91 cells/17 mL blood	[68]
Ikeya et al	2005	NRBCs	Histopaque 1095/Lectin method	89.4 $\pm$ 92.6 cells/10mL blood (confirmed with MGG and HbF)	[82]
Prieto et al	2001	Fetal NRBCs	Double DGC 1.077 & 1.107/MACS selection	194.6 cells/8.5mL blood	[70]
D'Souza et al	2009	Fetal NRBCs	Percoll DGC/FACS (CD71 <sup>+</sup> or GPA <sup>+</sup> , CD45 <sup>-</sup> , HbF <sup>+</sup> )	31–499 cells/7mL bloods	[65]
Calabrese et al	2012	NRBCs	Ficoll 1083 DGC/Dual-probe FISH (CD71 <sup>+</sup> /i-antigen <sup>+</sup> or CD34 <sup>+</sup> /i-antigen <sup>+</sup> )	71000 cells/25mL blood	[64]
Pongsritasana et al	2006	FNRCs	Ficoll DGC/FACS or MACS (CD45 <sup>+</sup> /CD71 <sup>+</sup> /CD235A <sup>+</sup> )	1.7–2.2 $\times 10^{-6}$ in MNCs	[26]
Jeon et al	2010	Fetal NRBCs	Double Percoll DGC 1.077 & 1.119/MACS CD71 selection	9.85–14.88 cells/10mL blood	[83]
Bischoff et al	2003	Progenitor cells	RosetteSep progenitor enrichment cocktail/CD45 depletion	6200 cells/20–30mL blood	[84]

DGC = density gradient centrifugation; FNRC = fetal nucleated red blood cells; MACS = magnetic activated cell sorting; NRBC = nucleated red blood cells.



immunomagnetic beads to hybridize against intracellular markers must consider if the beads are taken into the cells or conjugated through the primary antibody. Therefore, using intracellular cell marker for fnRBCs separation will encounter more difficulties.

Most of the group's fnRBCs separation countermeasures were to first remove most of the RBCs by DGC. The PBMC collected from interphase after DGC and CD45<sup>+</sup> cell separation were then subjected to MACS, FACS, immunomagnetic beads, or other selection technologies. However, novel separation procedures are still being developed. For example, the direct use of the lectin method after DGC crawl fnRBCs or size-base separation by using microfluidic chips [67,71,82]. Although the present method for fnRBCs separation has many options, the investigation of an optimum separation process is being investigated by scientists around the world.

### The isolation techniques of fnRBCs using microfluidic

With the constant development of micro- and nano-technology, novel process technology can be combined with biological materials into a tiny chip. This portable biochip, also called Lab-on-a-chip or micro total analysis systems, are widely used in cell separation technologies such as microfluidic, size-dependent hydrodynamic filtration, magnetic, and electrophoresis [93]. Microfluidic has been proven to effectively combine a variety of analytical systems that operate with a small amount of the sample for ease of sampling [94]. Microfluidic or microfluidic disk, with respect to flow cytometry, express higher sensitivity for rare cell separation [95]. The advantage of using microfluidic for cell separation include specific chip flow channel design for a single cell, automated experimental processes, high throughput, and reduced use of reagents and generated chemical wastes [96].

Microfluidic can separate fetal cells by physical properties such as size, deformability, electric, and optical properties. Antibody screening with specific cell markers can also be used with the system [96]. A previous study has used DGC followed by size-specific microfluidic chip and cell deformation to separate fnRBCs, in which it showed an average separation of 1.2 fetal cells/mm of maternal blood [67]. Another group has also used DGC, size-based microfluidic system, followed by magnetic beads for the selection of hemoglobin, which collected 274.38 nRBCs/mm of maternal blood [71]. These results indicated that the design of the microfluidic system can greatly affect the efficiency and recovery rate of separation. Currently, most practices use DGC followed by traditional methods to remove RBCs before microfluidic chip cell separation. However, DGC functional microfluidic disk has gained much attention in recent years. The microfluidic disk not only improved process automation, it was also demonstrated to separate rare cells from blood [95,97]. Despite this, there exists few reports that have utilized microfluidic disks for fetal cell separation from maternal blood. The further development of such devices still shows much promise.

### Operation after fnRBCs separation

The use of fnRBCs, resulting from different separation techniques is an important topic. Known fetus common hereditary diseases such as aneuploidy, chromosome defects, and genetic defects are diagnosed by the appearance of chromosomes, gene sequencing, PCR, multiplex ligation-dependent probe amplification, and fluorescence *in situ* hybridization (FISH). However, the resulting fetal cell isolation is often contaminated by maternal cells (background interference). Therefore, the sensitivity of the methods for the subsequent genetic analysis is very important. A previous study has demonstrated the use of highly sensitive FISH for the observation of triploid chromosome 21, 13, and 18, where

the sensitivity is up to 0.01% and the accuracy is >98% [85]. Techniques such as PCR and other methods may be more difficult than FISH for the observation of hereditary diseases under large background interference. Even after the amplification of specific gene fragments, the fetus signals are still easily obscured by the signals from maternal background cells. Therefore, samples with low purity are more suitable to use FISH for the direct calibration and observation of specific cells, which can avoid large interferences from the maternal background cells.

If diagnostics by PCR is required, with sequencing or other analogous molecular biology techniques, it is necessary to increase sample purity. Antibody and immunomagnetic based separation have a certain degree of limitation for improving purity, because with the use of antibodies it is difficult to avoid nonspecific calibration and adhesion. Therefore, micromanipulation may be the most effective way for improving sample purity [52]. Micromanipulation relies on the operator to identify the cell types, and a single cell is isolated from a large background of cells. Many molecular biology techniques are sufficient to support the operation of a single cell. For example, single cell PCR requires only one of the target cells for gene amplification. Capillary electrophoresis and short tandem repeat fragment analysis can then be used to distinguish the single cell from mother or fetus [64]. Many studies have also reported that mothers with abnormal fetuses have increased amount of maternal blood nRBC [24,98]. The number of normal maternal nRBCs is between one to 12 cells, however, a mother with a fetus suffering from beta-thalassemia will have increased nRBCs of about 22–158 cells; and a mother with abnormal ultrasonography fetal diagnosis also exhibits more nRBCs [98]. The reasons for this phenomenon is not yet clear, and perhaps, in the future, can be used as a noninvasive diagnostic indicator for the risk of fetal chromosomal abnormalities.

### Conclusion and perspectives

Using NIPD for prenatal diagnosis of aneuploidy and genetic disease, where risk-free collection of fetal samples has become the focus of recent prenatal diagnosis developments. Although cell free DNA in maternal blood has been widely used and accepted in clinical examinations, it relies heavily on new sequencing technologies that are relatively expensive. In addition, cell free DNA for the diagnosis of aneuploid and maternally inherited diseases is difficult. Therefore, fetal cells in maternal blood are an ideal solution for this limitation. With the progressive advancement in separation technology, microfluidic cell separation has become more simple and common. The microfluidic system is easily operated through automation, and requires only a trace amount of sample, which in term saves human resources and reduces costs. Furthermore, the subsequent processing of fetal cells is not limited to large fragment gene sequencing; it is able to support various molecular biology techniques for the ease of analysis.

### Conflicts of interest

The authors have no conflicts of interest relevant to this article.

### References

- [1] Röntgen WC. On a new kind of rays. *Science* 1896;3:227–31.
- [2] Steele MW, Breg Jr WR. Chromosome analysis of human amniotic-fluid cells. *Lancet* 1966;1:383–5.
- [3] Simoni G, Tibiletti MG, Dalpra L, Ferrari M, Brambati B. Routine chromosome analysis on fetal blood microaliquots obtained at fetoscopy. *Prenat Diagn* 1983;3:203–8.
- [4] Kuliev AM, Modell B, Jackson L, Simpson JL, Brambati B, Rhoads G, et al. Risk evaluation of CVS. *Prenatal Diagn* 1993;13:197–209.

- [5] Tabor A, Vestergaard CH, Lidegaard O. Fetal loss rate after chorionic villus sampling and amniocentesis: an 11-year national registry study. *Ultrasound Obstet Gynecol* 2009;34:19–24.
- [6] Eddleman KA, Malone FD, Sullivan L, Dukes K, Berkowitz RL, Kharbutli Y, et al. Pregnancy loss rates after midtrimester amniocentesis. *Obstet Gynecol* 2006;108:1067–72.
- [7] Odibo AO, Gray DL, Dicke JM, Stamilio DM, Macones GA, Crane JP. Revisiting the fetal loss rate after second-trimester genetic amniocentesis: a single center's 16-year experience. *Obstet Gynecol* 2008;111:589–95.
- [8] Buscaglia M, Ghisoni L, Bellotti M, Ferrazzi E, Levi-Setti P, Marconi AM, et al. Percutaneous umbilical blood sampling: indication changes and procedure loss rate in a nine years' experience. *Fetal Diagn Ther* 1996;11:106–13.
- [9] Antsaklis A, Daskalakis G, Papantoniou N, Michalas S. Fetal blood sampling—indication-related losses. *Prenatal Diagn* 1998;18:934–40.
- [10] Tabor A, Philip J, Madsen M, Bang J, Obel EB, Norgaard-Pedersen B. Randomised controlled trial of genetic amniocentesis in 4606 low-risk women. *Lancet* 1986;1:1287–93.
- [11] Wilson RD, Langlois S, Johnson JA, Society of Obstetricians and Gynaecologists of Canada. Mid-trimester amniocentesis fetal loss rate. *J Obstet Gynaecol Can* 2007;29:586–95.
- [12] Weise W, Gabriel D, Tanner B. Growth behavior of amnion cell cultures. *Zentralbl Allg Pathol* 1984;129:499–505.
- [13] Wapner RJ, Evans MI, Davis G, Weinblatt V, Moyer S, Krivchenia EL, et al. Procedural risks versus theology: chorionic villus sampling for Orthodox Jews at less than 8 weeks' gestation. *Am J Obstet Gynecol* 2002;186:1133–6.
- [14] Jackson LG, Zachary JM, Fowler SE, Desnick RJ, Golbus MS, Ledbetter DH, et al. A randomized comparison of transcervical and transabdominal chorionic-villus sampling. The U.S. National Institute of Child Health and Human Development Chorionic-Villus Sampling and Amniocentesis Study Group. *New Engl J Med* 1992;327:594–8.
- [15] Vantanasiri C, Kanokpongakdi S, Manassakorn J, Thitadilok W, Swasdimongkol S, Tontisirin P, et al. Percutaneous umbilical cord blood sampling. *J Med Assoc Thai* 1989;72:541–4.
- [16] Jauniaux E, Donner C, Simon P, Vanesse M, Hustin J, Rodesch F. Pathologic aspects of the umbilical cord after percutaneous umbilical blood sampling. *Obstet Gynecol* 1989;73:215–8.
- [17] Muller F, Benattar C, Audibert F, Roussel N, Dreux S, Cuckle H. First-trimester screening for Down syndrome in France combining fetal nuchal translucency measurement and biochemical markers. *Prenatal Diagn* 2003;23:833–6.
- [18] Snijders RJ, Noble P, Sebire N, Souka A, Nicolaides KH. UK multicentre project on assessment of risk of trisomy 21 by maternal age and fetal nuchal-translucency thickness at 10–14 weeks of gestation. Fetal Medicine Foundation First Trimester Screening Group. *Lancet* 1998;352:343–6.
- [19] Benn PA, Kaminsky LM, Ying J, Borgida AF, Egan JF. Combined second-trimester biochemical and ultrasound screening for Down syndrome. *Obstet Gynecol* 2002;100:1168–76.
- [20] Reynolds T. The triple test as a screening technique for Down syndrome: reliability and relevance. *Int J Womens Health* 2010;2:83–8.
- [21] Debiève F, Bouckaert A, Hubinont C, Thomas K. Multiple screening for fetal Down's syndrome with the classic triple test, dimeric inhibin A and ultrasound. *Gynecol Obstet Invest* 2000;49:221–6.
- [22] Fan HC, Blumenfeld YJ, Chitkara U, Hudgins L, Quake SR. Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proc Natl Acad Sci U S A* 2008;105:16266–71.
- [23] Hahn S, Lapaire O, Tercanli S, Kolla V, Hosli I. Determination of fetal chromosome aberrations from fetal DNA in maternal blood: has the challenge finally been met? *Expert Rev Mol Med* 2011;13:e16.
- [24] Bianchi DW, Lo YM. Fetomaternal cellular and plasma DNA trafficking: the Yin and the Yang. *Ann N Y Acad Sci* 2001;945:119–31.
- [25] Liao GJ, Gronowski AM, Zhao Z. Non-invasive prenatal testing using cell-free fetal DNA in maternal circulation. *Clin Chim Acta* 2014;428:44–50.
- [26] Pongsritasana T, Wongratanaheewin S, Prasertcharoensuk V, Sermswan RW. Isolation of fetal nucleated red blood cell from maternal blood using immunomagnetic beads for prenatal diagnosis. *Asian Pac J Allergy Immunol* 2006;24:65–71.
- [27] Wachtel SS, Shulman LP, Sammons D. Fetal cells in maternal blood. *Clin Genet* 2001;59:74–9.
- [28] Papageorgiou EA, Patsalis PC. Non-invasive prenatal diagnosis of aneuploidies: new technologies and clinical applications. *Genome Med* 2012;4:46.
- [29] Wald NJ, Cuckle HS, Densem JW, Nanchahal K, Royston P, Chard T, et al. Maternal serum screening for Down's syndrome in early pregnancy. *BMJ* 1988;297:883–7.
- [30] Shaw SW, Lin SY, Lin CH, Su YN, Cheng PJ, Lee CN, et al. Second-trimester maternal serum quadruple test for Down syndrome screening: a Taiwanese population-based study. *Taiwan J Obstet Gynecol* 2010;49:30–4.
- [31] Phillips OP, Elias S, Shulman LP, Andersen RN, Morgan CD, Simpson JL. Maternal serum screening for fetal Down syndrome in women less than 35 years of age using alpha-fetoprotein, hCG, and unconjugated estriol: a prospective 2-year study. *Obstet Gynecol* 1992;80:353–8.
- [32] Chao AS, Chung CL, Wu CD, Chang SD, Cheng PJ, Lin YT, et al. Second trimester maternal serum screening using alpha fetoprotein, free beta human chorionic gonadotropin and maternal age specific risk: result of chromosomal abnormalities detected in screen positive for Down syndrome in an Asian population. *Acta Obstet Gynecol Scand* 1999;78:393–7.
- [33] Hsiao CH, Cheng PJ, Shaw SW, Hsu JJ, Chen RC, Tseng YJ, et al. Extended First-trimester screening using multiple sonographic markers and maternal serum biochemistry: A five-year prospective study. *Fetal Diagn Ther* 2014;35:296–301.
- [34] Chen XQ, Stroun M, Magnenat JL, Nicod LP, Kurt AM, Lyautey J, et al. Microsatellite alterations in plasma DNA of small cell lung cancer patients. *Nat Med* 1996;2:1033–5.
- [35] Nawroz H, Koch W, Anker P, Stroun M, Sidransky D. Microsatellite alterations in serum DNA of head and neck cancer patients. *Nat Med* 1996;2:1035–7.
- [36] Lo YM, Corbetta N, Chamberlain PF, Rai V, Sargent IL, Redman CW, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet* 1997;350:485–7.
- [37] Poon LL, Leung TN, Lau TK, Lo YM. Presence of fetal RNA in maternal plasma. *Clin Chem* 2000;46:1832–4.
- [38] Lo YM, Tein MS, Lau TK, Haines CJ, Leung TN, Poon PM, et al. Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis. *Am J Hum Genet* 1998;62:768–75.
- [39] Saito H, Sekizawa A, Morimoto T, Suzuki M, Yanaiharu T. Prenatal DNA diagnosis of a single-gene disorder from maternal plasma. *Lancet* 2000;356:1170.
- [40] Chiu RW, Lau TK, Leung TN, Chow KC, Chui DH, Lo YM. Prenatal exclusion of beta thalassaemia major by examination of maternal plasma. *Lancet* 2002;360:998–1000.
- [41] Vrettou C, Traeger-Synodinos J, Tzetzis M, Malamis G, Kanavakis E. Rapid screening of multiple beta-globin gene mutations by real-time PCR on the LightCycler: application to carrier screening and prenatal diagnosis of thalassaemia syndromes. *Clin Chem* 2003;49:769–76.
- [42] Li Y, Di Naro E, Vitucci A, Zimmermann B, Holzgreve W, Hahn S. Detection of paternally inherited fetal point mutations for beta-thalassemia using size-fractionated cell-free DNA in maternal plasma. *JAMA* 2005;293:843–9.
- [43] Chan KC, Ding C, Gerovassili A, Yeung SW, Chiu RW, Leung TN, et al. Hyper-methylated RASSF1A in maternal plasma: A universal fetal DNA marker that improves the reliability of noninvasive prenatal diagnosis. *Clin Chem* 2006;52:2211–8.
- [44] Lun FM, Chiu RW, Allen Chan KC, Yeung Leung T, Kin Lau T, Dennis Lo YM. Microfluidics digital PCR reveals a higher than expected fraction of fetal DNA in maternal plasma. *Clin Chem* 2008;54:1664–72.
- [45] Lau TK, Chan MK, Lo PS, Chan HY, Chan WS, Koo TY, et al. Clinical utility of noninvasive fetal trisomy (NIFTY) test—early experience. *J Matern Fetal Neonatal Med* 2012;25:1856–9.
- [46] Ehrlich M, Deciu C, Zwiefelhofer T, Tynan JA, Cagasan L, Tim R, et al. Noninvasive detection of fetal trisomy 21 by sequencing of DNA in maternal blood: a study in a clinical setting. *Am J Obstet Gynecol* 2011;204. 205.e201–11.
- [47] Sehnert AJ, Rhee B, Comstock D, de Feo E, Heilek G, Burke J, et al. Optimal detection of fetal chromosomal abnormalities by massively parallel DNA sequencing of cell-free fetal DNA from maternal blood. *Clin Chem* 2011;57:1042–9.
- [48] Lau TK, Chen F, Pan X, Pooh RK, Jiang F, Li Y, et al. Noninvasive prenatal diagnosis of common fetal chromosomal aneuploidies by maternal plasma DNA sequencing. *J Matern Fetal Neonatal Med* 2012;25:1370–4.
- [49] Shaw SW, Hsiao CH, Chen CY, Ren Y, Tian F, Tsai C, et al. Noninvasive prenatal testing for whole fetal chromosomal aneuploidies: a multicenter prospective cohort trial in Taiwan. *Fetal Diagn Ther* 2014;35:13–7.
- [50] Jiang F, Ren J, Chen F, Zhou Y, Xie J, Dan S, et al. Noninvasive Fetal Trisomy (NIFTY) test: an advanced noninvasive prenatal diagnosis methodology for fetal autosomal and sex chromosomal aneuploidies. *BMC Med Genomics* 2012;5:57.
- [51] Lapaire O, Holzgreve W, Oosterwijk JC, Brinkhaus R, Bianchi DW. Georg Schmorl on trophoblasts in the maternal circulation. *Placenta* 2007 Jan;28(1):1–5.
- [52] Huang Z, Fong CY, Gauthaman K, Sukumar P, Choolani M, Bongso A. Novel approaches to manipulating foetal cells in the maternal circulation for non-invasive prenatal diagnosis of the unborn child. *J Cell Biochem* 2011;112:1475–85.
- [53] Sargent IL, Johansen M, Chua S, Redman CW. Clinical experience: isolating trophoblasts from maternal blood. *Ann New Y Acad Sci* 1994;731:154–61.
- [54] Durrant LG, Martin WL, McDowall KM, Liu DT. Isolation of fetal trophoblasts and nucleated erythrocytes from the peripheral blood of pregnant women for prenatal diagnosis of fetal aneuploidies. *Early Hum Dev* 1996;47:S79–83.
- [55] Johnson P, Cheng H, Molloy C, Stern C, Slade M. Human trophoblast-specific surface antigens identified using monoclonal antibodies. *Am J Reprod Immunol* 1981;1:246–54.
- [56] Bertero MT, Camaschella C, Serra A, Bergui L, Caligaris-Cappio F. Circulating 'trophoblast' cells in pregnancy have maternal genetic markers. *Prenatal Diagn* 1988;8:585–90.
- [57] Henderson KG, Shaw TE, Barrett IJ, Telenius AH, Wilson RD, Kalousek DK. Distribution of mosaicism in human placenta. *Hum Genetics* 1996;97:650–4.
- [58] Goldberg JD, Wohlferd MM. Incidence and outcome of chromosomal mosaicism found at the time of chorionic villus sampling. *Am J Obstet Gynecol* 1997;176:1349–52. discussion 1352–3.
- [59] Walknowska J, Conte FA, Grumbach MM. Practical and theoretical implications of fetal-maternal lymphocyte transfer. *Lancet* 1969;1:1119–22.
- [60] Johansen JK, Schacke E, Sturup AG. Foeto-maternal transfusion in relation to delivery. *Ugeskr Laeger* 1971;133:870–2.
- [61] Grosset L, Barrelet V, Odartchenko N. Antenatal fetal sex determination from maternal blood during early pregnancy. *Am J Obstet Gynecol* 1974;120:60–3.

- [62] Herzenberg LA, Bianchi DW, Schroder J, Cann HM, Iverson GM. Fetal cells in the blood of pregnant women: detection and enrichment by fluorescence-activated cell sorting. *Proc Natl Acad Sci U S A* 1979;76:1453–5.
- [63] Bianchi DW, Zickwolf GK, Weil GJ, Sylvester S, DeMaria MA. Male fetal progenitor cells persist in maternal blood for as long as 27 years postpartum. *Proc Natl Acad Sci U S A* 1996;93:705–8.
- [64] Calabrese G, Baldi M, Fantasia D, Sessa MT, Kalantar M, Holzhauser C, et al. Detection of chromosomal aneuploidies in fetal cells isolated from maternal blood using single-chromosome dual-probe FISH analysis. *Clin Genet* 2012;82:131–9.
- [65] D'Souza E, Ghosh K, Colah R. A comparison of the choice of monoclonal antibodies for recovery of fetal cells from maternal blood using FACS for noninvasive prenatal diagnosis of hemoglobinopathies. *Cytometry B Clin Cytome* 2009;76:175–80.
- [66] Babochkina T, Mergenthaler S, Lapaire O, Kiefer V, Yura H, Koike K, et al. Evaluation of a soybean lectin-based method for the enrichment of erythroblasts. *J Histochem Cytochem* 2005;53:329–30.
- [67] Mohamed H, Turner JN, Caggana M. Biochip for separating fetal cells from maternal circulation. *J Chromatog A* 2007;1162:187–92.
- [68] Samura O, Sekizawa A, Zhen DK, Falco VM, Bianchi DW. Comparison of fetal cell recovery from maternal blood using a high density gradient for the initial separation step: 1.090 versus 1.119 g/mL. *Prenatal Diagn* 2000;20:281–6.
- [69] Ponnusamy S, Mohammed N, Ho SS, Zhang HM, Chan YH, Ng YW, et al. In vivo model to determine fetal-cell enrichment efficiency of novel noninvasive prenatal diagnosis methods. *Prenatal Diagn* 2008;28:494–502.
- [70] Prieto B, Alonso R, Paz A, Cándenas M, Venta R, Ladenson JH, et al. Optimization of nucleated red blood cell (NRBC) recovery from maternal blood collected using both layers of a double density gradient. *Prenatal Diagn* 2001;21:187–93.
- [71] Huang R, Barber TA, Schmidt MA, Tompkins RG, Toner M, Bianchi DW, et al. A microfluidics approach for the isolation of nucleated red blood cells (NRBCs) from the peripheral blood of pregnant women. *Prenatal Diagn* 2008;28:892–9.
- [72] Holzgreve W, Ghezzi F, Di Naro E, Ganshirt D, Maymon E, Hahn S. Disturbed fetomaternal cell traffic in preeclampsia. *Obstet Gynecol* 1998;91:669–72.
- [73] Kavanagh DM, Kersaudy-Kerhoas M, Dhariwal RS, Desmulliez MP. Current and emerging techniques of fetal cell separation from maternal blood. *J Chromatogr. B Analyt Technol Biomed Life Sci* 2010;878:1905–11.
- [74] Choolani M, O'Donoghue K, Talbert D, Kumar S, Roberts I, Letsky E, et al. Characterization of first trimester fetal erythroblasts for non-invasive prenatal diagnosis. *Mol Hum Reprod* 2003;9:227–35.
- [75] Palis J. To condense is not to die. *Blood* 2005;106:1900.
- [76] Spivak JL. The anaemia of cancer: death by a thousand cuts. *Nat Rev Cancer* 2005;5:543–55.
- [77] Brittain T. Molecular aspects of embryonic hemoglobin function. *Mol Aspects Med* 2002;23:293–342.
- [78] Manning JM, Popowicz AM, Padovan JC, Chait BT, Manning LR. Intrinsic regulation of hemoglobin expression by variable subunit interface strengths. *FEBS J* 2012;279:361–9.
- [79] Choolani M, O'Donnell H, Campagnoli C, Kumar S, Roberts I, Bennett PR, et al. Simultaneous fetal cell identification and diagnosis by epsilon-globin chain immunophenotyping and chromosomal fluorescence in situ hybridization. *Blood* 2001;98:554–7.
- [80] Sorensen MD, Gonzalez Dosal R, Jensen KB, Christensen B, Kølvråa S, Jensen UB, et al. Epsilon haemoglobin specific antibodies with applications in noninvasive prenatal diagnosis. *J Biomed Biotechnol* 2009;2009:659219.
- [81] Al-Mufti R, Hambley H, Farzaneh F, Nicolaides KH. Fetal and embryonic hemoglobins in erythroblasts of chromosomally normal and abnormal fetuses at 10–40 weeks of gestation. *Haematologica* 2000;85:690–3.
- [82] Ikeya M, Shinya M, Kitagawa M. Basic investigation of the lectin method for separation and recovery of nucleated red blood cells in maternal blood, and a study into the frequency of nucleated red blood cells in fetomaternal disorders. *Congenit Anom (Kyoyo)* 2005;45:26–31.
- [83] Jeon YJ, Kwon KH, Kim JW, Pang M-G, Jung SC, Kim YJ. Comparison in the yield of fetal nucleated red blood cell between the first-and second-trimester using double density gradient centrifugation. *Korean J Obstet Gynecol* 2010;53:127–36.
- [84] Bischoff FZ, Marquez-Do DA, Martinez DI, Dang D, Horne C, Lewis D, et al. Intact fetal cell isolation from maternal blood: improved isolation using a simple whole blood progenitor cell enrichment approach (RosetteSep). *Clin Genet* 2003;63:483–9.
- [85] Yan J, Guilbault E, Masse J, Bronsard M, DeGrandpré P, Forest JC, et al. Optimization of the fluorescence in situ hybridization (FISH) technique for high detection efficiency of very small proportions of target interphase nuclei. *Clin Genet* 2000;58:309–18.
- [86] Posel C, Moller K, Frohlich W, Schulz I, Boltze J, Wagner DC. Density gradient centrifugation compromises bone marrow mononuclear cell yield. *PLoS One* 2012;7:e50293.
- [87] Purwosunu Y, Sekizawa A, Koide K, Okazaki S, Farina A, Okai T. Clinical potential for noninvasive prenatal diagnosis through detection of fetal cells in maternal blood. *Taiwan J Obstet Gynecol* 2006;45:10–20.
- [88] Ganshirt-Ahlert D, Borjesson-Stoll R, Burschkyk M, Dohr A, Garritsen HS, Helmer E, et al. Detection of fetal trisomies 21 and 18 from maternal blood using triple gradient and magnetic cell sorting. *Am J Reprod Immunol Sep-Oct* 1993;30:194–201.
- [89] Secker-Walker RH, Solomon LJ, Flynn BS, LePage SS, Crammond JE, Worden JK, et al. Training obstetric and family practice residents to give smoking cessation advice during prenatal care. *Am J Obstet Gynecol* 1992;166:1356–63.
- [90] Bianchi DW, Flint AF, Pizzimenti MF, Knoll JH, Latt SA. Isolation of fetal DNA from nucleated erythrocytes in maternal blood. *Proc Natl Acad Sci U S A* 1990;87:3279–83.
- [91] Henri A, Testa U, Tonthat H, Riou J, Titeux M, Vainchenker W, et al. Disappearance of Hb F and i antigen during the first year of life. *Am J Hematol* 1980;9:161–70.
- [92] Twu YC, Chen CP, Hsieh CY, Tzeng CH, Sun CF, Wang SH, et al. I branching formation in erythroid differentiation is regulated by transcription factor C/EBPalpha. *Blood* 2007;110:4526–34.
- [93] Liu C, Stakenborg T, Peeters S, Lagae L. Cell manipulation with magnetic particles toward microfluidic cytometry. *J Appl Phys* 2009;105. <http://dx.doi.org/10.1063/1.3116091>.
- [94] Lee BS, Lee JN, Park JM, Lee JG, Kim S, Cho YK, et al. A fully automated immunoassay from whole blood on a disc. *Lab Chip* 2009;9:1548–55.
- [95] Chen KC, Lee TP, Pan YC, Chiang CL, Chen CL, Yang YH, et al. Detection of circulating endothelial cells via a microfluidic disk. *Clin Chem* 2011;57:586–92.
- [96] Autebert J, Coudert B, Bidard FC, Pierga JY, Descroix S, Malaquin L, et al. Microfluidic: an innovative tool for efficient cell sorting. *Methods* 2012;57:297–307.
- [97] Chen CL, Chen KC, Pan YC, Lee TP, Hsiung LC, Lin CM, et al. Separation and detection of rare cells in a microfluidic disk via negative selection. *Lab Chip* 2011;11:474–83.
- [98] Mavrou A, Kouvidi E, Antsaklis A, Souka A, Kitsiou Tzeli S, Kolialexi A. Identification of nucleated red blood cells in maternal circulation: a second step in screening for fetal aneuploidies and pregnancy complications. *Prenatal Diagn* 2007;27:150–3.