



Research Letter

Prenatal diagnosis of chromosome 8p23.1 microdeletion by array comparative genomic hybridization using uncultured amniocytes in a pregnancy associated with fetal partial corpus callosum agenesis and schizencephaly



Chih-Ping Chen^{a, b, c, d, e, f, *}, Cheng-Ran Peng^a, Tung-Yao Chang^g, Wan-Yuo Guo^h, Yen-Ni Chen^a, Peih-Shan Wuⁱ, Dai-Dyi Town^a, Wayseen Wang^{b, j}

^a Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan

^c Department of Biotechnology, Asia University, Taichung, Taiwan

^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^g Taiji Fetal Medicine Center, Taipei, Taiwan

^h Department of Radiology, Taipei Veterans General Hospital, Taipei, Taiwan

ⁱ Gene Biodesign Co. Ltd, Taipei, Taiwan

^j Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 23 October 2015

Dear Editor

We previously reported a *de novo* 0.7-Mb deletion in sub-band 8p23.1 extending from 7,227,000 to 7,916,187 (NCBI 36/hg 18) overlapping the telomeric β -defensin gene cluster, but outside the 8p23.1 deletion critical region, in a fetus with congenital diaphragmatic hernia (CDH) and additional abnormalities [1]. Herein, we additionally report a fetus with the copy number variable β -defensin gene cluster on 8p23.1 and partial corpus callosum agenesis and schizencephaly. To our knowledge, such a case has not previously been described.

A 29-year-old, gravida 5, para 2, Taiwanese woman was referred for genetic counseling at 24 weeks of gestation because of fetal chromosome aberration and fetal brain abnormalities. This was the fifth pregnancy of the woman. She had one 3-year-old healthy child and had delivered a baby with chromosome 1p32-p31 deletion syndrome, ventriculomegaly, and corpus callosum hypogenesis. The parents had normal karyotypes. During this pregnancy, at 16 weeks of gestation, the woman underwent amniocentesis which revealed a karyotype of 45,X[4]/46,XY[31]. Prenatal ultrasound at

22 weeks of gestation revealed the absence of a septum pellucidum, partial agenesis of the corpus callosum, and wide splaying of the anterior horn of the lateral ventricles. Fetal magnetic resonance imaging examination showed partial corpus callosum agenesis with only the presence of the genu, and schizencephaly at the right occipital lobe. Repeated amniocentesis was performed at 24 weeks of gestation. Cytogenetic analysis of the cultured amniocytes revealed a karyotype of 45,X[6]/46,XY[9]. Array comparative genomic hybridization (aCGH) analysis by NimbleGen ISCA Plus Cytogenetic Array (Roche NimbleGen, Madison, WI, USA) using uncultured amniocytes revealed a 0.57-Mb 8p23.1 microdeletion or arr 8p23.1 (7,235,388–7,809,256) \times 1.3 (Build GRCh37, Feb 2009, hg19) encompassing 44 genes including 14 OMIM genes of FAM90A15P, FAM90A3P, FAM90A13P, FAM90A5P, FAM90A20P, DEFB103B, SPAG11B, FAM90A7P, FAM90A14P, FAM90A18P, FAM90A8P, FAM90A19P, FAM90A9P, and FAM90A10P. The parents did not have such a deletion. The parents decided to terminate the pregnancy, and a fetus was delivered with no abnormalities in gross appearance.

The β -defensin gene cluster on the chromosome band of 8p23.1 has been shown to be one of the most prominent examples of functional-relevant copy number variation (CNV), and various studies have demonstrated that multisite variants and CNV of the defensin-encoding genes are associated with increased risk for various diseases including cancer and inflammatory diseases [2–6].

The present case has CNV on the β -defensin gene cluster at 8p23.1, but without the involvement of the 8p23.1 deletion syndrome critical genes such as GATA4, NEIL2, SOX7, and TNKS. The 8p23.1 deletion syndrome includes congenital heart defects,

* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

microcephaly, mental retardation, CDH, psychomotor developmental delay, and behavior problems [7–12]. *GATA4*, *NEIL2*, and *SOX7* have been associated with congenital heart defects and CDH [12–19]. Haploinsufficiency of either *GATA4* or *SOX7* has been shown to contribute to the development of CDH in individuals with 8p23.1 deletions [20]. *TNKS* has also been proposed as a novel candidate for the Cornelia de Lange syndrome [21].

The peculiar aspect of this case is the concomitant occurrence of fetal partial corpus callosum agenesis, schizencephaly, and partial deletion of *DEFB103* and *SPAG11* on chromosome 8 β -defensin gene cluster. There is increasing evidence that abnormal defensin expression due to CNV of the β -defensin gene cluster of *DEFB107*, *DEFB106*, *DEFB105*, *DEFB104*, *DEFB103*, *DEFB4*, and *SPAG11B* is related to infectious diseases, sperm dysfunction, and infertility in males [22,23]. β -Defensins are a family of multifunctional peptides important for immunity, defense against pathogens, reproduction, and pigmentation [24,25].

A worldwide analysis of β -defensin CNV has found an unusually high frequency of high-*DEFB103*-expressing copies in East Asia [24]. A clinical report of central nervous system abnormalities associated with CNV on the β -defensin gene cluster at 8p23.1 has not previously been presented. The correlation of partial deletion of *DEFB103* with fetal brain abnormalities in this Taiwanese case is unclear and will require more case studies for confirmation.

In the present case, the conventional cytogenetic analysis of cultured amniocytes revealed mosaicism for 45,X and 46,XY, whereas, aCGH analysis on uncultured amniocytes revealed microdeletion of 8p23.1. The aCGH analysis has the advantage of detecting microdeletion syndrome in fetuses with a normal karyotype by conventional cytogenetic analysis [26,27]. Central nervous system abnormality is not the clinical phenotype of mosaic Turner syndrome. We think that aCGH provides the advantage of better understanding of the nature of microdeletion of 8p23.1 and genotype-phenotype correlation in this case.

Conflict of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by research grants NSC-101-2314-B-195-011-MY3 and MOST 103-2314-B-195-010 from the Ministry of Science and Technology and MMH-E-104-04 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Chen CP, Wang TH, Chen YJ, Chang TY, Liu YP, Tzen CY, et al. Prenatal diagnosis of Fryns syndrome associated with a microdeletion at 8p23.1. *Prenat Diagn* 2007;27:967–9.
- [2] Groth M, Szafranski K, Taudien S, Huse K, Mueller O, Rosenstiel P, et al. High-resolution mapping of the 8p23.1 beta-defensin cluster reveals strictly concordant copy number variation of all genes. *Hum Mutat* 2008;29:1247–54.
- [3] Groth M, Wiegand C, Szafranski K, Huse K, Kramer M, Rosenstiel P, et al. Both copy number and sequence variations affect expression of human *DEFB4*. *Genes Immun* 2010;11:458–66.
- [4] Taudien S, Groth M, Huse K, Petzold A, Szafranski K, Hampe J, et al. Haplotyping and copy number estimation of the highly polymorphic human beta-defensin locus on 8p23 by 454 amplicon sequencing. *BMC Genomics* 2010;11:252.
- [5] Taudien S, Gabel G, Kuss O, Groth M, Grützmann R, Huse K, et al. Association studies of the copy-number variable β -defensin cluster on 8p23.1 in adenocarcinoma and chronic pancreatitis. *BMC Res Notes* 2012;5:629.
- [6] Taudien S, Huse K, Groth M, Platzer M. Narrowing down the distal border of the copy number variable beta-defensin gene cluster on human 8p23. *BMC Res Notes* 2014;7:93.
- [7] Hutchinson R, Wilson M, Voullaire L. Distal 8p deletion (8p23.1→8pter): a common deletion? *J Med Genet* 1992;29:407–11.
- [8] Marino B, Reale A, Giannotti A, Digilio MC, Dallapiccola B. Nonrandom association of atrioventricular canal and del(8p) syndrome. *Am J Med Genet* 1992;42:424–7.
- [9] Digilio MC, Marino B, Dallapiccola B. Atrioventricular canal and 8p- syndrome. *Am J Med Genet* 1993;47:437–8.
- [10] Claeys I, Holvoet M, Eyskens B, Adriansens P, Gewillig M, Fryns J-P, et al. A recognisable behavioural phenotype associated with terminal deletions of the short arm of chromosome 8. *Am J Med Genet* 1997;74:515–20.
- [11] Faivre L, Morichon-Delvallez N, Viot G, Nancy F, Loison S, Mandelbrot L, et al. Prenatal diagnosis of an 8p23.1 deletion in a fetus with diaphragmatic hernia and review of the literature. *Prenat Diagn* 1998;18:1055–60.
- [12] Ballarati L, Cereda A, Caselli R, Selicorni A, Recalcati MP, Maitz S, et al. Genotype-phenotype correlations in a new case of 8p23.1 deletion and review of the literature. *Eur J Med Genet* 2011;54:55–9.
- [13] Pehlivan T, Pober BR, Brueckner M, Garrett S, Slaugh R, Van Rheeden R, et al. *GATA4* haploinsufficiency in patients with interstitial deletion of chromosome region 8p23.1 and congenital heart disease. *Am J Med Genet* 1999;83:201–6.
- [14] Reddy KS. A paternally inherited terminal deletion, del(8)(p23.1)pat, detected prenatally in an amniotic fluid sample: a review of deletion 8p23.1 cases. *Prenat Diagn* 1999;19:868–72.
- [15] Wat MJ, Shchelochkov OA, Holder AM, Breman AM, Dagli A, Bacino C, et al. Chromosome 8p23.1 deletions as a cause of complex congenital heart defects and diaphragmatic hernia. *Am J Med Genet* 2009;149A:1661–77.
- [16] Hoellen F, Weichert J. Sonographic diagnosis of multiple cardiovascular malformations in a fetus with an interstitial 8p23.1 deletion. *J Ultrasound Med* 2012;31:1689–92.
- [17] Longoni M, Lage K, Russell MK, Loscertales M, Abdul-Rahman OA, Baynam G, et al. Congenital diaphragmatic hernia interval on chromosome 8p23.1 characterized by genetics and protein interaction networks. *Am J Med Genet* 2012;158A:3148–58.
- [18] Guimiot F, Dupont C, Fuentes-Duarte A, Aboura A, Bazin A, Khung-Savatovsky S, et al. Maternal transmission of interstitial 8p23.1 deletion detected during prenatal diagnosis. *Am J Med Genet* 2013;161A:208–13.
- [19] Keitges EA, Pasion R, Burnside RD, Mason C, Gonzalez-Ruiz A, Dunn T, et al. Prenatal diagnosis of two fetuses with deletions of 8p23.1, critical region for congenital diaphragmatic hernia and heart defects. *Am J Med Genet* 2013;161A:1755–8.
- [20] Wat MJ, Beck TF, Hernández-García A, Yu Z, Veenma D, Garcia M, et al. Mouse model reveals the role of *SOX7* in the development of congenital diaphragmatic hernia associated with recurrent deletions of 8p23.1. *Hum Mol Genet* 2012;21:4115–25.
- [21] Baynam G, Goldblatt J, Walpole I. Deletion of 8p23.1 with features of Cornelia de Lange syndrome and congenital diaphragmatic hernia and a review of deletions of 8p23.1 to 8pter? A further locus for Cornelia de Lange syndrome. *Am J Med Genet* 2008;146A:1565–70.
- [22] Zhou YS, Webb S, Lettice L, Tardif S, Kilanowski F, Tyrrell C, et al. Partial deletion of chromosome 8 β -defensin cluster confers sperm dysfunction and infertility in male mice. *PLoS Genet* 2013;9:e1003826.
- [23] Jones EA, Kananurak A, Bevins CL, Hollox EJ, Bakaletz LO. Copy number variation of the beta defensin gene cluster on chromosome 8p influences the bacterial microbiota within the nasopharynx of otitis-prone children. *PLoS One* 2014;9:e98269.
- [24] Hardwick RJ, Machado LR, Zuccherato LW, Antolinos S, Xue Y, Shaw N, et al. A worldwide analysis of beta-defensin copy number variation suggests recent selection of a high-expressing *DEFB103* gene copy in East Asia. *Hum Mutat* 2011;32:743–50.
- [25] Ottolini B, Hornsby MJ, Abujaber R, MacArthur JAL, Badge RM, Schwarzscher T, et al. Evidence of convergent evolution in humans and macaques supports an adaptive role for copy number variation of the β -defensin-2 gene. *Genome Biol Evol* 2014;6:3025–38.
- [26] Chen CP, Chang SD, Wang TH, Wang LK, Tsai JD, Liu YP, et al. Detection of recurrent transmission of 17q12 microdeletion by array comparative genomic hybridization in a fetus with prenatally diagnosed hydronephrosis, hydro-ureter and multicystic kidney, and variable clinical spectrum in the family. *Taiwan J Obstet Gynecol* 2013;52:551–7.
- [27] Chen CP, Lin CJ, Chern SR, Liu YP, Kuo YL, Chen YN, et al. Prenatal diagnosis and molecular cytogenetic characterization of a 1.07-Mb microdeletion at 5q35.2-q35.3 associated with *NSD1* haploinsufficiency and Sotos syndrome. *Taiwan J Obstet Gynecol* 2014;53:583–7.