



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Case Report

Detection of mosaic 15q11.1-q11.2 deletion encompassing *NBEAP1* and *POTEB* in a fetus with diffuse lymphangiomatosis

Chih-Ping Chen^{a, b, c, d, e, f, *}, Kuo-Gon Wang^a, Hsu-Kuang Huang^g, Cheng-Ran Peng^g,
Schu-Rern Chern^b, Peih-Shan Wu^h, Yen-Ni Chen^a, Shin-Wen Chen^a, Chen-Chi Lee^a,
Wayseen Wang^{b, i}

^a Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan

^b Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan

^c Department of Biotechnology, Asia University, Taichung, Taiwan

^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan

^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan

^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan

^g Department of Obstetrics and Gynecology, Taiwan Adventist Hospital, Taipei, Taiwan

^h Gene Biodesign Co. Ltd, Taipei, Taiwan

ⁱ Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 12 January 2017

Keywords:

15q11.1-q11.2 deletion
diffuse lymphangiomatosis
NBEAP1
POTEB

ABSTRACT

Objective: We present cytogenetic and molecular cytogenetic diagnoses of mosaic deletion of chromosome 15q11.1-q11.2 in a fetus with diffuse lymphangiomatosis.

Case Report: A 33-year-old woman underwent amniocentesis at 22 weeks of gestation because of fetal diffuse lymphangiomatosis involving left-side chest, abdominal cavity, thigh and vulva, and intrauterine growth restriction. Amniocentesis revealed a karyotype of 46,XX,del(15)(q11.1q11.2)[9]/46,XX[26]. The mother had a karyotype of 46,XX. The father had a karyotype of 46,XY. The parents elected to terminate the pregnancy. A 610-g female fetus was delivered at 23 weeks of gestation with large cystic lymphangioma over the left abdomen, thigh, and vulva. The umbilical cord had a karyotype of 46,XX,del(15)(q11.1q11.2)[24]/46,XX[16]. The placental tissue had a karyotype of 46,XX,del(15)(q11.1q11.2)[23]/46,XX[17]. Array comparative genomic hybridization analysis of the umbilical cord and placenta revealed a 2.42-Mb deletion of 15q11.1-q11.2 encompassing the genes of *NBEAP1* and *POTEB*.

Conclusion: Deletion of 15q11.1-q11.2 encompassing *NBEAP1* and *POTEB* may be associated with diffuse lymphangiomatosis.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Diffuse lymphangiomatosis is a rare congenital malformation of the lymphatics caused by primary lymphatic dysplasia and a lymphatic proliferative process that may involve soft tissues, visceral organs, bones, retroperitoneum, and eyes [1–3]. The cause of diffuse lymphangiomatosis is poorly understood, and the genetic

basis of this disorder has rarely been investigated. We present perinatal detection of mosaic 15q11.1-q11.2 deletion encompassing *NBEAP1* and *POTEB* in a fetus with diffuse lymphangiomatosis. Our finding may provide a novel genotype–phenotype correlation of this disorder.

Case report

A 33-year-old gravida 1, para 0 woman underwent amniocentesis at 22 weeks of gestation because of abnormal fetal ultrasound findings. Her husband was 35 years old, and there was no family history of lymphangiomatosis. Level 2 fetal ultrasound

* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, 10449, Taiwan.
E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).

examination revealed unilateral diffuse lymphangiomatosis involving left-side chest, abdominal cavity, thigh and vulva, and intrauterine growth restriction (IUGR). Amniocentesis revealed a karyotype of 46,XX,del(15)(q11.1q11.2)[9]/ 46,XX[26]. The mother had a karyotype of 46,XX. The father had a karyotype of 46,XY. The parents elected to terminate the pregnancy. A 610-g female fetus was delivered at 23 weeks of gestation with large cystic lymphangioma over the left abdomen, thigh, and vulva. Postnatal cytogenetic analysis revealed a karyotype of 46,XX,del(15)(q11.1q11.2)[24]/46,XX[16] in the umbilical cord and a karyotype of 46,XX,del(15)(q11.1q11.2)[23]/46,XX[17] in the placental tissue (Figure 1). Array comparative genomic hybridization (aCGH) analysis of the umbilical cord using Roche ISCA Plus Cytogenetic Array (Roche NimbleGen, Madison, WI, USA) revealed a result of a 2.39-Mb deletion of 15q11.1-q11.2 or arr 15q11.1q11.2 (20,225,177–22,617,226)×1.15 encompassing 79 genes including two Online Mendelian Inheritance in Man (OMIM) genes of *NBEAP1* and *POTEB*. The log2 ratio was −0.8 equivalent to 85% mosaicism for the deletion. aCGH analysis of the placental tissue revealed a result of a 2.42-Mb deletion of 15q11.1-q11.2 or arr 15q11.1q11.2 (20,200,762–22,617,226)×1.2 encompassing 79 genes including two OMIM genes of *NBEAP1* and *POTEB* (Figure 2). The log2 ratio was −0.75 equivalent to 80% mosaicism for the deletion. aCGH analysis of the maternal peripheral blood revealed a result of a 0.44-Mb microdeletion of 15q11.2 or arr 15q11.2 (22,179,070–22,617,226)×1.19 encompassing 16 genes including no OMIM gene. aCGH analysis of the

paternal peripheral blood revealed a result of a 0.99-Mb microdeletion of 15q11.1-q11.2 or arr 15q11.1q11.2 (20,213,575–21,205,149)×1.18 encompassing 23 genes including no OMIM gene and a result of 0.47-Mb microdeletion of 15q11.2 or arr 15q11.2 (22,083,345–22,557,608)×1.16 encompassing 18 genes including no OMIM gene.

Discussion

The peculiar aspect of the present case is the association of 15q11.1-q11.2 deletion with diffuse lymphangiomatosis involving the deficiency of *NBEAP1* and *POTEB*. *POTEB* (OMIM 608912) belongs to the pote ankyrin domain family *POTE* (prostate-, ovary-, testis-, and placenta-expressed gene) [4,5]. *POTE* protein is associated with apoptotic cells and plays a role in apoptosis [6,7]. *NBEAP1* (OMIM 601889) or *BCL8* (B-cell CLL/lymphoma 8) is involved in translocation t(14;15)(q32;q11-13) breakpoint affecting band 15q11-13 in diffuse large-cell lymphoma [8]. *NBEAP1* belongs to an evolutionarily conserved human gene family that encodes proteins with protein kinase A anchoring function [9].

The present fetus and her family had copy number variation (CNV) at 15q11.1-q11.2. However, only the present affected fetus had deficiency of *NBEAP1* and *POTEB* and diffuse lymphangiomatosis. Whether such a correlation is a coincidence or a pathogenic effect is unclear and will require more investigations to prove.





Figure 1. A karyotype of 46,XX,del(15)(q11.1q11.2). The arrows indicate the breakpoints.

A



B




Figure 2. Array comparative genomic hybridization analysis on the placental tissue reveals a result of a 2.42-Mb deletion of 15q11.1-q11.2 encompassing *NBEAP1* and *POTE*. (A) Chromosome zoom-in view; (B) chromosome 15.

Conflicts of interest

The authors have no conflicts of interest relevant to this article.

Acknowledgments

This work was supported by research grants MOST-104-2314-B-195-009 and MOST-105-2314-B-195-012 from the Ministry of Science and Technology and MMH-E-105-04 from MacKay Memorial Hospital, Taipei, Taiwan.

References

- [1] Singh S, Baboo ML, Pathak IC. Cystic lymphangioma in children: report of 32 cases including lesions at rare sites. *Surgery* 1971;69:947–51.
- [2] Lohrmann C, Foeldi E, Bartholomä J-P, Langer M. Disseminated lymphangiomatosis with skeletal involvement: detection with magnetic resonance lymphangiography. *Lymphology* 2007;40:74–80.
- [3] Lohrmann C, Foeldi E, Langer M. Assessment of the lymphatic system in patients with diffuse lymphangiomatosis by magnetic resonance imaging. *Eur J Radiol* 2011;80:576–81.
- [4] Bera TK, Zimonjic DB, Popescu NC, Sathyanarayana BK, Kumar V, Lee B, et al. *POTE*, a highly homologous gene family located on numerous chromosomes and expressed in prostate, ovary, testis, placenta, and prostate cancer. *Proc Natl Acad Sci USA* 2002;99:16975–80.

- [5] Bera TK, Huynh N, Maeda H, Sathyanarayana BK, Lee B, Pastan I. Five *POTE* paralogs and their splice variants are expressed in human prostate and encode proteins of different lengths. *Gene* 2004;337:45–53.
- [6] Liu XF, Bera TK, Liu LJ, Pastan I. A primate-specific POTE-actin fusion protein plays a role in apoptosis. *Apoptosis* 2009;14:1237–44.
- [7] Bera TK, Walker DA, Sherins RJ, Pastan I. POTE protein, a cancer-testis antigen, is highly expressed in spermatids in human testis and is associated with apoptotic cells. *Biochem Biophys Res Commun* 2012;417:1271–4.
- [8] Dyomin VG, Rao PH, Dalla-Favera R, Chaganti RSK. *BCL8*, a novel gene involved in translocations affecting band 15q11–13 in diffuse large-cell lymphoma. *Proc Natl Acad Sci USA* 1997;94:5728–32.
- [9] Dyomin VG, Chaganti SR, Dyomina K, Palanisamy N, Murty VVVS, Dalla-Favera R, et al. *BCL8* is a novel, evolutionarily conserved human gene family encoding proteins with presumptive protein kinase A anchoring function. *Genomics* 2002;80:158–65.