



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Short Communication

Can biparietal diameter-to-femur length ratio be a useful sonographic marker for screening thanatophoric dysplasia since the first trimester? A literature review of case reports and a retrospective study based on 10,293 routine fetal biometry measurements



Liangcheng Wang^a, Yasushi Takai^{a,*}, Kazunori Baba^a, Yukiko Mikami^a, Masahiro Saito^a, Isao Horiuchi^b, Ryo Konno^b, Kenjiro Takagi^b, Hiroyuki Seki^a

^a Center for Maternal, Fetal and Neonatal Medicine, Saitama Medical Center, Saitama Medical University, 1981 Kamoda, Kawagoe, Saitama 350-8550, Japan

^b Department of Obstetrics and Gynecology, Saitama Medical Center, Jichi Medical University, 1-847, Amanuma-cho, Omiya-ku, Saitama 330-8503, Japan

ARTICLE INFO

Article history:
Accepted 6 March 2017

Keywords:
Thanatophoric dysplasia
Biparietal diameter
Femur length
Skeletal dysplasia
Lethal skeletal dysplasia

ABSTRACT

Objective: The aim of the study was to determine whether the biparietal diameter/femur length (BPD/FL) ratio can be used to detect thanatophoric dysplasia in the first trimester of pregnancy.

Materials and Methods: Twenty-four reported cases of thanatophoric dysplasia diagnosed based on ultrasonographic results with molecular or radiographic diagnosis were included. All sonographic measurement records were extracted and reviewed, and the BPD/FL ratio was calculated for each gestational week. In addition, 10,293 routine fetal biometry measurements from 1395 cases of patients without skeletal dysplasia were compared.

Results: The BPD/FL ratio in the control group decreased to less than 3 prior to gestational week 13, and to less than 2 prior to week 18. Of the 27 BPD/FL ratios obtained from 24 cases of thanatophoric dysplasia, none was in the control range.

Conclusion: The BPD/FL ratio may be used to detect lethal skeletal dysplasias such as thanatophoric dysplasia since the first trimester.

© 2017 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Thanatophoric dysplasia (TD) is the most common neonatal lethal skeletal dysplasia [1]. Clinically, it is primarily characterized by remarkable shortening of limbs, narrow thorax, and macrocephaly. Pulmonary hypoplasia, secondary to fetal thoracic deformities, may result in severe respiratory insufficiency at birth. However, perinatal treatment remains very difficult. Consequently, early prenatal diagnosis is important because patients may choose termination if legal permission is limited by gestational weeks, or may choose not to undergo cesarean delivery of a fetus that will not survive. With advances in ultrasound technology, it is no longer difficult to observe details of fetal development in the first trimester such as sex [2], facial bones [3], and placental function [4].

However, TD is very rare, with prevalence as low as 1–6 per 100,000 [1]; only few were found in the first trimester in the reported cases. Therefore, establishing an easy early-screening method for less-experienced obstetricians remains a challenge. In general, shortened femurs on routine ultrasonography indicate skeletal dysplasia, and the prenatal sonographic detection rate for all cases is as high as 98.8% across all gestational ages [5]; however, lethal skeletal dysplasia is not distinguishable until the late second trimester or later [6,7]. Even if a short femoral bone is suspected on ultrasonography in the first or early second trimester, using a single ultrasound finding of short limbs to determine lethality might be unreliable because the accuracy of a single parameter may be affected by an incorrect estimated date of confinement. Therefore, combination with an additional fetal biometric parameter with minimal individual variation such as the biparietal diameter is considered necessary.

The aim of the present study was to determine whether the biparietal diameter/femur length ratio (BPD/FL) can be used to

* Corresponding author. Fax: +81 49 226 1495.

E-mail address: yastakai@saitama-med.ac.jp (Y. Takai).

screen for lethal skeletal dysplasias such as thanatophoric dysplasia since the first trimester.

Methods

This study was approved by both, the Ethics Committees of the Center for Maternal, Fetal and Neonatal Medicine, Saitama Medical University, Saitama Medical Center (SMUSMC), and the Ethics Committees of Jichi Medical University, Saitama Medical Center (JMUSMC). The need for informed consent was waived, as only retrospective review of medical records was performed.

We included all reported cases of TD with details of fetal ultrasound biometry that were confirmed with molecular analysis or postpartum radiology. The PubMed database was searched using the terms “thanatophoric dysplasia,” “first trimester,” “prenatal diagnosis,” and “molecular diagnosis.”

For comparison, fetal biometry records of controls without skeletal dysplasia evaluated at JMUSMC during 2009–2013 were examined. All fetal biometry measurement was performed by abdominal (Voluson E8; GE Healthcare, Zipf, Austria) or vaginal (SONOVISTA FX; Siemens Healthcare, Korea) ultrasound. All fetal ultrasonography measurement records were extracted from an electronic database, and outlier data due to entry errors were excluded; the maximum, minimum, median, and standard difference of the BPD/FL ratios at each gestational week were calculated.

Results

Twenty-four cases of TD with details of prenatal sonographic and molecular diagnosis were extracted from a PubMed database

search. The prenatal fetal biometry measurements and ultrasound findings of 24 cases are listed in Table 1. The median BPD/FL ratio of 27 records extracted from the 24 cases was 3.38 (range: 2.00–4.49). At JMUSMC, 10,293 routine fetal biometry measurements from 1395 cases without skeletal dysplasia were extracted, and 124 patients with recorded outliers due to entry errors were excluded. The median and range of BPD/FL ratios at weeks 12, 13, and 14 were 2.82 (1.89–4.06), 2.26 (1.68–2.87), and 2.05 (1.55–2.59), respectively; those at weeks 17, 18 and 19 were 1.70 (1.39–2.06), 1.65 (1.37–1.98) and 1.57 (1.30–1.87), respectively (Table 2). For all gestational ages, the BPD/FL ratio showed a continuous absolute decrease as a function of gestational weeks, with a rapid decrease during weeks 12–19. From weeks 20 to 36, the median of the BPD/FL ratio decreased slowly from 1.52 to 1.35, with a convergent range of 1.19–1.53 at week 36 (Table 2). Of 27 BPD/FL ratios obtained at each gestational week from 24 cases, none was in the control range (Fig. 1).

Discussion

In a normal pregnancy, the BPD/FL ratio shows a continuous absolute decrease as a function of gestational weeks, with a rapid decrease during weeks 12–18. The BPD/FL ratio decreases to less than 3 prior to gestational week 13, and to less than 2 prior to week 18. Therefore, we assume that by using the routine fetal biometry measurement, an extreme outlier such as TD can be easily detected from as early as the first trimester.

Three decades ago, the use of the BPD/FL ratio to detect trisomy 21 had been considered, but studies concluded that its usefulness was limited due to insignificant differences before 18 weeks [23,24]. The usefulness of measuring fetal nuchal

Table 1
The characteristics of fetal biometry and other clinical findings in thanatophoric dysplasia.

Case	Authors	Maternal age	Gestational week	BPD (mm)	FL (mm)	BPD/FL	Diagnosis	Molecular analyses	Radiological diagnosis
1	Sawai [8]	33	27	83.7	20.9	4.00	TD1	+	+
2	Chen [9]	28	27	82.0	19.8	4.14	TD1	+	+
3	Chen [9]	28	31	92.0	21.1	4.36	TD1	+	+
4	Chen [9]	30	18	45.4	12.9	3.52	TD1	+	+
5	Chen [9]	30	24	80.0	19.7	4.06	TD2	+	+
6	Hatzaki [10]	28	18	47.0	17.0	2.76	TD1	+	–
7	Hatzaki [10]	24	19	49.0	14.7	3.33	TD1	+	+
8	Hatzaki [10]	26	22	60.0	19.2	3.13	TD1	+	+
9	Giancotti [11]	37	12	25.3	6	4.22	TD1	+	–
10	Li [12]	34	21	54	24	2.25	TD2	+	+
11	Chen [13]	35	18	42.0	18.2	2.31	TD2	+	+
			21	51.5 ^a	16	3.22			
12	Cho [14]	30	15	35.7	11.6	3.08	TD1	+	–
			18	49.4	16.8	2.94			
13	Lindy [15]	35	12	26.0	6.1	4.26	TD1	+	+
14	Ngo [16]	35	14	35 ^a	10	3.50	TD1	+	+
15	Delahaye [17]	45	12	20 ^a	4.6	4.34	TD1	+	–
16	Tonni [18]	38	19	46.6	19.6	2.38	TD1	+	+
17	Li [19]	26	22	54	16	3.38	TD1	+	+
18	Li [19]	34	28	75	19	3.95	TD1	+	+
19	Li [19]	38	20	47	12	3.92	TD1	+	–
20	Schild [20]	23	26	72	36	2.00	TD2	–	+
21	Schild [20]	29	20	56	21	2.67	TD1	–	+
22	Schild [20]	29	20	60	18	3.33	TD1	–	+
23	Calongos [21]	36	13	27.4	8.2	3.34	TD1	–	+
			20	53	11.8	4.49			
24	Teele [22]	NA	18	44.8 ^a	18	2.48	TD1	–	+

AC, abdominal circumference; BPD, biparietal diameter; FL, femur length; NA, not available; TC, thoracic circumference; TD1, thanatophoric dysplasia type 1; TD2, thanatophoric dysplasia type 2.

^a Estimated value.

Table 2
The BPD/FL ratios of 1271 controls without skeletal dysplasia from gestational week 12 to 36.

Distribution	Gestational age (weeks)																		
	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30
100.0%	4.06	2.87	2.59	2.41	2.23	2.06	1.98	1.87	1.78	1.74	1.69	1.68	1.69	1.66	1.65	1.63	1.64	1.60	1.57
99.5%	4.06	2.87	2.59	2.41	2.23	2.06	1.98	1.86	1.78	1.74	1.69	1.68	1.69	1.65	1.64	1.63	1.61	1.60	1.56
97.5%	3.91	2.87	2.54	2.34	2.17	1.99	1.90	1.79	1.73	1.67	1.67	1.62	1.63	1.60	1.58	1.58	1.58	1.56	1.54
90.0%	3.54	2.67	2.35	2.20	2.04	1.91	1.81	1.73	1.66	1.62	1.60	1.57	1.56	1.55	1.53	1.54	1.52	1.51	1.49
75.0%	3.18	2.43	2.20	2.04	1.91	1.80	1.74	1.66	1.60	1.55	1.53	1.52	1.51	1.49	1.49	1.48	1.48	1.46	1.44
50.0%	2.82	2.26	2.05	1.89	1.78	1.70	1.65	1.57	1.52	1.49	1.48	1.45	1.45	1.44	1.42	1.43	1.42	1.41	1.40
25.0%	2.43	2.08	1.89	1.79	1.68	1.62	1.58	1.51	1.46	1.44	1.42	1.42	1.39	1.39	1.37	1.37	1.32	1.35	1.36
10.0%	2.07	1.93	1.75	1.69	1.59	1.53	1.49	1.44	1.40	1.37	1.37	1.35	1.34	1.34	1.33	1.33	1.31	1.31	1.30
2.5%	1.92	1.77	1.62	1.58	1.50	1.42	1.42	1.37	1.34	1.31	1.30	1.31	1.28	1.29	1.28	1.28	1.26	1.26	1.26
0.5%	1.89	1.68	1.55	1.46	1.39	1.39	1.37	1.30	1.28	1.26	1.25	1.30	1.24	1.26	1.26	1.22	1.26	1.21	1.25
0.0%	1.89	1.68	1.55	1.46	1.38	1.39	1.37	1.30	1.27	1.26	1.25	1.30	1.24	1.26	1.24	1.22	1.22	1.20	1.24
Mean	2.81	2.27	2.06	1.92	1.80	1.71	1.65	1.58	1.53	1.50	1.48	1.46	1.45	1.44	1.43	1.43	1.42	1.41	1.40
Std dev	0.51	0.26	0.22	0.19	0.17	0.14	0.12	0.11	0.10	0.09	0.09	0.08	0.09	0.08	0.08	0.08	0.07	0.08	0.07

translucency thickness and nasal bones at 10-weeks to detect trisomy 21 was subsequently reported [25,26], and the BPD/FL ratio in early gestational weeks is no longer being used to screen for Down syndrome. TD is subdivided into 2 types according to clinical features and mutations of the fibroblast growth factor receptor 3 (*FGFR3*) gene: type 1 (TD1) is characterized by curved femurs, with or without a cloverleaf skull, as a result of several mutations of *FGFR3*; most cases are caused by missense mutation of 742C → T [8,9]. Type 2 (TD2) is characterized by short straight femurs with a cloverleaf skull as a result of the mutation of 1948A → C [9]. These anomalies in TD possibly appear at the beginning of fetal skeletal development before 11 weeks, resulting in significant skeletal dysplasia, which can be observed at weeks 12–14 [11,16,27]. As a result, either macrocephaly or severe growth restriction of the femur will cause an increased denominator and decreased fraction of the BPD/FL ratio, with a higher ratio at an early stage of pregnancy. This can explain why the BPD/FL ratio may be useful in predicting TD since the first trimester.

Rahemtullah and Ramus et al. suggested that the fetal femur length-to-abdominal circumference ratio (FL/AC) can predict a lethal outcome, although Rahemtullah reported 2 cases of achondroplasia and Ramus reported 1 case of osteogenesis imperfecta type III with an FL/AC ratio less than 0.16, who survived after birth [28,29]. Parilla et al. also reported inaccuracy in 3 of 31 cases studied [6]. In addition, compared to Rosati's study of fetal biometry in normal pregnancy, FL/AC < 0.16 appears to be within the normal range before gestational week 16 [30] (Table 3). Therefore, the feasibility of using such a method to diagnose TD or other lethal disorders at all gestational ages remains questionable, despite Weaver et al. reporting that an FL/AC ratio of 0.124 can be a better clinical cutoff to predict lethality, based on a retrospective study of 23 cases at 21–38 weeks of gestation [31].

Ngo et al. reported that in approximately 69% of severe skeletal dysplasia cases, nuchal translucency (NT) can be seen in the first trimester [16], as well as in Down syndrome. Subsequently, several case studies reported the usefulness of this finding in skeletal disorders such as osteogenesis imperfecta type 1 [32], type 2 [33,34], and type 3 [35]. These studies highlighted the importance of closely monitoring skeletal development when increased NT is seen. However, in our experience, and based on previous reports, increased NT is not always seen in TD [11,36] or other lethal skeletal dysplasias [37]. In addition, a temporary increase in NT can be observed in some normal pregnancies [38]; therefore, the usefulness of NT to predict a rare TD is considered to be limited.

Despite our results suggesting that performing routine fetal biometry measurements since the first trimester can help detect TD, we do not recommend using this measurement alone to diagnose fetal lethality of an unknown skeletal dysplasia, without further confirmation. To avoid misdiagnosis, repeat sonographic evaluation, including that of the thorax, head, and heart, and another reliable examination such as cell-free fetal DNA in maternal plasma [7,39,40] or molecular analysis should be performed. Of note, normal BPD/FL ratio cannot deny different skeletal disorders caused by different mutations in the same *FGFR3* gene before the late second trimester, such as hypochondroplasia [41] or achondroplasia [42], as the shortened femur in these disorders is not obvious during this period [7].

TD is rare; however, according to our results, the BPD/FL ratio can probably be a useful marker to detect an extreme outlier such as TD since early pregnancy. If a higher BPD/FL ratio is obtained, fetal skeletal development should be monitored carefully.




Fig 1. The distribution of the BPD/FL ratio between the thanatophoric dysplasia and control groups. Of 27 BPD/FL ratios obtained from 24 cases, none was within the control range. AC, abdominal circumference; BPD, biparietal diameter; FL, femur length, TD, thanatophoric dysplasia.

Table 3
Fetal biometry measurement at gestational weeks 12–16 (modified from Rosati's study, 1997).

Week	BPD		AC		FL		FL/AC			BPD/FL		
	5%	95%	5%	95%	5%	95%	Minimum (5%/95%)	Average (50%/50%)	Maximum (95%/5%)	Minimum (5%/95%)	Average (50%/50%)	Maximum (95%/5%)
12	18.5	22.1	51.8	65.6	6.3	8.1	0.10	0.12	0.16	2.28	2.82	3.51
13	22.5	26.0	66.2	77.4	8.4	10.8	0.11	0.13	0.16	2.08	2.53	3.10
14	26.3	30.0	77.4	91.0	11.3	14.0	0.12	0.15	0.18	1.88	2.23	2.65
15	30.8	33.8	93.7	104.0	14.4	16.7	0.14	0.16	0.18	1.84	2.08	2.35
16	34.8	38.2	104.0	125.0	18.1	21.2	0.14	0.17	0.20	1.64	1.86	2.11

AC, abdominal circumference; BPD, biparietal diameter; FL, femur length.

Conflict of interest

The authors report no conflicts of interest.

Acknowledgment

The authors would like to thank Editage (www.editage.jp) for the English language review.

References

- [1] Orioli IM, Castilla EE, Barbosa-Neto JG. The birth prevalence rates for the skeletal dysplasias. *J Med Genet* 1986;23:328–32.
- [2] Kong CW, Tong LY, Lam WC, Chan LW, To WW. Learning curve in determining fetal sex by first trimester ultrasound scan. *J Med Ultrasound* 2016;24:54–9.
- [3] Shyu IL, Yang MJ, Wang HI, Wang PH, Chang CM, Juang CM, et al. Fetal maxillary and mandibular length in normal pregnancies from 11 weeks' to 13(+6) weeks' gestation: a Taiwanese study. *Taiwan J Obstet Gynecol* 2014;53:53–6.
- [4] Chang CP, Wang HI, Wang PH, Yang MJ, Chang CM, Juang CM, et al. Umbilical artery Doppler velocimetry in normal pregnancies from 11(+0) to 13(+6) gestational weeks: a Taiwanese study. *Taiwan J Obstet Gynecol* 2014;53:193–6.
- [5] Schramm T, Gloning KP, Minderer S, Daumer-Haas C, Hortnagel K, Nerlich A, et al. Prenatal sonographic diagnosis of skeletal dysplasias. *Ultrasound Obstet Gynecol* 2009;34:160–70.
- [6] Parilla BV, Leeth EA, Kambich MP, Chilis P, MacGregor SN. Antenatal detection of skeletal dysplasias. *J Ultrasound Med* 2003;22:255–8.
- [7] Chitty LS, Khalil A, Barrett AN, Pajkrt E, Griffin DR, Cole TJ. Safe, accurate, prenatal diagnosis of thanatophoric dysplasia using ultrasound and free fetal DNA. *Prenat Diagn* 2013;33:416–23.
- [8] Sawai H, Komori S, Ida A, Henmi T, Bessho T, Koyama K. Prenatal diagnosis of thanatophoric dysplasia by mutational analysis of the fibroblast growth factor receptor 3 gene and a proposed correction of previously published PCR results. *Prenat Diagn* 1999;19:21–4.
- [9] Chen CP, Chern SR, Shih JC, Wang W, Yeh LF, Chang TY, et al. Prenatal diagnosis and genetic analysis of type I and type II thanatophoric dysplasia. *Prenat Diagn* 2001;21:89–95.
- [10] Hatzaki A, Sifakis S, Apostolopoulou D, Bouzarelou D, Konstantinidou A, Kappou D, et al. FGFR3 related skeletal dysplasias diagnosed prenatally by ultrasonography and molecular analysis: presentation of 17 cases. *Am J Med Genet A* 2011;155A:2426–35.

- [11] Giancotti A, Castori M, Spagnuolo A, Binni F, D'Ambrosio V, Pasquali G, et al. Early ultrasound suspect of thanatophoric dysplasia followed by first trimester molecular diagnosis. *Am J Med Genet A* 2011;155A:1756–8.
- [12] Li D, Liao C, Ma X, Li Q, Tang X. Thanatophoric dysplasia type 2 with encephalocele during the second trimester. *Am J Med Genet A* 2006;140:1476–7.
- [13] Chen CP, Chang TY, Lin MH, Chern SR, Su JW, Wang W. Rapid detection of K650E mutation in FGFR3 using uncultured amniocytes in a pregnancy affected with fetal cloverleaf skull, occipital pseudoencephalocele, ventriculomegaly, straight short femurs, and thanatophoric dysplasia type II. *Taiwan J Obstet Gynecol* 2013;52:420–5.
- [14] Cho I, Shim JY, Kim GH, Yoo HW, Lee EJ, Won HS, et al. Thanatophoric dysplasia in a dichorionic twin confirmed by genetic analysis at the early second trimester: a case report and literature review. *Obstet Gynecol Sci* 2014;57:151–4.
- [15] Lindy AS, Basehore MJ, Munisha M, Williams AL, Friez MJ, Writzl K, et al. Identification of a novel insertion mutation in FGFR3 that causes thanatophoric dysplasia type 1. *Am J Med Genet A* 2016;170:1573–9.
- [16] Ngo C, Viot G, Aubry MC, Tsatsaris V, Grange G, Cabrol D, et al. First-trimester ultrasound diagnosis of skeletal dysplasia associated with increased nuchal translucency thickness. *Ultrasound Obstet Gynecol* 2007;30:221–6.
- [17] Delahaye S, Rosenblatt J, Costa JM, Bazin A, Benifla JL, Jouannic JM. First-trimester molecular prenatal diagnosis of a thanatophoric dysplasia. *Prenat Diagn* 2010;30:1222–3.
- [18] Tonni G, Azzoni D, Ventura A, Ferrari B, Felice CD, Baldi M. Thanatophoric dysplasia type I associated with increased nuchal translucency in the first trimester: early prenatal diagnosis using combined ultrasonography and molecular biology. *Fetal Pediatr Pathol* 2010;29:314–22.
- [19] Li D, Liao C, Ma X. Prenatal diagnosis and molecular analysis of type 1 thanatophoric dysplasia. *Int J Gynaecol Obstet* 2005;91:268–70.
- [20] Schild RL, Hunt GH, Moore J, Davies H, Horwell DH. Antenatal sonographic diagnosis of thanatophoric. *Ultrasound Obstet Gynecol* 1996;8:62–7.
- [21] Calongos G, Hori M, Ogino M, Sawai H. A case of thanatophoric dysplasia type I with fetal hydrops in the first trimester. *Case Rep Obstet Gynecol* 2016;2016:1821230.
- [22] Teele RL. A guide to the recognition of skeletal disorders in the fetus. *Pediatr Radiol* 2006;36:473–84.
- [23] Marquette GP, Boucher M, Desrochers M, Dallaire L. Screening for trisomy 21 with ultrasonographic determination of biparietal diameter/femur length ratio. *Am J Obstet Gynecol* 1990;163:1604–5.
- [24] Snijders RJ, Platt LD, Greene N, Carlson D, Krakow D, Gregory K, et al. Femur length and trisomy 21: impact of gestational age on screening efficiency. *Ultrasound Obstet Gynecol* 2000;16:142–5.
- [25] Snijders RJ, Noble P, Sebire N, Souka A, Nicolaides KH. UK multicentre project on assessment of risk of trisomy 21 by maternal age and fetal nuchal-translucency thickness at 10–14 weeks of gestation. *Lancet* 1998;352:343–6.
- [26] Dagklis T, Borenstein M, Peralta CF, Faro C, Nicolaides KH. Three-dimensional evaluation of mid-facial hypoplasia in fetuses with trisomy 21 at 11 + 0 to 13 + 6 weeks of gestation. *Ultrasound Obstet Gynecol* 2006;28:261–5.
- [27] Wong HS, Kidd A, Zuccollo J, Tuohy J, Strand L, Tait J, et al. A case of thanatophoric dysplasia: the early prenatal 2D and 3D sonographic findings and molecular confirmation of diagnosis. *Fetal Diagn Ther* 2008;24:71–3.
- [28] Rahemtullah A, McGillivray B, Wilson RD. Suspected skeletal dysplasias: femur length to abdominal circumference ratio can be used in ultrasonographic prediction of fetal outcome. *Am J Obstet Gynecol* 1997;177:864–9.
- [29] Ramus RM, Martin LB, Twickler DM. Ultrasonographic prediction of fetal outcome in suspected skeletal dysplasias with use of the femur length-to-abdominal circumference ratio. *Am J Obstet Gynecol* 1998;179:1348–52.
- [30] Rosati P, Guariglia L. Transvaginal fetal biometry in early pregnancy. *Early Hum Dev* 1997;49:91–6.
- [31] Weaver KN, Johnson J, Kline-Fath B, Zhang X, Lim FY, Tinkle B, et al. Predictive value of fetal lung volume in prenatally diagnosed skeletal dysplasia. *Prenat Diagn* 2014;34:1326–31.
- [32] Takahashi H, Horie K, Hayashi S, Matsubara S. Ultrasound findings of a fetus with osteogenesis imperfecta type I. *Arch Gynecol Obstet* 2016;294:213–4.
- [33] Hsieh CT, Yeh GP, Wu HH, Wu JL, Chou PH, Lin YM. Fetus with osteogenesis imperfecta presenting as increased nuchal translucency thickness in the first trimester. *J Clin Ultrasound* 2008;36:119–22.
- [34] Chen CP, Su YN, Chang TY, Huang MC, Pan CH, Chern SR, et al. Osteogenesis imperfecta type II: prenatal diagnosis and association with increased nuchal translucency and hypochogenicity of the cranium. *Taiwan J Obstet Gynecol* 2012;51:315–8.
- [35] Vimercati A, Panzarino M, Totaro I, Chincoli A, Selvaggi L. Increased nuchal translucency and short femur length as possible early signs of osteogenesis imperfecta type III. *J Prenat Med* 2013;7:5–8.
- [36] Beuke L, Koduri P, Martinez F, Weiner S. Stippling: a first trimester marker for thanatophoric dysplasia type I. *Prenat Diagn* 2010;30:1220–1.
- [37] Chen CP, Chen CY, Chern SR, Su JW, Wang W. First-trimester prenatal diagnosis of Ellis-van Creveld syndrome. *Taiwan J Obstet Gynecol* 2012;51:643–8.
- [38] Bilardo CM, Muller MA, Pajkrt E, Clur SA, van Zalen MM, Bijlsma EK. Increased nuchal translucency thickness and normal karyotype: time for parental reassurance. *Ultrasound Obstet Gynecol* 2007;30:11–8.
- [39] Chitty LS, Mason S, Barrett AN, McKay F, Lench N, Daley R, et al. Non-invasive prenatal diagnosis of achondroplasia and thanatophoric dysplasia: next-generation sequencing allows for a safer, more accurate, and comprehensive approach. *Prenat Diagn* 2015;35:656–62.
- [40] Cheng WL, Hsiao CH, Tseng HW, Lee TP. Noninvasive prenatal diagnosis. *Taiwan J Obstet Gynecol* 2015;54:343–9.
- [41] Chen CP, Su YN, Lin TH, Chang TY, Su JW, Wang W. Detection of a de novo Y278C mutation in FGFR3 in a pregnancy with severe fetal hypochondroplasia: prenatal diagnosis and literature review. *Taiwan J Obstet Gynecol* 2013;52:580–5.
- [42] Yang PY, Liao HG, Yeh GP, Hsieh CT. Prenatal diagnosis of achondroplasia with ultrasound, three-dimensional computed tomography and molecular methods. *J Med Ultrasound* 2012;20:176–9.