



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Research Letter

Pfeiffer syndrome with *FGFR2* C342R mutation presenting extreme proptosis, craniosynostosis, hearing loss, ventriculomegaly, broad great toes and thumbs, maxillary hypoplasia, and laryngomalaciaChih-Ping Chen^{a, b, c, d, e, f, *}, Shuan-Pei Lin^{b, g, h, i}, Yu-Peng Liu^{j, k}, Schu-Rern Chern^b, Shin-Wen Chen^a, Shih-Ting Lai^a, Wayseen Wang^{b, l}^a Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan^b Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan^c Department of Biotechnology, Asia University, Taichung, Taiwan^d School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan^e Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan^f Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan^g Department of Pediatrics, MacKay Memorial Hospital, Taipei, Taiwan^h Department of Medicine, MacKay Medical College, New Taipei City, Taiwanⁱ Department of Early Childhood Care, National Taipei University of Nursing and Health Sciences, Taipei, Taiwan^j Department of Radiology, Hsinchu MacKay Memorial Hospital, Hsinchu, Taiwan^k MacKay Medicine, Nursing and Management College, Taipei, Taiwan^l Department of Bioengineering, Tatung University, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 30 March 2017

Dear Editor,

The male proband was the first child of a healthy unrelated couple. The mother was 32 years old, and the father was 37 years old at his birth. There was no family history of congenital malformations. The pregnancy was uncomplicated and associated with unremarkable prenatal ultrasound findings. He was born at 39 weeks by cesarean section due to breech presentation. Birth weight was 2740 g, length was 49 cm, and head circumference was 33.5 cm. After birth, he was found to have severe proptosis, bilateral hearing loss, brachycephaly, hypertelorism, broad big toes and thumbs, low-set ears, and midface hypoplasia (Figs. 1 and 2). He had suffered from respiratory distress and cardiac arrest because of laryngomalacia that required a surgery to release and repair larynx. Radiographs showed multisynostoses of sagittal and coronal sutures, shallow orbits, brachycephaly and ventriculomegaly (Fig. 3). A diagnosis of type 3 Pfeiffer syndrome was made. Cytogenetic analysis revealed a karyotype of 46,XY. Molecular analysis of peripheral blood at age three months revealed a heterozygous

c.1024T>C, TGC>CGC transition, leading to a p.Cys342Arg (C342R) mutation in the *FGFR2* gene (Fig. 4).

We previously reported Pfeiffer syndrome with *FGFR2* mutation in two infants [1,2]. In this presentation, we additionally report a case of Pfeiffer syndrome with a *FGFR2* C342R mutation. Both Apert syndrome and Pfeiffer syndrome can be caused by *FGFR2* mutations. Apert syndrome is usually associated with *FGFR2* S252W and P253R mutations, and may manifest severe syndactyly of hands and feet but no cloverleaf skull or proptosis, whereas Pfeiffer syndrome is usually associated with *FGFR2* W290C, Y340C, C342R and S351C mutations, and can be associated with a cloverleaf skull, proptosis and broad great toes and thumbs [3,4]. The *FGFR2* mutations of W290C (p.Trp290Cys), Y340C (p.Tyr340Cys), C342R (p.Cys342Arg) and S351C (p.Ser351Cys) have been reported to be associated with severe phenotypic features of Pfeiffer syndrome such as a cloverleaf skull, extreme proptosis, midface hypoplasia, hydrocephalus, tracheal sleeve, Chiari malformation, radio-ulnar-humeral synostosis and early infant death [3,4]. In a review of 25 patients with *FGFR2* C342R mutation, Lajeunie et al [3] found 22 cases with Pfeiffer syndrome, two cases with Crouzon syndrome and one case with Jackson-Weiss syndrome. *FGFR2* C342R mutation has been occasionally observed in patients with Crouzon syndrome [5,6], Jackson-Weiss syndrome [7] and Antley-Bixler syndrome [8], but frequently in patients with severe Pfeiffer syndrome [3,9].

In a review of 18 cases with prenatal diagnosis of Pfeiffer syndrome, Giancotti et al [10] found frequent sonographic features of skull shape anomalies (72.2%), nasal abnormalities (50%), proptosis and hypertelorism (44.4%), frontal bossing (22.2%), thumbs' anomalies (33.3%) and toes' abnormalities (38.9%). However, our

* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei 10449, Taiwan. Fax: +886 2 25433642, +886 2 25232448.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).



Fig. 2. (A) Broad thumb and (B) broad big toe.

Fig. 1. (A) and (B). Craniofacial appearance of extreme proptosis, brachycephaly, midface hypoplasia, hypertelorism and low-set ears of the propositus at age 3 months.




Fig. 3. Radiographs show multisynostoses of (A) sagittal and (B) coronal sutures (arrows) and shallow orbits, and (C) ventriculomegaly.

case did not manifest identifiable abnormalities on prenatal ultrasound indicating that prenatal diagnosis of Pfeiffer syndrome simply by ultrasound screening is not easy.

In summary, we demonstrate a genotype-phenotype correlation of *FGFR2* C342R mutation with severe Pfeiffer syndrome. We suggest a mutational analysis of *FGFR2* is helpful in genetic counseling of newborns with severe Pfeiffer syndrome.

Conflict of interest

The authors have no conflicts of interest relevant to this article.




Fig. 4. Mutational analysis of the *FGFR2* gene shows a heterozygous c.1024T>C, TGC>CGC transition, leading to a p.Cys342Arg (C342R) mutation in the proband. WT = wild type.

Acknowledgements

This work was supported by research grants MOST-104-2314-B-195-009 and MOST-105-2314-B-195-012 from the Ministry of Science and Technology and MMH-E-105-04 from MacKay Memorial Hospital, Taipei, Taiwan.

References

- Chen C-P, Lin S-P, Su Y-N, Chien S-C, Tsai F-J, Wang W. Craniosynostosis and congenital tracheal anomalies in an infant with Pfeiffer syndrome carrying the W290C *FGFR2* mutation. *Genet Couns* 2008;19:165–72.
- Chen C-P, Huang H-K, Liu Y-P, Chern S-R, Su J-W, Wang W. Pfeiffer syndrome with *FGFR2* W290C mutation perinatally presenting extreme proptosis. *Taiwan J Obstet Gynecol* 2013;52:607–10.
- Lajeunie E, Heuertz S, El Ghouzzi V, Martinovic J, Renier D, Le Merrer M, et al. Mutation screening in patients with syndromic craniosynostoses indicates that a limited number of recurrent *FGFR2* mutations accounts for severe forms of Pfeiffer syndrome. *Eur J Hum Genet* 2006;14:289–98.
- Robin NH, Falk MJ, Haldeman-Englert CR. *FGFR*-related craniosynostosis syndromes. *GeneReviews*TM [Internet]. In: Pagon RA, Adam MP, Ardinger HH, Wallace SE, Amemiya A, Bean LJH, et al, editors. Seattle, WA: University of Washington; 1993–2016. [Updated 2011 Jun 7].
- Reardon W, Winter RM, Rutland P, Pulleyn LJ, Jones BM, Malcolm S. Mutations in the fibroblast growth factor receptor 2 gene cause Crouzon syndrome. *Nat Genet* 1994;8:98–103.
- Ke R, Yang X, Tianyi C, Ge M, Lei J, Mu X. The C342R mutation in *FGFR2* causes Crouzon syndrome with elbow deformity. *J Craniofac Surg* 2015;26:584–6.
- Park W-J, Meyers GA, Li X, Theda C, Day D, Orlow SJ, et al. Novel *FGFR2* mutations in Crouzon and Jackson-Weiss syndromes show allelic heterogeneity and phenotypic variability. *Hum Mol Genet* 1995;4:1229–33.
- Reardon W, Smith A, Honour JW, Hindmarsh P, Das D, Rumsby G, et al. Evidence for digenic inheritance in some cases of Antley-Bixler syndrome? *J Med Genet* 2000;37:26–32.
- Rutland P, Pulleyn LJ, Reardon W, Baraitser M, Hayward R, Jones B, et al. Identical mutations in the *FGFR2* gene cause both Pfeiffer and Crouzon syndrome phenotypes. *Nat Genet* 1995;9:173–6.
- Giancotti A, D'Ambrosio V, Marchionni E, Squarcella A, Aliberti C, La Torre R, et al. Pfeiffer syndrome: literature review of prenatal sonographic findings and genetic diagnosis. *J Matern Fetal Neonatal Med* 2016. <http://dx.doi.org/10.1080/14767058.2016.1243099>.