

Contents lists available at [ScienceDirect](http://www.sciencedirect.com)

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Original Article

Serum vitamin D concentrations in young Turkish women with primary dysmenorrhea: A randomized controlled study

Osman Karacin ^a, Ilknur Mutlu ^b, Mesut Kose ^a, Fatih Celik ^a, Mine Kanat-Pektas ^{a,*}, Mehmet Yilmazer ^a^a Department of Obstetrics and Gynecology, Afyon Kocatepe University Hospital, Afyonkarahisar, Turkey^b Department of Obstetrics and Gynecology, Novaart In Vitro Fertilization Center, Ankara, Turkey

ARTICLE INFO

Article history:

Accepted 14 December 2016

Keywords:

Dairy product
Dysmenorrhea
Parathyroid hormone
Visual analogue scale
Vitamin D

ABSTRACT

Objective: This study aims to investigate the possible role of vitamin D deficiency in primary dysmenorrhea by assessing serum 25-hydroxyvitamin D₃ levels in a cohort which includes young Turkish women with primary dysmenorrhea and healthy controls.**Materials and methods:** A total of 683 women who were aged between 18 and 25 years and who were consecutively admitted to the study center were eligible. After the exclusion of 55 women, 184 women with primary dysmenorrhea were randomly assigned into the dysmenorrhea group and 184 women without dysmenorrhea were randomly allocated into the control group.**Results:** The dysmenorrhea group had significantly less consumption of dairy products ($p = 0.001$), lower serum calcium ($p = 0.001$), lower serum vitamin D ($p = 0.001$) and higher serum parathyroid hormone ($p = 0.001$) than those of the control group. Hyperparathyroidism was significantly less frequent whereas vitamin D deficiency was significantly more frequent in the dysmenorrhea group ($p = 0.001$ for each). The dysmenorrhea patients with vitamin D deficiency had significantly higher visual analogue scale (VAS) scores ($p = 0.001$). Depression, irritability, mood swings, fatigue, headache and breast tenderness were significantly more frequent in the vitamin D deficiency group ($p < 0.05$ for all). The VAS scores of the dysmenorrhea patients correlated positively and significantly with serum parathyroid hormone levels ($r = 0.666$, $p = 0.001$) whereas these VAS scores correlated negatively and significantly with serum vitamin D levels ($r = -0.713$, $p = 0.001$).**Discussion:** The significant and positive correlation between vitamin D levels and VAS scores and the significant reduction in serum vitamin D levels of the dysmenorrhea patients designate the possible role of vitamin D deficiency in the primary dysmenorrhea.© 2018 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Primary dysmenorrhea is characterized by uterine cramping which causes suprapubic pain that occurs just before or during menstruation in the absence of any pelvic pathologic conditions. Primary dysmenorrhea is observed in at least half of the menstruating women, often resulting in the interruption of school and work schedules and ultimately leading to educational and economical considerations. Although the pathogenesis of primary dysmenorrhea has not been clarified, it has been proposed that an

excessive release of prostaglandins triggers uterine contractions and lower abdominal pain [1,2].

Previously published studies have identified younger age, younger age at menarche, nulliparity, higher and longer menstrual flow, smoking and positive family history as the associated risk factors. These studies have also indicated the possible role of certain nutrients and dietary habits in the etiopathogenesis of dysmenorrhea. For instance, dysmenorrhea is related with irregular eating, obesity and history of an attempt to lose weight. Moreover, it has been found that the intake of caffeine and sugar is increased and the consumption of vegetables and fruits is reduced in dysmenorrhea patients (Table 1) [3–17]. Recently, two Jordanian studies have reported about the relationship between low calcium intake or vitamin D insufficiency and dysmenorrhea in adolescents and young women with dysmenorrhea [18,19]. Such a relationship

* Corresponding author. Selcuklu Mah. Adnan Kahveci Cad. No: 16/2, D: 4, Afyonkarahisar, Turkey. Fax. +90 2722463322.

E-mail address: minekanat@hotmail.com (M. Kanat-Pektas).

Table 1
Previously published studies on the etiopathogenesis of primary dysmenorrhea.

| Authors | Year | Country | Population | Risk factors |
|--------------------------|------|----------------------|------------------|--|
| Hailemeskel S [3] | 2016 | Ethiopia | 440 women | Nulliparity Positive family history Lower monthly income History of depression/anxiety History of attempt to lose weight Drinking > 4 glasses of tea/day Drinking > 1 bottle of coke/day |
| Tomás-Rodríguez MI [4] | 2016 | Spain | 306 women | Higher menstrual flow |
| Pejčić A, Janković S [5] | 2016 | Serbia | 288 women | Younger age at menarche Positive family history Longer menstrual flow |
| Ju H [6] | 2016 | Australia | 9067 women | Smoking |
| Habibi N [7] | 2015 | Iran | 311 women | Smoking Younger age Positive family history Higher menstrual flow Shorter intermenstrual period |
| Kural M [8] | 2015 | India | 310 adolescents | Residing at home Longer menstrual flow Higher menstrual flow |
| Kazama M [9] | 2015 | Japan | 1167 adolescents | Positive family history Less physical activity Less sleep (< 6 h/day) |
| Shiferaw MT [10] | 2014 | Ethiopia | 470 women | Longer menstrual flow Positive family history |
| Sahin S [11] | 2014 | Turkey | 520 women | Circumcision Menstrual irregularity Smoking |
| Beal SJ [12] | 2014 | USA | 262 adolescents | Positive family history History of depression/anxiety |
| Ju H [13] | 2014 | Review of 15 studies | | Younger age/Nulliparity Smoking/Substance abuse History of attempt to lose weight Positive family history History of depression/anxiety |
| Jang IA [14] | 2013 | Vietnam | 3017 women | Younger age Nulliparity Younger age at menarche Higher menstrual flow |
| Grandi G [15] | 2012 | Italy | 408 women | Younger menstrual flow Younger age at menarche Longer menstrual flow |
| Gagua T [16] | 2012 | Georgia | 2561 women | Smoking Smoking Positive family history Eating more sugar Skipping meals |
| Tavallaee M [17] | 2011 | Iran | 381 women | Younger age Positive family history History of depression/anxiety Eating less vegetables and fruits |

has been attributed to the physiological effects of calcium on muscle contractility and relaxation [20,21]. Since calcium homeostasis is maintained by the conjoint actions of calcitonin, parathyroid hormone and 25-hydroxyvitamin D₃, it can be expected that these three hormones would also participate in the pathophysiology of primary dysmenorrhea [22,23].

The present study aims to investigate the possible role of vitamin D₃ deficiency in primary dysmenorrhea by assessing serum 25-hydroxyvitamin D₃ levels in a cohort which includes young Turkish women with primary dysmenorrhea and healthy controls.

Material and methods

This prospective, randomized, case-controlled study was approved by the Institutional Review Board and Ethical Committee of Afyon Kocatepe University where it was undertaken from January 2015 to January 2016. Written informed consent was obtained from all participants included in the study.

Study design and patients

A total of 423 women who were aged between 18 and 25 years and who were consecutively admitted to the department of gynecology at the study center were eligible for the study. Twenty-two women who had pelvic pathologies, twelve women who had systemic diseases, ten women who had acute infections, six women who regularly used calcium or vitamin D supplements and five women who refused to participate were excluded. A total of 184 women who had regular menstrual cycles (occurring in 21–35 days, with menstruation lasting 3–7 days) and who experienced at least 4 consecutive painful periods in the past 6 months with the pain starting one day before or on the day of onset of bleeding were randomly assigned into the dysmenorrhea group. Another total of 184 women without dysmenorrhea were randomly allocated into the control group. Randomization was performed by sequentially numbered, sealed, opaque envelopes (Refer Fig. 1).




Fig. 1. Flow consort chart of the study.

The participants were instructed to complete a guided self-assessment questionnaire including their demographic features and clinical characteristics about menstruation. Body mass index was calculated as follows: $\text{Body mass index (BMI)} = \text{Body weight (kg)} / \text{Body height}^2 (\text{m}^2)$.

The participants also answered questions regarding their consumption of dairy products. The intake of dairy products was determined on daily basis as less than 1, 1 to 2 and more than 2 servings per day. A dairy serving was defined as: 1 cup of milk or yogurt, 2 tablespoons of butter, or 2 tablespoons (or $\frac{1}{4}$ cup) of cheese.

Moreover, the participants were requested to answer questions about their recurrent experience of 12 physical and psychological symptoms emerging during the premenstrual period. Psychological symptoms of depression, irritability, mood swings, social withdrawal, change in appetite and cravings for sweet or salty foods are recorded, along with physical symptoms such as general fatigue, headaches, nausea, abdominal bloating, and breast tenderness. The severity of dysmenorrhea pain was assessed by using visual analogue scale (VAS) scoring system which is based on numerical rating between 1 and 10.

Laboratory studies

Venous blood samples of 10 ml were collected into heparinized tubes. Serum concentrations of 25-hydroxyvitamin D₃ and parathyroid hormone were measured by a chemiluminescent assay (Eleclys 2010 analyzer, Roche Diagnostics, Mannheim, Germany). The intra-assay and inter-assay coefficients of variation (CVs) for vitamin D were 1.3% and 1.8% respectively while the intra-assay and inter-assay CVs for parathormone were 1.7% and 1.5% respectively. Serum 25-hydroxyvitamin D₃ levels <12 ng/ml indicated vitamin D

deficiency and serum parathyroid hormone concentrations >54 pg/ml referred to hyperparathyroidism.

Statistical analysis

Collected data were analyzed by Statistical Package for Social Sciences version 18.0 (SPSS-IBM Inc., Armonk, NY, USA). Continuous variables were expressed as mean \pm standard deviation (range: minimum–maximum) whereas categorical variables were denoted as numbers or percentages. Student t-test, chi-square test and Mann Whitney U-test were used for the comparisons. Spearman correlation test was used to detect the correlations among the variables. Two-tailed p values less than 0.05 were accepted to be statistically significant. A *post hoc* analysis was carried out to make a retrospective power analysis and it was determined that a cohort size of 368 women (184 healthy controls and 184 patients with primary dysmenorrhea) had 84.6% power to detect a difference at the 0.05 significance level.

Results

Table 2 compares the demographic, clinical and biochemical characteristics of the dysmenorrhea and control groups. When compared with the healthy controls, the dysmenorrhea patients had significantly more menstrual bleeding ($p = 0.004$), less consumption of dairy products ($p = 0.001$), lower calcium ($p = 0.001$), lower serum 25-hydroxyvitamin D₃ ($p = 0.001$), higher serum parathyroid hormone ($p = 0.001$) and higher VAS scores ($p = 0.001$). Positive family history, vitamin D deficiency and hyperparathyroidism were significantly more frequent in the dysmenorrhea group ($p = 0.001$ for each). Table 3 demonstrates that depression, irritability, mood swings, fatigue, headache, breast

Table 2
Demographic, clinical and biochemical characteristics of the participants.

| | Dysmenorrhea patients (n = 184) | Healthy controls (n = 184) | p |
|--------------------------------------|---------------------------------|----------------------------|------------------------------|
| Age (years) | 20.8 ± 1.9 | 20.8 ± 2.0 | 0.957 |
| Weight (kg) | 53.8 ± 5.0 | 53.9 ± 5.4 | 0.904 |
| Height (m) | 1.55 ± 0.05 | 1.55 ± 0.10 | 0.793 |
| Body mass index (kg/m ²) | 22.46 ± 2.75 | 22.06 ± 2.43 | 0.140 |
| Education | | | 0.587 ($\chi^2 = 1067$) |
| Primary | 50 (27.2%) | 59 (32.1%) | |
| Secondary | 34 (18.5%) | 31 (16.8%) | |
| High | 100 (54.3%) | 94 (51.1%) | |
| Employment | | | 0.119 ($\chi^2 = 4262$) |
| Unemployed | 173 (94.0%) | 164 (89.1%) | |
| Employed | 11 (6.0%) | 20 (10.9%) | |
| Menarche (years) | 12.2 ± 1.3 | 12.1 ± 1.2 | 0.366 |
| Menstrual cycle length (days) | 27.3 ± 2.1 | 27.1 ± 2.3 | 0.705 |
| Menstrual cycle duration (days) | 5.2 ± 0.9 | 5.0 ± 1.0 | 0.442 |
| Menstrual bleeding (pads/day) | 6.2 ± 1.1 | 5.9 ± 1.2 | 0.004* |
| Positive family history | 115 (62.5%) | 53 (28.8%) | 0.001* ($\chi^2 = 42,101$) |
| Consumption of dairy products | | | 0.001* ($\chi^2 = 27,129$) |
| <1 serving/day | 139 (75.5%) | 91 (49.5%) | |
| 1-2 servings/day | 34 (18.5%) | 75 (40.8%) | |
| >2 servings/day | 11 (6.0%) | 18 (9.8%) | |
| Serum calcium (mg/dl) | 8.3 ± 0.7 | 8.8 ± 0.9 | 0.001* |
| Serum magnesium (mg/dl) | 2.1 ± 0.4 | 2.2 ± 0.5 | 0.910 |
| Alkaline phosphatase (IU/l) | 80.4 ± 24.8 | 78.8 ± 23.7 | 0.492 |
| 25-hydroxyvitamin D (ng/ml) | 7.1 ± 3.8 | 14.9 ± 2.5 | 0.001* |
| 25-hydroxyvitamin D deficiency | 155 (84.2%) | 22 (12.0%) | 0.001* |
| Parathyroid hormone (pg/ml) | 66.4 ± 19.5 | 49.1 ± 13.0 | 0.001* |
| Hyperparathyroidism | 80 (43.5%) | 22 (12.0%) | 0.001* ($\chi^2 = 45,627$) |
| Visual analog scale score | 7.3 ± 1.4 | 2.2 ± 0.9 | 0.001* |

*p < 0.05 was accepted to be statistically significant.

Table 3
Clinical symptoms of the participants.

| | Dysmenorrhea patients (n = 184) | Healthy controls (n = 184) | p |
|--------------------|---------------------------------|----------------------------|------------------------------|
| Depression | 93 (50.5%) | 35 (19.0%) | 0.001* ($\chi^2 = 40,298$) |
| Irritability | 90 (48.9%) | 30 (16.3%) | 0.001* ($\chi^2 = 44,516$) |
| Mood swings | 86 (46.7%) | 53 (28.8%) | 0.001* ($\chi^2 = 12,590$) |
| General fatigue | 86 (46.7%) | 44 (23.9%) | 0.001* ($\chi^2 = 20,981$) |
| Headache | 82 (44.6%) | 57 (31.0%) | 0.007* ($\chi^2 = 7226$) |
| Nausea | 44 (23.9%) | 52 (28.3%) | 0.342 ($\chi^2 = 0,902$) |
| Social withdrawal | 35 (19.0%) | 50 (27.2%) | 0.064 ($\chi^2 = 3442$) |
| Abdominal bloating | 30 (16.3%) | 37 (20.1%) | 0.344 ($\chi^2 = 0,894$) |
| Breast tenderness | 26 (14.1%) | 11 (6.0%) | 0.009* ($\chi^2 = 6761$) |
| Change in appetite | 23 (12.5%) | 11 (6.0%) | 0.031* ($\chi^2 = 4666$) |
| Food craving | 22 (12.0%) | 12 (6.5%) | 0.072 ($\chi^2 = 3241$) |
| Anxiety | 14 (7.6%) | 11 (6.0%) | 0.534 ($\chi^2 = 0,386$) |

*p < 0.05 was accepted to be statistically significant.

tenderness and appetite change were significantly more frequent in the dysmenorrhea group (p = 0.001, p = 0.001, p = 0.001, p = 0.001, p = 0.007, p = 0.009 and p = 0.031 respectively).

Table 4 shows that dysmenorrhea patients with 25-hydroxyvitamin D₃ deficiency had significantly shorter height, longer menstrual cycle, longer menstrual bleeding, less menstrual bleeding, less consumption of dairy products and higher VAS scores (respectively p = 0.014, p = 0.002, p = 0.001, p = 0.001, p = 0.001 and p = 0.001). The rates of low education and unemployment were also significantly more frequent in this group (p = 0.001 for both).

Depression, irritability, mood swings, fatigue, headache and breast tenderness were significantly more frequent in the 25-hydroxyvitamin D₃ deficiency group (p < 0.05 for all). However, nausea, social withdrawal, change in appetite and food craving were significantly less frequent in the 25-hydroxyvitamin D₃ deficiency group (p = 0.001 for each) (Table 5).

The VAS scores of dysmenorrhea patients correlated positively and significantly with their BMI (r = 0.242, p = 0.001), menstrual

cycle length (r = 0.305, p = 0.001), menstrual bleeding duration (r = 0.271, p = 0.001) and parathyroid hormone levels (r = 0.666, p = 0.001). The VAS scores of dysmenorrhea patients correlated negatively and significantly with their menarche age (r = -0.246, p = 0.001) and serum 25-hydroxyvitamin D₃ levels (r = -0.713, p = 0.001).

Discussion

Vitamin D is a biologically inert molecule which is activated by hydroxylation, first to 25-hydroxyvitamin D₃ by 25 α -hydroxylase in the liver and then to 1,25-dihydroxyvitamin D₃ by 1 α -hydroxylase in the kidney. This activation process induces the intestinal absorption of calcium and phosphate. When serum vitamin D level decreases, intestinal calcium absorption would be reduced significantly. Then, calcium in the extracellular fluid would decrease and parathyroid hormone release would increase. In turn, parathyroid hormone would enhance the renal reabsorption of calcium and intestinal absorption of calcium and phosphate [23–26].

Risk factors that are associated with dysmenorrhea include younger age, younger age at menarche, nulliparity, higher and longer menstrual flow, smoking and positive family history. The findings of the present study also designate higher menstrual flow and positive family history as the underlying risk factors for dysmenorrhea [3–17]. The possible relationship between dietary habits and dysmenorrhea as well as the widespread location of vitamin D receptors throughout the human body and the expression of 1 α -hydroxylase in decidual cells imply that vitamin D may participate in the pathogenesis of dysmenorrhea [27,28].

This study indicates the overall prevalence of vitamin D deficiency (<12 ng/ml) as 44.5% in a cohort of 398 young Turkish women. This number is compatible with those previously published in literature. The prevalence of vitamin D deficiency was found to be 50.4% in a cohort of 258 healthy Turkish women [29]. Later, Ergur et al. detected moderate vitamin D deficiency in 54.3%

Table 4
Demographic, clinical and biochemical characteristics of dysmenorrhea patients.

| | 25-hydroxyvitamin D ₃ deficiency (n = 155) | 25-hydroxyvitamin D ₃ normal (n = 29) | p |
|--------------------------------------|---|--|-----------------------------------|
| Age (years) | 20.8 ± 1.8 | 21.0 ± 2.3 | 0.543 |
| Weight (kg) | 53.6 ± 5.2 | 55.2 ± 4.2 | 0.112 |
| Height (m) | 1.55 ± 0.05 | 1.57 ± 0.05 | 0.014* |
| Body mass index (kg/m ²) | 22.48 ± 2.90 | 22.36 ± 1.68 | 0.747 |
| Education | | | 0.001* (χ ² = 24,846) |
| Primary | 50 (32.3%) | 0 (0.0%) | |
| Secondary | 33 (21.2%) | 1 (3.4%) | |
| High | 72 (46.5%) | 28 (96.6%) | |
| Employment | | | 0.001* (χ ² = 123,933) |
| Unemployed | 144 (92.9%) | 0 (0.0%) | |
| Employed | 11 (7.1%) | 29 (100.0%) | |
| Menarche (years) | 12.2 ± 1.3 | 12.6 ± 0.9 | 0.070 |
| Menstrual cycle length (days) | 27.5 ± 2.1 | 26.2 ± 1.6 | 0.002* |
| Menstrual cycle duration (days) | 5.2 ± 0.9 | 4.7 ± 0.7 | 0.001* |
| Menstrual bleeding (pads/day) | 5.7 ± 1.2 | 6.7 ± 1.2 | 0.001* |
| Family history of dysmenorrhea | 98 (63.2%) | 17 (58.6%) | 0.638 (χ ² = 0,221) |
| Consumption of dairy products | | | 0.001* (χ ² = 98,757) |
| <1 serving/day | 136 (87.7%) | 3 (10.3%) | |
| 1-2 servings/day | 19 (12.3%) | 15 (51.7%) | |
| >2 servings/day | 0 (0.0%) | 11 (37.9%) | 0.001* |
| Serum calcium (mg/dl) | 8.3 ± 0.7 | 8.4 ± 0.8 | 0.455 |
| Serum magnesium (mg/dl) | 2.0 ± 0.3 | 2.2 ± 0.4 | 0.070 |
| Alkaline phosphatase (IU/l) | 81.4 ± 25.1 | 76.2 ± 23.5 | 0.271 |
| Parathyroid hormone (pg/ml) | 70.5 ± 9.0 | 47.2 ± 4.4 | 0.001* |
| Hyperparathyroidism | 74 (47.7%) | 6 (20.7%) | 0.007* (χ ² = 7275) |
| Visual analogue scale score | 7.6 ± 1.2 | 5.3 ± 0.5 | 0.001* |

*p < 0.05 was accepted to be statistically significant.

of women with singleton term pregnancies and 45.2% of non-pregnant fertile women [30]. In a similar study, the prevalence of severe vitamin D deficiency was 45.9% in a cohort of 229 women with singleton first-trimester pregnancies [31].

The prevalence of vitamin D deficiency and hyperparathyroidism is 84.2% and 43.5% respectively in this cohort of 184 Turkish women with dysmenorrhea. In a study by Abdul-Razzak et al., the prevalence of vitamin D deficiency (<10 ng/ml), vitamin D insufficiency and hyperparathyroidism was 9%, 80% and 48% respectively among 56 young Jordanian women with severe dysmenorrhea [19]. However, vitamin D insufficiency and hyperparathyroidism were observed in respectively 56.5% and 27.6% of young Jordanian women with dysmenorrhea [32]. These contradictory results may be due to the differences in the cut-off values and the timing of blood sampling for parathyroid hormone measurements.

This study indicates that 75.5% of the dysmenorrhea patients consumed <1 serving of dairy products per day and only 6% of these patients consumed >2 servings of dairy products per day. In a study, only 36% of the students who consumed 3 or 4 servings of dairy products per day had severe dysmenorrhea whereas 97% of the students who did not ingest any dairy products complained of severe dysmenorrhea [19]. In a smaller cohort, nearly 50% of the young women with dysmenorrhea had <1 serving of dairy products per day [18]. Vitamin D insufficiency or deficiency has been linked to low socioeconomic status in prior studies [33–35]. Similarly,

Table 5
Clinical Symptoms of Dysmenorrhea Patients with respect to Vitamin D Status.

| | 25-hydroxyvitamin D ₃ deficiency (n = 155) | 25-hydroxyvitamin D ₃ normal (n = 29) | p |
|--------------------|---|--|----------------------------------|
| Depression | 93 (60.0%) | 0 (0.0%) | 0.001* (χ ² = 35,182) |
| Irritability | 90 (58.1%) | 0 (0.0%) | 0.001* (χ ² = 33,416) |
| Mood swings | 86 (55.5%) | 0 (0.0%) | 0.001* (χ ² = 30,210) |
| General fatigue | 86 (55.5%) | 0 (0.0%) | 0.001* (χ ² = 30,210) |
| Headache | 82 (52.9%) | 0 (0.0%) | 0.001* (χ ² = 27,676) |
| Nausea | 25 (16.1%) | 19 (65.5%) | 0.001* (χ ² = 32,750) |
| Social withdrawal | 23 (14.8%) | 12 (41.4%) | 0.001* (χ ² = 11,172) |
| Abdominal bloating | 23 (14.8%) | 7 (24.1%) | 0.213 (χ ² = 1548) |
| Breast tenderness | 26 (16.8%) | 0 (0.0%) | 0.017* (χ ² = 5665) |
| Food craving | 11 (7.1%) | 11 (37.9%) | 0.001* (χ ² = 22,064) |
| Change in appetite | 10 (6.5%) | 13 (44.8%) | 0.001* (χ ² = 24,216) |
| Anxiety | 14 (9.0%) | 0 (0.0%) | 0.092 (χ ² = 2835) |

*p < 0.05 was accepted to be statistically significant.

dysmenorrhea patients with vitamin D deficiency have a low education rate of 53.5% and an unemployment rate of 93% in this study.

The present study points out depression (50.5%), irritability (48.9%), mood swings (46.7%), fatigue (46.7%) and headache (44.6%) as the most frequent premenstrual symptoms. In a Jordanian study, the most commonly encountered premenstrual symptoms were fatigue (72.9%), mood swings (72.3%), anxiety (68.9%), abdominal bloating (68.9%) and depression (58.8%) [32]. Calcium has been addressed as a micronutrient which is directly associated with the severity of premenstrual symptoms. As vitamin D and parathyroid hormone regulate calcium homeostasis, it can be hypothesized that they may also have a role in promoting premenstrual symptoms [36,37]. Accordingly, Bertone-Johnson et al. has related low dietary intake of vitamin D with the emergence of premenstrual symptoms [38]. In addition, it has been demonstrated that high calcium intake or vitamin D supplementation contributes to the alleviation of premenstrual symptoms [39–43]. However, Obeidat et al. were unable to show a significant difference between vitamin D levels of women with and without premenstrual symptoms. They also failed to detect any significant relationship between high dietary calcium intake and premenstrual symptoms except headache and social withdrawal [32].

As for the present study, depression, irritability, mood swings, fatigue, headache and breast tenderness are significantly more frequent while nausea, social withdrawal, change in appetite and food craving are significantly less frequent in the vitamin D deficiency group. Such discrepancies may be attributed to the differences in demographic and clinical characteristics of the reviewed participants and seasonal variations in serum concentrations of vitamin D. The possible correlation between vitamin D deficiency and premenstrual symptoms suggest that vitamin D participates in the etiopathogenesis of these symptoms and vitamin D can be used to treat these symptoms. As known, nearly every tissue and cell type in the body (especially myocardial cells, neurons and adipocytes) has receptors for vitamin D meaning that they all require vitamin D for adequate functioning. Additionally, vitamin D regulates genes that control cell growth and development, immune function, and metabolic control. Any interference with vitamin D metabolism may lead to the psychological, neurological and gastrointestinal symptoms that accompany dysmenorrhea [23–26].

To the best of our knowledge, this is the first randomized controlled study which aims to assess serum vitamin D levels of

young women with primary dysmenorrhea and healthy controls. The significant and positive correlation between vitamin D levels and VAS scores and the significant reduction in serum vitamin D levels of the dysmenorrhea patients designate the possible role of vitamin D deficiency in primary dysmenorrhea. However, the power of this study is limited by two factors. First, this study was undertaken throughout a year and seasonal variations in serum vitamin D concentrations could not be taken into consideration. Second, the patients with known pelvic pathologies and systemic diseases were excluded but clinical entities which could cause pelvic pain and have an accurate diagnosis by interventional methods such as laparoscopy could not be ruled out. Further research is warranted to clarify the role of vitamin D in the pathogenesis of primary dysmenorrhea.

Funding

This research was funded by Scientific Research Projects Board of Afyon Kocatepe University Medical Faculty (grant no: 15. HIZ.DES.73).

Conflict of interest statement

The authors declare that they have no conflict of interest.

References

- [1] De Sanctis V, Soliman A, Bernasconi S, Bianchini L, Bona G, Bozzola M, et al. Primary dysmenorrhea in adolescents: prevalence, impact and recent knowledge. *Pediatr Endocrinol Rev* 2015;13(2):512–20.
- [2] Harel Z. Dysmenorrhea in adolescents and young adults: an update on pharmacological treatments and management strategies. *Expert Opin Pharmacother* 2012;13(15):2157–70.
- [3] Hailemeskel S, Demissie A, Assefa N. Primary dysmenorrhea magnitude, associated risk factors, and its effect on academic performance: evidence from female university students in Ethiopia. *Int J Womens Health* 2016 Sep 19;8:489–96.
- [4] Tomás-Rodríguez MI, Palazón-Bru A, Martínez-St John DR, Navarro-Cremades F, Toledo-Marhuenda JV, Gil-Guillén VF. Factors associated with increased pain in primary dysmenorrhoea: analysis through a multivariate ordered logistic regression model. *J Pediatr Adolesc Gynecol* 2016 Sep 29;30(2):199–202.
- [5] Pejčić A, Janković S. Risk factors for dysmenorrhea among young adult female university students. *Ann Inst Soc Sci* 2016;52(1):98–103.
- [6] Habibi N, Huang MS, Gan WY, Zulida R, Safavi SM. Prevalence of primary dysmenorrhea and factors associated with its intensity among undergraduate students: a cross-sectional study. *Pain Manag Nurs* 2015;16(6):855–61.
- [7] Kural M, Noor NN, Pandit D, Joshi T, Patil A. Menstrual characteristics and prevalence of dysmenorrhea in college going girls. *J Family Med Prim Care* 2015;4(3):426–31.
- [8] Kazama M, Maruyama K, Nakamura K. Prevalence of dysmenorrhea and its correlating lifestyle factors in Japanese female junior high school students. *Tohoku J Exp Med* 2015;236(2):107–13.
- [9] Ju H, Jones M, Mishra GD. Smoking and trajectories of dysmenorrhoea among young Australian women. *Tob Control* 2016;25(2):195–202.
- [10] Shiferaw MT, Wubshet M, Tegabu D. Menstrual problems and associated factors among students of bahir dar university, Amhara National regional state, Ethiopia: a cross-sectional survey. *Pan Afr Med J* 2014 Apr 1;17:246.
- [11] Sahin S, Ozdemir K, Unsal A, Arslan R. Review of frequency of dysmenorrhea and some associated factors and evaluation of the relationship between dysmenorrhea and sleep quality in university students. *Gynecol Obstet Invest* 2014;78(3):179–85.
- [12] Beal SJ, Dorn LD, Sucharew HJ, Sontag-Padilla L, Pabst S, Hillman J. Characterizing the longitudinal relations between depressive and menstrual symptoms in adolescent girls. *Psychosom Med* 2014;76(7):547–54.
- [13] Jang IA, Kim MY, Lee SR, Jeong KA, Chung HW. Factors related to dysmenorrhea among Vietnamese and Vietnamese marriage immigrant women in South Korea. *Obstet Gynecol Surv* 2013;56(4):242–8.
- [14] Ju H, Jones M, Mishra G. The prevalence and risk factors of dysmenorrhoea. *Epidemiol Rev* 2014;36:104–13.
- [15] Grandi G, Ferrari S, Xholli A, Cannoletta M, Palma F, Romani C, et al. Prevalence of menstrual pain in young women: what is dysmenorrhoea? *J Pain Res* 2012;5:169–74.
- [16] Gagaa T, Ktshelashvili B, Gagaa D. Primary dysmenorrhea: prevalence in adolescent population of Tbilisi, Georgia and risk factors. *J Turk Ger Gynecol Assoc* 2012;13(3):162–8.
- [17] Tavallaei M, Joffres MR, Corber SJ, Bayanzadeh M, Rad MM. The prevalence of menstrual pain and associated risk factors among Iranian women. *J Obstet Gynaecol Res* 2011;37(5):442–51.
- [18] Abdul-Razzak KK, Ayoub NM, Abu-Taleb AA, Obeidat BA. The influence of dietary intake of dairy products on dysmenorrhea. *J Obstet Gynaecol Res* 2010;36(2):377–83.
- [19] Abdul-Razzak KK, Obeidat BA, Al-Farras MI, Dauod AS. Vitamin D and PTH status among adolescent and young females with severe dysmenorrhea. *J Pediatr Adolesc Gynecol* 2014;27(2):78–82.
- [20] Wray S, Burdyga T, Noble D, Noble K, Borysova L, Arrowsmith S. Progress in understanding electro-mechanical signalling in the myometrium. *Acta Physiol (Oxf)* 2015;213(2):417–31.
- [21] Noble K, Matthew A, Burdyga T, Wray S. A review of recent insights into the role of the sarcoplasmic reticulum and Ca entry in uterine smooth muscle. *Eur J Obstet Gynecol Reprod Biol* 2009;144(Suppl. 1):S11–9.
- [22] Maimoun L, Sultan C. Effect of physical activity on calcium homeostasis and calcitropic hormones: a review. *Calcif Tissue Int* 2009;85(4):277–86.
- [23] Holick MF. Vitamin D: a D-Lightful health perspective. *Nutr Rev* 2008;66(10 Suppl. 2):S182–94.
- [24] Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *Am J Clin Nutr* 2008;87(4):1080S–6S.
- [25] Saggese G, Vierucci F, Boot AM, Czech-Kowalska J, Weber G, Camargo Jr CA, et al. Vitamin D in childhood and adolescence: an expert position statement. *Eur J Pediatr* 2015;174(5):565–76.
- [26] Winzenberg T, Jones G. Vitamin D and bone health in childhood and adolescence. *Calcif Tissue Int* 2013;92(2):140–50.
- [27] Viganò P, Lattuada D, Mangioni S, Ermellino L, Vignali M, Caporizzo E, et al. Cycling and early pregnant endometrium as a site of regulated expression of the vitamin D system. *J Mol Endocrinol* 2006;36(3):415–24.
- [28] Adams JS, Hewison M. Extrarenal expression of the 25-hydroxyvitamin D-1-hydroxylase. *Arch Biochem Biophys* 2012;523(1):95–102.
- [29] Halicioğlu O, Aksit S, Koc F, Akman SA, Albudak E, Yaprak I, et al. Vitamin D deficiency in pregnant women and their neonates in spring time in western Turkey. *Paediatr Perinat Epidemiol* 2012;26(1):53–60.
- [30] Ergur AT, Berberoglu M, Atasay B, Siklar Z, Bilir P, Arsan S, et al. Vitamin D deficiency in Turkish mothers and their neonates and in women of reproductive age. *J Clin Pediatr Endocrinol* 2009;1(6):266–9.
- [31] Ates S, Sevket O, Ozcan P, Ozkal F, Kaya MO, Dane B. Vitamin D status in the first-trimester: effects of Vitamin D deficiency on pregnancy outcomes. *Afr Health Sci* 2016;16(1):36–43.
- [32] Obeidat BA, Alchalabi HA, Abdul-Razzak KK, Al-Farras MI. Premenstrual symptoms in dysmenorrhoeic college students: prevalence and relation to vitamin D and parathyroid hormone levels. *Int J Environ Res Publ Health* 2012;9(11):4210–22.
- [33] Parlak M, Kalay S, Kalay Z, Kirecci A, Guney O, Koklu E. Severe vitamin D deficiency among pregnant women and their newborns in Turkey. *J Matern Fetal Neonatal Med* 2015;28(5):548–51.
- [34] Pehlivan I, Hatun S, Aydogan M, Babaoglu K, Gokalp AS. Maternal vitamin D deficiency and vitamin D supplementation in healthy infants. *Turk J Pediatr* 2003;45(4):315–20.
- [35] Olmez D, Bober E, Buyukgebiz A, Cimrin D. The frequency of vitamin D insufficiency in healthy female adolescents. *Acta Paediatr* 2006;95(10):1266–9.
- [36] Whelan AM, Jurgens TM, Naylor H. Herbs, vitamins and minerals in the treatment of premenstrual syndrome: a systematic review. *Can J Clin Pharmacol* 2009;16(3):e407–29.
- [37] Thys-Jacobs S, McMahon D, Bilezikian JP. Cyclical changes in calcium metabolism across the menstrual cycle in women with premenstrual dysphoric disorder. *J Clin Endocrinol Metab* 2007;92(8):2952–9.
- [38] Bertone-Johnson ER. Vitamin D and the occurrence of depression: causal association or circumstantial evidence? *Nutr Rev* 2009;67(8):481–92.
- [39] Bertone-Johnson ER, Hankinson SE, Bendich A, Johnson SR, Willett WC, Manson JE. Calcium and vitamin D intake and risk of incident premenstrual syndrome. *Arch Intern Med* 2005;165(11):1246–52.
- [40] Ghanbari Z, Haghollahi F, Shariat M, Foroshani AR, Ashrafi M. Effects of calcium supplement therapy in women with premenstrual syndrome. *Taiwan J Obstet Gynecol* 2009;48(2):124–9.
- [41] Bertone-Johnson ER, Chocano-Bedoya PO, Zagarins SE, Micka AE, Ronnenberg AG. Dietary vitamin D intake, 25-hydroxyvitamin D3 levels and premenstrual syndrome in a college-aged population. *J Steroid Biochem Mol Biol* 2010;121(1–2):434–7.
- [42] Khajehei M, Abdali K, Parsanezhad ME, Tabatabaee HR. Effect of treatment with dydrogesterone or calcium plus vitamin D on the severity of premenstrual syndrome. *J Gynecol Obstet* 2009;105(2):158–61.
- [43] Lasco A, Catalano A, Benvenega S. Improvement of primary dysmenorrhoea caused by a single oral dose of vitamin D: results of a randomized, double-blind, placebo-controlled study. *Arch Intern Med* 2012;172(4):366–7.