



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Case Report

Detection of fetal trisomy 9 mosaicism by noninvasive prenatal testing through maternal plasma DNA sequencing



Chung-Yuan Lee^{a, b, c}, Hsing-Ju Su^b, Yu-Tzu Cheng^b, Yu-Lun Ku^b, Yeh Giin Ngo^b,
Che-Min Chen^b, Yu-Che Ou^b, Meng-Chih Lee^a, Sheng-Wen Steven Shaw^{d, e, *}

^a Institute of Medicine, Chung Shan Medical University, Taichung, Taiwan^b Department of Obstetrics and Gynecology, Chia-Yi Chang Gung Memorial Hospital, Chia-Yi, Taiwan^c Department of Nursing, Chang Gung University of Science and Technology, Chia-Yi Campus, Chia-Yi, Taiwan^d Department of Obstetrics and Gynecology, Taipei Chang Gung Memorial Hospital, Taipei, Taiwan^e College of Medicine, Chang Gung University, Taoyuan, Taiwan

ARTICLE INFO

Article history:

Accepted 3 April 2018

Keywords:

Amniocentesis

Karyotyping

Maternal plasma DNA

Noninvasive prenatal testing

Trisomy 9 mosaicism

ABSTRACT

Objective: Noninvasive prenatal testing (NIPT) is widely used as a powerful screening tool to detect common aneuploidies. However, its application for detection of rare chromosomal abnormalities remains inconclusive.

Case report: A 38-year-old woman (gravida 2, para 0) requested NIPT as a primary screening test for fetal aneuploidies at 13 weeks and 1 day of gestation. An unexpected Trisomy 9 (T9) abnormality was highly suspected. Amniocentesis was arranged for further diagnosis at 18 weeks of gestation. Final karyotyping reported 47,XX,+9 [18]/46,XX [12], indicating 60% T9 mosaicism.

Conclusion: This case shows strong evidence that NIPT can be a powerful screening tool to detect rare fetal trisomies at very early gestation.

© 2018 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Background

Most pregnant women without indication for chorionic villous sampling or amniocentesis receiving prenatal screening for fetal aneuploidies rely on measurement of multiple biochemical markers in maternal serum as well as ultrasound examination of fetal nuchal translucency or biparietal diameter in the first and second trimesters. If fetal chromosomal abnormalities are suspected, chorionic villous sampling or amniocentesis remain the gold standard for prenatal diagnosis and carry little critical risks, such as miscarriage, abortion, and intrauterine infection [1]. Recently, noninvasive prenatal testing (NIPT) by massively parallel sequencing of cell-free DNA in maternal circulation, which contains an average of 10%–20% fetal DNA during the second trimester, has been applied for prenatal aneuploidy screening [2]. The benefit of NIPT is the safe and accurate detection of fetal aneuploidies at early gestation. Based on several large prospective trials [3–6], the

general detection rate of trisomy (T) 21, T18, and T13 is >99%, while the false positive and negative rates are 0.1%–0.2%, if the maternal plasma-free fetal DNA fraction is adequate.

According to positive NIPT results, it is strongly suggested to confirm the results by invasive prenatal diagnosis methods, because the fetal DNA tested originated from placental trophoblasts [7]. Nonetheless, NIPT can detect common fetal autosomal aneuploidies, such as T21, T18, and T13, and sex chromosomal aneuploidies, while several other studies have reported the technical potential of NIPT for detecting other chromosomal aneuploidies, mosaicism, and small copy number variations [4–6,8,9]. In addition to identification of these rare genetic conditions, NIPT may also have clinical utility for detection of rare fetal trisomies. Here, we present a rare case of fetal T9 mosaicism that was originally detected by NIPT and confirmed by traditional invasive prenatal diagnosis methods. We also compare this case with other reported cases of T9 mosaicism.

Case report

The patient was a 38-year-old woman (gravida 2, para 0). She had no family history of chromosomal abnormalities or congenital

* Corresponding author. Department of Obstetrics and Gynecology, Taipei Chang Gung Memorial Hospital, No. 199, Dun-Hua North Road, Taipei, 105, Taiwan. Fax: +886 3 3288252.

E-mail address: dr.shaw@me.com (S.-W.S. Shaw).



Fig. 1. Karyotyping analysis of amniotic fluid cells. Final karyotyping reported 47,XX,+9 [18]/46,XX [12], indicating 60% trisomy 9 mosaicism (red circle).

fetal malformations. There was no sign of spontaneous abortion during early pregnancy. All prenatal laboratory data were within normal range. Because of her advanced maternal age and hesitation for the unavoidable risks of invasive prenatal diagnosis methods, she selected NIPT as a primary screening test for fetal autosomal aneuploidies at 13 weeks and 1 day of gestation after counseling.

Sample preparation, maternal plasma DNA sequencing, and bioinformatics analysis were performed as previously described [10]. Briefly, 5 mL of maternal blood was collected in a Streck tube for sample preparation within 72 h. Cell-free DNA extraction, library construction, and massively parallel sequencing were performed at ISO17025-certified clinical laboratories using BGISEQ-500 platforms (BGI, China). Bioinformatics analysis was carried out using the proprietary algorithm previously reported [10], which

uses the binary hypothesis T-score to classify a high-risk sample (T-score > 3 or < -3) or low-risk sample (T-score > -3 or < 3). Assessment of all 23 pairs of chromosomes was included in the experimental protocol.

Under the qualification of 7.87% cell-free fetal DNA fraction (3.5% is the least reliable cell-free fetal DNA level), NIPT results showed low risk of T21 (T-score = -1.35), T18 (T-score = -0.59), T13 (T-score = -0.51), T6 (T-score = -1.89), and T22 (T-score = 1.19). Sex chromosomes, including monosomy X (X0), XXY, XXX, and XYY, were also within normal limits. Other deletion syndromes, such as Cri du chat syndrome (5p15deletion), 1p36 deletion syndrome, 2q33.1 deletion syndrome, 16p12.2-p11.2 deletion syndrome, type II DiGeorge syndrome (10p14-p13 deletion), Jacobsen syndrome (11q23 deletion), Prader-Willi/Angelman syndrome (15q11.2 deletion), and Van der Woude syndrome (1q32.2 deletion), were also considered low risk. Aneuploidy of other chromosomes or chromosomal deletions/duplications were not detected. An unexpected T9 abnormality (T-score = 7.16) was highly suspected.

After counseling for the NIPT results, the patient agreed to undergo amniocentesis for further analysis at 18 weeks of gestation. Final karyotyping reported 47,XX,+9 [18]/46,XX [12], indicating 60% T9 mosaicism (Fig. 1). Subsequent confirmation fetal ultrasound showed bilateral low-set ears, micrognathia, clitoromegaly, and partial depletion of the corpus callosum (Fig. 2).

After a complete prenatal survey, a final consultation was given and the couple decided to terminate the pregnancy. Medical termination was performed at 20 weeks and 2 days of gestation. Gross examination of the abortus showed female sex, bilateral low-set ears, micrognathia, and clitoromegaly, compatible with the previous ultrasound (Fig. 3).

Discussion

In 1973, Haslam and colleagues reported the first case of T9 mosaicism, while Feingold and colleagues reported the first example of a child with full T9 using blood lymphocytes in the same




Fig. 2. The arrow denotes the site of fetal ultrasound showing bilateral low-set ears (A), micrognathia (B), clitoromegaly (C), and partial depletion of the corpus callosum (D).




Fig. 3. The arrow denotes the site of abortus showing bilateral low-set ears and micrognathia (A), and female sex and clitoromegaly (B), similar to the previous ultrasound images.

Table 1

Characteristics of patients with trisomy 9 mosaicism.

Case	Maternal age, y	G/P Screening			Diagnosis			Fetus			Author	
		Method	GA, weeks	Result	Method	GA, weeks	Result	Outcome	GA, weeks	Presentation		
1	39	G4/ P1	NIPT	17	T9 high risk	Amniocentesis	20	47,XX,+9 [55]/46,XX [33], T9 mosaicism (62.5%)	Termination	26	Abnormal fetal head morphology, congenital absence of parietal bone o/n both sides	Sun et al. [14]
2	NA	NA	NIPT	16	T9 mosaicism (30.1%)	Percutaneous umbilical blood sampling	34	46,XX	Alive	41 (vaginal delivery; birth weight, 1960 g)	Congenital cerebral dysplasia, intracranial hemorrhage, thrombocytopenia, congenital heart disease (ventricular septal defect, patent ductus arteriosus)	Ma et al. [15]
3	38	G2/ P0	NIPT	13	T9 high risk	Amniocentesis	18	47,XX,+9 [18]/46, XX [12], T9 mosaicism (60%)	Termination	20	Low-set ears, micrognathia, clitoromegaly, partial depletion of corpus callosum	Present case

G, gravida; GA, gestational age; NA, not available; NIPT, noninvasive prenatal testing; P, para.

year [11,12]. The etiology of this disorder is trisomy of chromosome 9. The incidence and severity of malformations and intellectual disability correlate with the percentage of trisomic cells in different tissues [13]. The majority of T9 mosaicism cases die during the early postnatal period. If the patient survives, most cases present with symptoms of failure to thrive and severe motor and intellectual disabilities [13]. The usual abnormalities of T9 mosaicism include prenatal onset of growth deficiency, severe intellectual disability, and structural deformity of the skeletal and craniofacial area [13]; approximately two-thirds of cases also have congenital heart defects [13].

T9 is relatively rare compared with T21, T18, and T13. In the past 2 decades, young pregnant women without indication for invasive prenatal diagnosis methods can undergo noninvasive prenatal screening using biochemical and ultrasound markers. However, measurement of serum biomarkers typically requires a relatively narrow gestational age window, while ultrasound examination requires extensive experience and cannot reveal genetic defects [1]. Recently, NIPT by maternal plasma DNA sequencing has been demonstrated to be a safe and effective method to detect T21 (sensitivity, 100%; 95% confidence interval [CI], 95.9–100), T18 (sensitivity, 97.2%; 95% CI, 85.5–99.9), and T13 (sensitivity, 78.6%; 95% CI, 49.2–95.3) [4]. However, the application of NIPT for detection of T9 remains inconclusive, and T9 cases detected by NIPT are sporadic in the literature. We compared our case with 2 other published articles (Table 1) [14,15]. In comparison, our case is the earliest gestational age to detect T9 by NIPT and with confirmation of T9 mosaicism by subsequent amniocentesis. Ultrasound was also performed for fetal

structural abnormality. Therefore, our present case could provide useful evidence that NIPT is a powerful tool to detect T9 at very early gestation based on an adequate cell-free fetal DNA fraction.

Conclusion

In conclusion, we reported a case of fetal T9 mosaicism identified by NIPT at very early gestation. This case may provide strong evidence that NIPT is a powerful screening tool for detection of rare fetal trisomies at early gestation under an adequate cell-free fetal DNA fraction.

Conflict of interests

None.

Ethics approval

Institutional review board approval was obtained in advance for this study.

Author contribution

Conception and design of study: C. Y. Lee, S. W. Steven Shaw; Acquisition of data: H. J. Su, Y. T. Cheng, Y. L. Ku, Y. G. Ngo, C. M. Chen;

Analysis and/or interpretation of data: C. Y. Lee, S. W. Steven Shaw;

Drafting the manuscript: C. Y. Lee, H. J. Su;

Revising the manuscript critically for important intellectual content:

Y. C. Ou, S. W. Steven Shaw;

Approval of the version of the manuscript to be published: C. Y. Lee, H. J. Su, Y. T. Cheng, Y. L. Ku., Y. G. Ngo, C. M. Chen, M.C. Lee, Y.C. Ou, S.W. Steven Shaw.

Acknowledgment

The authors would like to thank Gene Health Co.Ltd. for offering a high-quality NIPT service, NIFTY PLUS. We are also grateful to the Chang Gung Medical Foundation Institutional Review Board who approved the study.

References

- [1] Tabor A, Alfirevic Z. Update on procedure-related risks for prenatal diagnosis techniques. *Fetal Diagn Ther* 2010;27:1–7.
- [2] Benn P, Cuckle H, Pergament E. Non-invasive prenatal testing for aneuploidy: current status and future prospects. *Ultrasound Obstet Gynecol* 2013;42:15–33.
- [3] Bianchi DW, Parker RL, Wentworth J, Madankumar R, Saffer C, Das AF, et al. DNA sequencing versus standard prenatal aneuploidy screening. *N Engl J Med* 2014;370:799–808.
- [4] Bianchi DW, Platt LD, Goldberg JD, Abuhamad AZ, Sehnert AJ, Rava RP, et al. Genome-wide fetal aneuploidy detection by maternal plasma DNA sequencing. *Obstet Gynecol* 2012;119:890–901.
- [5] Dan S, Wang W, Ren J, Li Y, Hu H, Xu Z, et al. Clinical application of massively parallel sequencing-based prenatal noninvasive fetal trisomy test for trisomies 21 and 18 in 11,105 pregnancies with mixed risk factors. *Prenat Diagn* 2012;32:1225–32.
- [6] Liang D, Lv W, Wang H, Xu L, Liu J, Li H, et al. Non-invasive prenatal testing of fetal whole chromosome aneuploidy by massively parallel sequencing. *Prenat Diagn* 2013;33:409–15.
- [7] American College of Obstetricians and Gynecologists Committee on Genetics. Committee opinion No. 545: noninvasive prenatal testing for fetal aneuploidy. *Obstet Gynecol* 2012;120:1532–4.
- [8] Peters D, Chu T, Yatsenko SA, Hendrix N, Hogge WA, Surti U, et al. Noninvasive prenatal diagnosis of a fetal microdeletion syndrome. *N Engl J Med* 2011;365:1847–8.
- [9] Chen CP, Tsai C, Lin MH, Chern SR, Chen SW, Lai ST, et al. Application of non-invasive prenatal testing in late gestation in a pregnancy associated with intrauterine growth restriction and trisomy 22 confined placental mosaicism. *Taiwan J Obstet Gynecol* 2017;56:691–3.
- [10] Jiang F, Ren J, Chen F, Zhou Y, Xie J, Dan S, et al. Noninvasive fetal trisomy (NIFTY) test: an advanced noninvasive prenatal diagnosis methodology for fetal autosomal and sex chromosomal aneuploidies. *BMC Med Genom* 2012;5:57.
- [11] Feingold M, Atkins L. A case of trisomy 9. *J Med Genet* 1973;10:184–7.
- [12] Haslam RH, Broske SP, Moore CM, Thomas GH, Neill CA. Trisomy 9 mosaicism with multiple congenital anomalies. *J Med Genet* 1973;10:180–4.
- [13] Jones KL, Jones MC, Del Campo M. Smith's recognizable patterns of human malformation. 7th ed. New York: Elsevier; 2013.
- [14] Sun L, Zhang L, Zhou J, Yang X, Duan T, Wang W. Noninvasive prenatal testing for fetal trisomy 9 mosaicism by maternal plasma DNA sequencing. *Case Rep Perinat Med* 2015;4:37–40.
- [15] Ma J, Cram DS, Zhang J, Shang L, Yang H, Pan H. Birth of a child with trisomy 9 mosaicism syndrome associated with paternal isodisomy 9: case of positive noninvasive prenatal test result unconfirmed by invasive prenatal diagnosis. *Mol Cytogenet* 2015;8:44.