



Contents lists available at ScienceDirect

Taiwanese Journal of Obstetrics & Gynecology

journal homepage: www.tjog-online.com

Case Report

Prenatal diagnosis and molecular cytogenetic characterization of a small supernumerary marker chromosome derived from chromosome 15 in a pregnancy associated with recurrent Down syndrome

Chih-Ping Chen^{a, b, c, d, e, f, *}, Chia-Hao Chan^g, Schu-Rern Chern^b, Peih-Shan Wu^h, Shin-Wen Chen^a, Fang-Tzu Wu^a, Dai-Dyi Town^a, Meng-Shan Lee^a, Wayseen Wang^b^a Department of Obstetrics and Gynecology, MacKay Memorial Hospital, Taipei, Taiwan^b Department of Medical Research, MacKay Memorial Hospital, Taipei, Taiwan^c School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan^d Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan^e Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan^f Department of Medical Laboratory Science and Biotechnology, Asia University, Taichung, Taiwan^g Department of Obstetrics and Gynecology, Hsinchu MacKay Memorial Hospital, Hsinchu, Taiwan^h Gene Biodesign Co. Ltd, Taipei, Taiwan

ARTICLE INFO

Article history:

Accepted 30 September 2020

Keywords:

Chromosome 15

Recurrent down syndrome

Small supernumerary marker chromosome

ABSTRACT

Objective: We present prenatal diagnosis and molecular cytogenetic characterization of a small supernumerary marker chromosome (sSMC) derived from chromosome 15 in a pregnancy associated with recurrent Down syndrome.

Case report: A 33-year-old, gravida 4, para 2, woman underwent amniocentesis at 16 weeks of gestation because of a previous child with Down syndrome and a karyotype of 46,XY,der(14;21)(q10;q10),+21. In this pregnancy, amniocentesis revealed a karyotype of 47,XX,+21[12]/48,XX,+21,+mar[3]. The parental karyotypes were normal. The pregnancy was terminated, and a malformed fetus was delivered with characteristic craniofacial appearance of Down syndrome and hypoplastic middle phalanx of the fifth fingers. The placenta had a karyotype of 47,XX,+21[37]/48,XX,+21,+mar[3]. The umbilical cord had a karyotype of 47,XX,+21[38]/48,XX,+21,+mar[2]. In addition to trisomy 21, array comparative genomic hybridization (aCGH) on the DNA extracted from umbilical cord revealed 40–50% mosaicism for a 2.604-Mb duplication of 15q25.2–q25.3, or arr 15q25.2q25.3 (83,229,665–85,834,131) × 2.4 [GRCh37 (hg19)] encompassing 19 Online Mendelian Inheritance in Man (OMIM) genes. Quantitative fluorescent polymerase chain reaction (QF-PCR) using the DNAs extracted from cultured amniocytes and parental bloods revealed maternal origin of the sSMC(15) and the extra chromosome 21.

Conclusion: aCGH is useful for identification of the nature of sSMC, and QF-PCR is useful for determination of the parental origin of the aberrant chromosomes.

© 2021 Taiwan Association of Obstetrics & Gynecology. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

We previously reported identification of a small supernumerary marker chromosome (sSMC) derived from chromosome 15 [1–3]. Here, we present prenatal diagnosis and molecular cytogenetic

characterization of an sSMC derived from chromosome 15 in a pregnancy associated with recurrent Down syndrome.

Case report

A 33-year-old, gravida 4, para 2, woman underwent amniocentesis at 16 weeks of gestation because of a previous child with Down syndrome and a karyotype of 46,XY,der(14;21)(q10;q10),+21. In this pregnancy, amniocentesis revealed a karyotype of 47,XX,+21 [12]/48,XX,+21,+mar [3] (Fig. 1). The parental karyotypes were normal. The pregnancy was terminated, and a malformed fetus was

* Corresponding author. Department of Obstetrics and Gynecology, MacKay Memorial Hospital 92, Section 2, Chung-Shan North Road, Taipei 104215, Taiwan. Fax: +886 2 25433642, +886 2 25232448.

E-mail address: cpc_mmh@yahoo.com (C.-P. Chen).



Fig. 1. A karyotype of 48,XX,+21,+mar. mar = marker chromosome.

delivered with characteristic craniofacial appearance of Down syndrome and hypoplastic middle phalanx of the fifth fingers. The placenta had a karyotype of 47,XX,+21[37]/48,XX,+21,+mar [3]. The umbilical cord had a karyotype of 47,XX,+21[38]/48,XX,+21,+mar [2]. In addition to trisomy 21, array comparative genomic hybridization (aCGH) on the DNA extracted from umbilical cord revealed 40–50% mosaicism for a 2.604-Mb duplication of 15q25.2–q25.3, or arr 15q25.2q25.3 (83,229,665–85,834,131) \times 2.4 [GRCh37 (hg19)] encompassing 19 Online Mendelian Inheritance in Man (OMIM) genes of *CPEB1*, *AP3B2*, *SCARNA15*, *WHAMM*, *HOMER2*, *FAM103A1*, *BTBD1*, *TM6SF1*, *HDGFRP3*, *BNC1*, *SH3GL3*, *ADAMTSL3*, *WDR73*, *NMB*, *SEC11A*, *ZNF592*, *ALPK3*, *SLC28A1* and *PDE8A* (Fig. 2). Quantitative fluorescent polymerase chain reaction (QF-PCR) using the DNAs extracted from cultured amniocytes and parental bloods revealed maternal origin of the sSMC(15) and the extra chromosome 21 (Fig. 3).

Discussion

The present case had a 2.604-Mb gene dosage increase on 15q25.2–q25.3 encompassing 19 OMIM genes including *HOMER2*, *BNC1*, *AP3B2*, *WDR73*, *ALPK3*, *SLC28A1* and *HDGFRP3*. *HOMER2* (OMIM 604799) is associated with autosomal dominant deafness 68 (OMIM 616707). *BNC1* (OMIM 601930) is associated with autosomal dominant premature ovarian failure 16 (OMIM 618723). *AP3B2* (OMIM 602166) is associated with autosomal recessive early infantile epileptic encephalopathy 48 (OMIM 617276). *WDR73* (OMIM 616144) is associated with autosomal recessive Galloway-Mowat

syndrome 1 (OMIM 251300). *ALPK3* (OMIM 617608) is associated with autosomal recessive familial hypertrophic cardiomyopathy 27 (OMIM 618052). *SLC28A1* (OMIM 606207) is associated with autosomal recessive uridine-cytidineuria (OMIM 618477). *HDGFRP3* (OMIM 616643) encodes hepatoma-derived growth factor-related protein 3 which is a novel angiogenic factor [4] and is involved in the progression of hepatocellular carcinoma [5]. *HDGFRP3* is a mitogenic, angiogenic and anti-apoptotic factor, and is also a neurotrophic and a neuroprotective factor that is related to proliferation, differentiation and maintenance of neurons [6].

Microdeletions and microduplications of 15q25.2 have been reported to be associated with phenotypic abnormalities. Monzani et al. [7] reported co-occurrence of duplication on Xp22.1 flanking the *SHOX* coding sequence and an additional duplication of 1.6–2.5 Mb on 15q25.2 that included 13 genes such as *RPS17*, *CPEB1* and *HOMER2* in a girl with short stature, precocious puberty, urogenital malformations and bone anomalies. Wat et al. [8] reported individual with recurrent microdeletions of 15q25.2 involving *CPEB1*, *AP3B2*, *HOMER2* and *HDGFRP3* in association with abnormal central nervous system development, cognitive deficits, congenital diaphragmatic hernia and anaemia. Monkam et al. [9] reported a case with a 15q25.2 microduplication, delayed speech, intellectual disability, splay foot, toe syndactyly, thin hand and short fifth fingers. DECIPHER database v10.1 [10] reported four patients with 15q25.2 microduplication with abnormal phenotype, i.e., #316713 with abnormality of toe, delayed speech, eczema and retrognathia; #289343 with intellectual disability and macrocephaly; #260289 with global developmental delay, nasal speech and seizure; and




Fig. 2. (A), (B) and (C) Array comparative genomic hybridization analysis by SurePrint G3 Unrestricted CGH ISCA v2, 8 × 60 K Array (Agilent Technologies, Santa Clara, CA, USA) using the DNA extracted from umbilical cord shows the result of arr 15q25.2q25.3 (83,229,665–85,834,131) × 2.4, arr 21q11.2q22.3 (15,485,008–48,090,317) × 3.0 [GRCh37 (hg19)].




Fig. 3. Quantitative fluorescent polymerase chain reaction assays on the DNAs extracted from cultured amniocytes and parental bloods using the informative markers of D15S643 (15q22.2), D15S821 (15q25.2), D15S230 (15q26.2), D21S1437 (21q21.1) and D21S2052 (21q21.3) show that D15S821 (15q25.2) has gene dosage increase in the maternal allele of 265 bp, D21S2052 (21q21.3) has gene dosage increase in the maternal allele of 137 bp, and D21S1437 (21q21.1) contains three different alleles.

#263566 with delayed speech, macrocephaly, micrognathia and preaxial hand polydactyly. However, in our case, no other abnormalities in addition to Down syndrome have been observed.

The present case was associated with recurrent Down syndrome in two pregnancies. However, both parents had normal phenotypes by conventional cytogenetic analysis on peripheral bloods. The recurrent risk for Down syndrome in phenotypically normal young parents is about 1–2% [11]. The recurrent risk has been proposed to be caused by parental mosaic trisomy 21, parental gonadal mosaicism, maternal age-associated risk and genetic predisposition to non-disjunction [12,13]. Magalhães et al. [14] reported a woman with three trisomy 21 pregnancies. The woman and her husband had normal karyotypes by conventional cytogenetic analysis on peripheral bloods. However, interphase fluorescence *in situ* hybridization (FISH) analysis on 230 maternal blood cells detected one cell with trisomy 21, and FISH analysis on maternal oral mucosal cells detected three trisomy-21 cells in 100 cells. Hultén et al. [15] suggested the importance of detection of germinal and somatic trisomy 21 mosaicism in individual carrier parents. Down syndrome associated with parental mosaic trisomy 21 has been well reported previously [16–22]. The incidence of parental mosaic trisomy 21 in families with only one child with Down syndrome is about 2.7–4.3% [17]. Harris et al. [18] estimated that 3% of couples producing a child with trisomy 21 can be explained by parental mosaicism. However, the incidence of parental mosaic trisomy 21 has been reported to be higher (5/13) in families with recurrent trisomy 21 [21].

In summary, we present prenatal diagnosis and molecular cytogenetic characterization of an sSMC derived from chromosome 15 in a pregnancy associated with recurrent Down syndrome. aCGH is useful for identification of the nature of sSMC, and QF-PCR is useful for determination of the parental origin of the aberrant chromosomes.

Declaration of competing interest

The authors have no conflicts of interest relevant to this article.

Acknowledgements

This work was supported by research grants MOST-107-2314-B-195-005 from the Ministry of Science and Technology, Taiwan, and MMH-E-109-04 from Mackay Memorial Hospital, Taipei, Taiwan.

References

- [1] Chen C-P, Chen M, Su Y-N, Chern S-R, Wu P-S, Chang S-P, et al. Prenatal diagnosis and molecular cytogenetic characterization of mosaicism for a small supernumerary marker chromosome derived from chromosome 15. *Taiwan J Obstet Gynecol* 2014;53:129–32.
- [2] Chen C-P, Lin S-P, Chern S-R, Wu P-S, Chen Y-N, Chen S-W, et al. Molecular cytogenetic characterization of an inv dup(15) chromosome presenting as a small supernumerary marker chromosome associated with the inv dup(15) syndrome. *Taiwan J Obstet Gynecol* 2016;55:728–32.
- [3] Chen C-P, Lin H-Y, Wang L-K, Chern S-R, Wu P-S, Chen S-W, et al. Prenatal diagnosis and molecular cytogenetic characterization of a small supernumerary marker chromosome derived from inv dup(15). *Taiwan J Obstet Gynecol* 2020;59:580–5.
- [4] LeBlanc ME, Wang W, Caberoy NB, Chen X, Guo F, Alvarado G, et al. Hepatoma-derived growth factor-related protein-3 is a novel angiogenic factor. *PLoS One* 2015;10:e0127904.
- [5] Enomoto H, Nakamura H, Liu W, Nishiguchi S. Hepatoma-derived growth factor: its possible involvement in the progression of hepatocellular carcinoma. *Int J Mol Sci* 2015;16:14086–97.
- [6] Eltahir HM. Insight into the roles of hepatoma derived growth factor related protein-3 under physiological and pathological conditions. *Biochem Cell Biol* 2018;96:707–12.
- [7] Monzani A, Babu D, Mellone S, Genoni G, Fanelli A, Prodam F, et al. Co-occurrence of genomic imbalances on Xp22.1 in the *SHOX* region and 15q25.2 in a girl with short stature, precocious puberty, urogenital malformations and bone anomalies. *BMC Med Genom* 2019;12:5.
- [8] Wat MJ, Enciso VB, Wiszniewski W, Resnick T, Bader P, Roeder ER, et al. Recurrent microdeletions of 15q25.2 are associated with increased risk of congenital diaphragmatic hernia, cognitive deficits and possibly Diamond-Blackfan anaemia. *J Med Genet* 2010;47:777–81.
- [9] Monkam CY, Kemeny S, Miret A, Pebrel-Richard C, Sarret C. A case which further refines the critical region for 15q25.2 microduplication phenotypes. *Acta Neurol Belg* 2016;116:683–5.
- [10] DECIPHER database: Database of genomic variation and Phenotype in Humans using Ensembl Resources. Available at: <https://decipher.sanger.ac.uk/>. [Accessed 5 August 2020].
- [11] Kovaleva NV. Germ-line transmission of trisomy 21: data from 80 families suggest an implication of grandmaternal age and a high frequency of female-specific trisomy rescue. *Mol Cytogenet* 2010;3:7.

- [12] Bruyère H, Rupps R, Kuchinka BD, Friedman JM, Robinson WP. Recurrent trisomy 21 in a couple with a child presenting trisomy 21 mosaicism and maternal uniparental disomy for chromosome 21 in the euploid cell line. *Am J Med Genet* 2000;94:35–41.
- [13] Warburton D, Dallaire L, Thangavelu M, Ross L, Levin B, Kline J. Trisomy recurrence: a reconsideration based on North American data. *Am J Hum Genet* 2004;75:376–85.
- [14] Magalhães M, Marques C, Ramos F, Jardim A, Franco S, Coelho F, et al. Why could a woman have three Trisomy 21 pregnancies? - a case report. *Clin Case Rep* 2017;5:1222–5.
- [15] Hultén MA, Öijerstedt L, Iwarsson E, Jonasson J. Maternal germinal trisomy 21 in Down syndrome. *J Clin Med* 2014;3:167–75.
- [16] Sachs ES, Jahoda MG, Los FJ, Pijpers L, Wladimiroff JW. Trisomy 21 mosaicism in gonads with unexpectedly high recurrence risks. *Am J Med Genet* 1990;S7: 186–8.
- [17] Hsu LYF, Gertner M, Leiter E, Hirschhorn K. Paternal trisomy 21 mosaicism and Down's syndrome. *Am J Hum Genet* 1971;23:592–601.
- [18] Harris DJ, Begleiter ML, Chamberlin J, Hankins L, Magenis RE. Parental trisomy 21 mosaicism. *Am J Hum Genet* 1982;34:125–33.
- [19] Uchida IA, Freeman VC. Trisomy 21 Down syndrome. Parental mosaicism. *Hum Genet* 1985;70:246–8.
- [20] Nielsen KG, Poulsen H, Mikkelsen M, Steuber E. Multiple recurrence of trisomy 21 Down syndrome. *Hum Genet* 1988;78:103–5.
- [21] Pangalos CG, Talbot Jr CC, Lewis JG, Adelsberger PA, Petersen MB, Serre JL, et al. DNA polymorphism analysis in families with recurrence of free trisomy 21. *Am J Hum Genet* 1992;51:1015–27.
- [22] Tseng L-H, Chuang S-M, Lee T-Y, Ko T-M. Recurrent Down's syndrome due to maternal ovarian trisomy 21 mosaicism. *Arch Gynecol Obstet* 1994;255: 213–6.